






Geographic range extension of the Spix's Disk-winged Bat, *Thyroptera tricolor* Spix, 1823 (Chiroptera, Thyropteridae)—first record from Santa Catarina, southern Brazil

FERNANDO CARVALHO^{1,2}, KAROLAINE P. SUPI^{1,2}, BEATRIZ F. L. LUCIANO^{1,2}, ARTUR S. SOBRINHO³, SAM GENEROSO², VICTOR QUINTAS², SÉRGIO L. ALTHOFF^{3,4}

1 Post-Graduate Program in Environmental Sciences, University of the Extremo Sul Catarinense, Criciúma, SC, Brazil • FC: f.carvalho@unesc.net  <https://orcid.org/0000-0002-7216-7083> • KPS: karolainesupi@outlook.com  <https://orcid.org/0000-0002-2667-3882> • BFL: beatrizfll@hotmail.com  <https://orcid.org/0000-0002-5715-9165>

2 Laboratory of Zoology and Ecology of Vertebrates, University of the Extremo Sul Catarinense, Criciúma, SC, Brazil • SG: samygeneroso@gmail.com  <https://orcid.org/0009-0007-4815-4390> • VQ: quintassc@gmail.com  <https://orcid.org/0009-0005-5778-0057>

3 Post-Graduate Program in Biodiversity, Regional University of Blumenau, Blumenau, SC, Brazil • ASS: assbio@gmail.com  <https://orcid.org/0000-0001-9015-529X> • SLA: slalthoff@gmail.com  <https://orcid.org/0000-0002-0877-5156>

4 Laboratory of Zoology, Regional University of Blumenau, Blumenau, SC, Brazil

* Corresponding author

Abstract. We captured the Spix's Disk-winged Bat, *Thyroptera tricolor* Spix, 1823, at two locations in Santa Catarina state, southern Brazil. These are the first records of occurrence of *T. tricolor* from the state and expand the southern limit of the species' geographical range by 408 km. With our new data, 52 species of bat are now confirmed to occur in Santa Catarina; this state has the second richest bat fauna in southern Brazil.

Keywords. Atlantic Forest, distribution, bats, mist nets, Parque das Nascentes, São Francisco Reserve

Academic editor: Héctor Ramírez-Chaves

Received 25 April 2023, accepted 25 September 2023, published 11 October 2023

Carvalho F, Supi KP, Luciano BFL, Sobrinho AS, Generoso S, Quintas V, Althoff SL (2023) Geographic range extension of the Spix's Disk-winged Bat, *Thyroptera tricolor* Spix, 1823 (Chiroptera, Thyropteridae)—first record from Santa Catarina, southern Brazil. Check List 19 (5): 719–725. <https://doi.org/10.15560/19.5.719>

Introduction

Species of the bat family Thyropteridae occur in moist, lowland, Neotropical forests from Mexico to southeastern Brazil (Lee 2019). Members of this family are easily distinguished from other Neotropical bats by the presence of adhesive disks on the soles of their feet and on their thumbs (Boada et al. 2010; Lee 2019). They are small bats with an average weight of 4 g and have funnel-shaped ears, a long snout, rounded nostrils, and small eyes (Findley and Wilson 1974; Wilson 2008; Chaverri and Vonhof 2011; Semedo et al. 2020). Thyropteridae is represented by a single genus, *Thyroptera* Spix, 1823, and five species, *Thyroptera tricolor* Spix, 1823, *Thyroptera discifera* (Lichtenstein & Peters, 1854),

Thyroptera lavalii Pine, 1993, *Thyroptera devivoi* Gregorin, Gonçalves, Lim & Engstrom 2006, and *Thyroptera wynneae* Velazco, Gregorin, Voss & Simmons, 2014 (Wilson 2008; Velazco et al. 2014), all with confirmed records in Brazil (Garbino et al. 2022). In the Atlantic Forest, only *T. wynneae* and *T. tricolor* have been confirmed (Hoppe et al. 2014; Solari 2015, 2016; Solari and Velazco 2016; Tavares and Mantilla 2016; Solari and Villada-Cadavid 2018).

Thyroptera tricolor is distributed from southern Mexico, through Central America, northern and western South America, to the northern region and the coast of Brazil (Wilson 2008; Tavares and Mantilla 2016; Lee 2019). It occurs at elevations up to 1,800 m above sea level (Reyes-Amaya et al. 2016). The southernmost dis-

tribution limit of this species is the southern Brazilian state of Paraná (Passos et al. 2010). *Thyroptera tricolor* is as a small-bodied species having an average weight of 4 g and a forearm length of 35.9 mm, and females are larger than males (Chaverri and Vonnhoff 2011). This species has a slow and maneuverable flight, and flies 3–5 m above the ground, which results in a low rate of capture in mist nets in the understory (Findley and Wilson 1974; Castaño-Salazar et al. 2004; Esbérard et al. 2007; Boada et al. 2010; Garbino 2016). Manual captures in roosts (Velazco et al. 2014) have shown that over 100 individuals are present in a single site (e.g. Vonnhoff and Fenton 2004; Chaverri 2010; Chaverri and Vonnhoff 2011; Sagot et al. 2022).

Due to its ability to adhere to and move on smooth surfaces, *T. tricolor* roosts in the young leaves of genera *Heliconia* L., *Musa* L., and occasionally *Calathea* G. Mey. and *Cecropia* Loefl. (Findley and Wilson 1974; Vonnhoff and Fenton 2004; Velazco et al. 2014). Bats of this species form cohesive groups in shelters with strong social relationships and composed of up to 11 individuals, which are possibly related; bats roost in rows, positioned with their heads facing upwards (Findley and Wilson 1974; Vonnhoff and Fenton 2004; Vonnhoff et al. 2004; Chaverri 2010).

Throughout the geographic range of *T. tricolor*, the species is usually recorded within protected areas (Dechmann et al. 2006; Passos et al. 2010; Chaverri 2010; Chaverri and Kunz 2011; Montero and Gillam 2015). However, it can also be found less frequently in degraded or human-modified landscapes, such as rubber-tree plantations and small forest fragments (Castaño-Salazar et al. 2004; Esbérard et al. 2007; Boada et al. 2010). Although sensitive to environmental changes, *T. tricolor* is currently classified as Least Concern according to IUCN Red List criteria (Tavares and Mantilla 2016), and it is not on the Official List of Endangered Species of the Brazilian Fauna (MMA 2022). Here, we present an geographic range extension of *T. tricolor* and the first record from Santa Catarina based on newly collected specimens at localities within the Atlantic Forest.

Methods

Our new records of *T. tricolor* were obtained at two localities in the state of Santa Catarina (Fig. 1), both in Dense Ombrophilous Forest inserted in the Atlantic Forest phytogeographic domain (IBGE 2012). According to the Köppen climate classification system, the predominant climate in both locations is the *Cfa* type, which is characterized as humid subtropical with hot summers and no defined dry season (Alvares et al. 2013).

The first location was in the Nascentes do Garcia Municipal Natural Park in the city of Blumenau, northern Santa Catarina (Fig. 1). The park has an area of approximately 53 km² and is within the area of the Serra do Itajaí National Park, and as a result, the park integrates a large and continuous forest remnant, which

extends beyond its boundary and forms an important ecological corridor between conservation units on the slopes of Serra Geral. The second location was in the São Francisco Natural Reserve in the city of Nova Veneza, in southernmost Santa Catarina (Fig. 1). Currently, the reserve has an area of approximately 15 km² and is connected to the Aguaí State Biological Reserve, an important protected area in Santa Catarina. On a macro scale, considering the landscape matrix, the two locations are connected by the Serra Geral ecological corridor.

In the Nascente do Garcia Municipal Park the bat was captured inside a house, located on the edges of a forest remnant. In São Francisco Natural Reserve the capture occurred during sampling for a research project on the bat fauna. In this area, sampling began in April 2022, with three sampling nights in each month. For each sampling night, 10 mist nets (two of 12 m × 2.6 m, four of 9 × 2.6 m, and four of 6 × 2.6 m) were installed and remained open for 6 h after sunset. The nets were placed on trails, forest edges, over bodies of water, and in banana plantations.

Two captured individuals were collected, fixed and preserved in 70% alcohol and had their skull prepared and deposited in the Zoology collection of the Regional University of Blumenau (CZFUB-SLA) and in the collection of the Zoology Laboratory of the University of the Extreme South of Santa Catarina (LABZEV). We measured one specimen (LABZEV 1007) using calipers (to the nearest 0.001 mm), measuring forearm length, condyloincisive length, maxillary tooththrow length, mandibular tooththrow length, length of mandible, greatest length of skull, braincase breadth, zygomatic breadth, width at M3, postorbital breadth, and rostral length.

Results

Materials examined. BRAZIL – Santa Catarina • Blumenau, Nascentes do Garcia Municipal Natural Park; 27°03'16"S, 049°05'07"W; 301 m alt.; 07.XI.2004; A.S. Sobrinho & S.L. Althoff leg.; inside a residence next to a forest remnant/manual capture; 1 ♂, CZFUB-SLA 2700 • Nova Veneza, São Francisco Reserve; 28°39'10"S, 049°37'52"W; 207 m alt.; 24.VIII.2022; F. Carvalho & V. Quintas leg.; using a polyester mist net (Ecotone®), installed in the midst of a banana plantation; 1 ♂, LABZEV 1007.

Identification. The specimens were identified based on morphological characters described by Miranda et al. (2011), Velazco et al. (2014), and Díaz et al. (2016); the following diagnostic morphological traits were observed: (1) circular disks on the thumbs; (2) forearm length 32–39mm; (3) white or yellowish ventral pelage which contrasts with the dorsal pelage; (4) proximal portion of the forearm sparsely haired; (5) dark ears; (6) calcar with two lappets posteriorly (Fig. 2, Table 1). The morphological data indicates that, of the 11 analyzed measurements, the individual from the São Francisco Reserve (LABZEV 1007) was larger in six




Figure 1. Distribution map of *Thyroptera tricolor* in Central and South America with addition of new records in state of Santa Catarina, where: Garcia Municipal Natural Park - PNMNG and, São Francisco Natural Reserve – RNSF.

Table 1. Measurements of the external and cranial morphological characters (in mm) of the individual captured in the São Francisco Reserve, in the south of Santa Catarina (LABZEV 1007), and those presented by Velazco et al. (2014), Passos et al. (2010) and Lee (2019).

Characters	Present study	Passos et al. (2010)	Velazco et al. (2014)		Lee (2019)*
	LABZEV 1007 ♂	♂	♀	♂	♂, ♀
Forearm length	35.98	34.35	37.10	36.70	—
Condylolincisive length	14.27	14.05	13.70	13.50	12.90–14.40
Maxillary tooththrow length	6.68	6.05	5.90	5.90	5.50–6.30
Mandibular tooththrow length	6.39	6.10	6.20	6.10	—
Length of mandible	11.04	10.50	10.60	10.40	—
Greatest length of skull	15.67	15.45	14.60	14.30	13.60–15.70
Braincase breadth	7.62	7.80	7.30	7.30	—
Zygomatic breadth	7.94	7.80	7.50	7.40	6.90–7.70
Width at M3	5.39	—	5.20	5.20	—
Postorbital breadth	2.98	2.85	2.70	2.70	—
Rostral length	5.45	—	5.30	—	5.30–5.50

*The range shown represents the minimums and maximums observed in Lee’s (2009) study.

measurements (mandibular and maxillary tooththrow length, length of mandible, zygomatic breadth, width at M3 and postorbital breadth: Table 1) than compared to those from other locations.

Discussion

The records that we provide here represent the first two confirmed localities of *T. tricolor* in Santa Catarina; this




Figure 2. *Thyroptera tricolor* captured in the São Francisco Reserve. **A.** Specimen. **B–D.** Skull, in lateral, dorsal, and ventral view, respectively. **E, F.** Mandible, in lateral view and dorsal view. Scale bars: 0.5 cm.

species had not been reported from the state until now (e.g. Cherem et al. 2004; Pacheco et al. 2007; Passos et al. 2010; Carvalho et al. 2013; Bôlla et al. 2017). Additionally, the São Francisco Reserve record extends the species' known southern distributional limit by 408 km from the Natural Patrimony Private Reserve Salto Morato, which is on the north coast of Paraná (Passos et al. 2010). Our new records of *T. tricolor* from Santa Catarina aid in understanding the distribution patterns of *T. tricolor* in the Atlantic Forest—the Atlantic Forest phytogeographic domain has few records of this species (Esbérard et al. 2007; Passos et al. 2010; Garbino 2016). This is due to the species' low rate of capture in mist nets installed in the understory (Castaño-Salazar et al. 2004; Velazco et al. 2014). In areas where other sampling methods were used, the number of captures of *T. tricolor* is higher, with more than 100 individuals detected (e.g. Chaverri 2010; Chaverri and Vonnhof 2011; Sagot et al. 2022). Thus, there are large gaps in knowledge about the distribution of *T. tricolor*, and new data, such as those presented here, have importance, especially for the southern portion of the Atlantic Forest.

Both locations where we found *T. tricolor* in Santa Catarina are protected areas, and these are part of larger, conserved forest remnant. The occurrence of this species in intact forest environments has already been reported in other countries (e.g. Findley and Wilson 1974; Chaverri 2010; Chaverri and Vonnhof 2011; Turcio-Casco and Medina-Fitoria 2018) and in Brazil (Sampaio et al. 2003; Bernard and Fenton 2007; Passos

et al. 2010; Tavares et al. 2017; Semedo et al. 2020). *Thyroptera tricolor* has an expressive dependence on plants suitable for roosting, such as those of the genera *Heliconia* and *Calathea* (Findley and Wilson, 1974; Vonnhof and Fenton 2004), which are characteristically found in the understory (Chaverri 2010; Velazco et al. 2014) of intact tropical forests. However, the Nascentes do Garcia Park individual was located inside a residence, while the São Francisco Reserve specimen was caught in a banana plantation.

In areas where preferred plants are absent, *T. tricolor* may use other plant species for shelter (Chaverri and Kunz 2011) or even roost in cavities (Esbérard et al. 2007) and man-made structures, as reported here. This suggests that the species may also be tolerant to moderately impacted or modified environments. However, in the Atlantic Forest, the environmental change has been great and may have drastically reduced the availability of shelter sites for *T. tricolor*. Bats that use ephemeral roosts, such as this species, tend to need multiple roost sites (Vonnhof and Fenton 2004). Thus, the loss of shelter sites may affect the occurrence of this species, as well as the behavior and social cohesion of groups (e.g. Chaverri 2010; Chaverri and Kunz 2011), as *T. tricolor* displays extremely well-developed social behavior (e.g. Vonnhof et al. 2004; Sagot et al. 2022).

The morphological measurements of the specimen captured in the São Francisco reserve are more similar to those reported by Passos et al. (2010) for an individual captured in southern Brazil than compared to

individuals captured at further south (Table 1). The reason for this is not known, and few individuals have been captured in these southern areas. However, it may be important for guiding further research.

With our new records of *T. tricolor*, 52 bat species are confirmed from Santa Catarina; at least six species have been added in the last 11 years (Passos et al. 2010; Carvalho and Fábian 2011; Cherem and Althoff 2015; Carvalho et al. 2017; Althoff et al. 2017, 2018). This is reflective of increased research groups and funding of science by government agencies, such as, for example, the Foundation for the Support of Research and Innovation of Santa Catarina (FAPESC). Despite the progress made in recent years, there is still much more to learn about the bat fauna of Santa Catarina, and the continuity of projects and funding is of fundamental importance for a better understanding of the patterns of diversity and occurrence of bat species in the state.

Acknowledgements

We acknowledge FAPESC for financial support to this project (2021TR1825) and the São Francisco Reserve for logistical support during our fieldwork. We thank Prof. Jori Ramos Pereira for preparing the species distribution map. We would like to thank the two reviewers and the editor for the comments made throughout the manuscript evaluation process, which contributed to improving the article.

Author Contributions

Conceptualization: FC, BFL, ASS, SG, VQ, KPS, SLA. Data curation: FC, VQ. Formal analysis: FC, BFL, ASS, SG, VQ, KPS, SLA. Funding acquisition: FC. Investigation: FC, BFL, ASS, SG, VQ, KPS, SLA. Methodology: FC, BFL, ASS, SG, VQ, KPS, SLA. Resources: FC, BFL, ASS, SG, VQ, KPS, SLA. Supervision: FC, BFL, ASS, SG, VQ, KPS, SLA. Visualization: FC, BFL, ASS, SG, VQ, KPS, SLA. Project administration: FC. Validation: FC, BFL, ASS, SG, VQ, KPS, SLA. Visualization: FC. Writing – original draft: FC, BFL, ASS, SG, VQ, KPS, SLA. Writing – review and editing: FC, BFL, ASS, SG, VQ, KPS, SLA.

References

- Althoff SL, Carvalho F, Luciano BFL, Garcia JP, Stanke A (2018) First record of *Molossops neglectus* Williams and Genoways, 1980 (Chiroptera, Molossidae) from the state of Santa Catarina, southern Brazil. Check List 14 (1): 167–172. <https://doi.org/10.15560/14.1.167>
- Althoff SL, Tribess B, Reinert MJ, Ferreira MAR, Carvalho F (2017) Expansion of the southern limit of *Vampyroides caraccioli* Thomas, 1889 (Chiroptera, Phyllostomidae) and first record for Santa Catarina state, southern Brazil. Check List 13 (6): 871–877. <https://doi.org/10.15560/13.6.871>
- Alvares CA, Stape JL, Sentelhas PC, Gonçalves JLM, Sparovek G (2013) Köppen's climate classification map for Brazil. Meteorologische Zeitschrift 22 (6): 711–728. <https://doi.org/10.1127/0941-2948/2013/0507>
- Bernard E, Fenton MB (2007) Bats in a fragmented landscape: species composition, diversity and habitat interactions in savannas of Santarém, Central Amazonia, Brazil. Biological Conservation 134 (3): 332–343. <https://doi.org/10.1016/j.biocon.2006.07.021>
- Boada CE, Tirira DG, Camacho MA, Burneo SF (2010) Mammalia, Chiroptera, Thyropteridae, *Thyroptera tricolor* Spix, 1823: distribution extension in Ecuador. Check List 6 (2): 227–229. <https://doi.org/10.15560/6.2.227>
- Bôlla DAS, Ceron K, Carvalho F, Matia DL, Luiz MR, Panatta KA, Pavei DD, Mendonça RA, Zocche JJ (2017) Mastofauna terrestre do Sul de Santa Catarina: mamíferos de médio e grande porte e voadores. Revista Tecnologia e Ambiente 23: 61–78.
- Bornholdt R, Oliveira LR, Fabián ME (2008) Size and shape variability in the skull of *Myotis nigricans* (Schinz, 1821) (Chiroptera: Vespertilionidae) from two geographic areas in Brazil. Brazilian Journal of Biology 68 (3): 623–631. <https://doi.org/10.1590/S1519-69842008000300021>
- Carvalho F, Fábian ME (2011) Mammalia, Chiroptera, Phyllostomidae, *Platyrrhinus recifinus* (O. Thomas, 1901): first confirmed record in the state of Santa Catarina, southern Brazil. Check-List 7 (2): 139–141 <https://doi.org/10.15560/7.2.139>
- Carvalho F, Fabián ME, Menegheti JO (2013) Vertical structure of an assemblage of bats (Mammalia: Chiroptera) in a fragment of Atlantic Forest in Southern Brazil. Zoologia 30 (5): 491–498. <https://doi.org/10.1590/S1984-46702013000500004>
- Carvalho F, Bolla DAS, Patel FM, Miranda JMD, Althoff SL, Zocche JJ (2017) Ampliação de distribuição de *Eumops patagonicus* Thomas, 1924, (Chiroptera: Molossidae) e primeiro registro em ambiente de Restinga na Costa Leste do Brasil. Mastozoología Neotropical 24 (2): 443–450.
- Castaño-Salazar JH, Cardona-Ramírez DM, Orrego O (2004) Notas sobre el primer registro para Caldas del murciélago de ventosas buchiblanco (*Thyroptera tricolor* Spix, 1823). Boletín Científico Centro de Museos Museo de Historia Natural 8: 261–266.
- Chaverri G (2010) Comparative social network analysis in a leaf-roosting bat. Behavioral Ecology and Sociobiology 64: 1619–1630. <https://doi.org/10.1007/s00265-010-0975-3>
- Chaverri G, Kunz TH (2011) All-offspring natal philopatry in a Neotropical bat. Animal Behaviour 82 (5): 1127–1133. <https://doi.org/10.1016/j.anbehav.2011.08.007>
- Chaverri G, Vonnhof MJ (2011) Reproduction and growth in a neotropical insectivorous bat. Acta Chiropterologica 13 (1): 147–155. <https://doi.org/10.3161/150811011x578697>
- Cherem JJ, Althoff SL (2015) Mamíferos de uma área de estepe ombrófila nos estados do Paraná e Santa Catarina, sul do Brasil. Boletim da Sociedade de Mastozoologia 72: 42–50.
- Cherem JJ, Simões-López PC, Althoff SL, Graipel ME (2004) Lista dos mamíferos do Estado de Santa Catarina, Sul do Brasil. Mastozoologia Neotropical 11 (2): 151–184.
- Dechmann DKN, Safi K, Vonnhof MJ (2006) Matching

- Morphology and diet in the disc-winged bat *Thyroptera tricolor* (Chiroptera). *Journal of Mammalogy* 87 (5): 1013–1019. <https://doi.org/10.1644/05-mamm-a-424R2.1>
- Díaz MM, Solari S, Aguirre LF, Aguiar LMS, Barquez RM** (2016) Clave de identificación de los murciélagos de Sudamérica. Programa de Conservación de los Murciélagos de Argentina, Tucumán, Argentina, 160 pp.
- Esbérard CEL, Santos BS, Faria D** (2007) New *Thyroptera tricolor* Spix records in the Atlantic Forest, Brazil (Chiroptera, Thyropteridae). *Brazilian Journal of Biology* 67 (2): 631–637. <https://doi.org/10.1590/S1519-69842007002000028>
- Findley JS, Wilson DE** (1974) Observations on the Neotropical Disk-Winged Bat, *Thyroptera tricolor* Spix. *Journal of Mammalogy* 55 (3): 562–571. <https://doi.org/10.2307/1379546>
- Garbino GST** (2016) Research on bats (Chiroptera) from the state of São Paulo, southeastern Brazil: annotated species list and bibliographic review. *Arquivos de Zoologia* 47 (3): 43–128. <https://doi.org/10.11606/issn.2176-7793.v47i3p43-128>
- Garbino GST, Gregorin R, Lima IP, Loureiro L, Moras LM, Moratelli R, Nogueira MR, Pavan AC, Tavares VC, Nascimento MC, Peracchi AL** (2022) Updated checklist of Brazilian bats: versão 2022. Comitê da Lista de Morcegos do Brasil, Sociedade Brasileira para o Estudo de Quirópteros (SBEQ).
- Hoppe JPM, Pimenta VT, Ditchfield AD**. First occurrence of the recently described Patricia's Disk-winged Bat *Thyroptera wynneae* (Chiroptera: Thyropteridae) in Espírito Santo, southeastern Brazil. *Check List* 10 (3): 645–647 <https://doi.org/10.15560/10.3.645>
- IBGE** (Instituto Brasileiro de Geografia e Estatística) (2012) Manual técnico da vegetação brasileira. 2nd edition. Ministério do Meio Ambiente, Rio de Janeiro, Brazil, 271 pp.
- Kerr J, Packer L** (1998) The impact of climate change on mammal diversity in Canada. *Environmental Monitoring and Assessment* 49: 263–270. <https://doi.org/10.1023/a:1005846910199>
- Lee T Jr.** (2019) Family Thyropteridae (disk-winged bats). In: Wilson DE, Mittermeier RA (Eds.) *Handbook of the mammals of the world. Volume 9: bats*. Lynx Edicions, Barcelona, Spain, 418–423.
- Marinho-Filho J** (1996) Distribution of bat diversity in the southern and southeastern Brazilian Atlantic Forest. *Chiroptera Neotropical* 2 (2): 51–54.
- McCain CM** (2019) Assessing the risks to United States and Canadian mammals caused by climate change using a trait-mediated model. *Journal of Mammalogy* 100 (6): 1808–1817. <https://doi.org/10.1093/jmammal/gyz155>
- Melo AS** (2008) O que ganhamos ‘confundindo’ riqueza de espécies e equitabilidade em um índice de diversidade? *Biota Neotropical* 8 (3): 21–27. <https://doi.org/10.1590/s1676-06032008000300001>
- Miranda JMD, Bernardi IP, Passos FC** (2011) Chave ilustrada para a determinação dos morcegos da Região Sul do Brasil. Universidade Federal do Paraná, Curitiba, Brazil, 51 pp.
- MMA** (Ministério do Meio Ambiente) (2022) Lista oficial de espécies da fauna brasileira ameaçadas de extinção, portaria n°. 444. Brazil, 116 pp. https://www.icmbio.gov.br/cepsul/images/stories/legislacao/Portaria/2020/P_mma_148_2022_altera_anexos_P_mma_443_444_445_2014_atualiza_especies_ameacadas_extincao.pdf. Accessed on: 2023-06-05.
- Montero BK, Gillam EH** (2015) Behavioural strategies associated with using an ephemeral roosting resource in Spix's Disc-winged Bat. *Animal Behaviour* 108: 81–89. <https://doi.org/10.1016/j.anbehav.2015.07.014>
- Moratelli R, Oliveira JA** (2011) Morphometric and morphological variation in South American populations of *Myotis albescens* (Chiroptera: Vespertilionidae). *Zoologia* 28 (6): 789–802. <https://doi.org/10.1590/S1984-46702011000600013>
- Pacheco SM, Sekiama ML, Oliveira KPA, Quintela F, Weber MM, Marques RV, Geiger D, Silveira DD** (2007) Biogeografia de quirópteros da Região Sul. *Ciência e Ambiente* 35: 181–202.
- Passos FC, Miranda JMD, Bernardi IP, Kaku-Oliveira NY, Munster LC** (2010) Morcegos da Região Sul do Brasil: análise comparativa da riqueza de espécies, novos registros e atualizações nomenclaturais (Mammalia, Chiroptera). *Iheringia, Série Zoologia* 100 (1): 25–34. <https://doi.org/10.1590/s0073-47212010000100004>
- Reyes-Amaya N, Lozano-Flórez J, Flores D, Solari S** (2016) Distribution of the spix's disk-winged bat, *Thyroptera tricolor* spix, 1823 (Chiroptera: Thyropteridae) in Colombia, with first records for the middle Magdalena Valley. *Mastozoologia Neotropical* 23 (1): 127–137.
- Russi CH, Carvalho F, Althoff SL** (2021) Phyllostomid bats distribution and richness gradient in a subtropical Brazilian state. *Papéis Avulsos de Zoologia* 61: 1–12. <https://doi.org/10.11606/1807-0205/2021.61.60>
- Sagot M, Giacomini G, Chaves-Ramírez S, Hernández-Pinsón HA, Chaverri G** (2022) Vocal behavior and the use of social information during roost finding. *Frontiers in Ecology and Evolution* 10: 1–9. <https://doi.org/10.3389/fevo.2022.905925>
- Sampaio EM, Kalko EKV, Bernard E, Rodríguez-Herrera B, Handley CO Jr** (2003) A Biodiversity assessment of bats (Chiroptera) in a tropical lowland rainforest of central Amazonia, including methodological and conservation considerations. *Studies on Neotropical Fauna and Environment* 38 (1): 17–31. <https://doi.org/10.1076/snfe.38.1.17.14035>
- Semedo FBT, Garbino ST, Ardente N, Colas-Rosas PF, Torres MA, Rossi RV** (2020) New records of disk-winged bats *Thyroptera tricolor* Spix, 1823 and *T. devivoi* Gregorin, Gonçalves, Lim & Engstrom, 2006 (Chiroptera: Thyropteridae) for the Brazilian Amazonia and Cerrado. *Boletim do Museu Paraense Emílio Goeldi. Ciências Naturais* 15 (3): 817–827. <https://doi.org/10.46357/bcnaturais.v15i3.213>
- Solari S** (2015) *Thyroptera devivoi*. The IUCN Red List of Threatened Species 2015: e.T136594A21996185. <https://doi.org/10.2305/iucn.uk.2015-4.rlts.t136594a21996185.en>. Accessed on: 2023-06-05.
- Solari S** (2016) *Thyroptera wynneae*. The IUCN Red List

- of Threatened Species 2016: e.T88151033A88151036. <https://doi.org/10.2305/iucn.uk.2016-3.rlts.t88151033a88151036.en>. Accessed on: 2023-06-05.
- Solari S, Velazco P** (2016) *Thyroptera lavalii*. The IUCN Red List of Threatened Species 2016: e.T21878A21985717. <https://doi.org/10.2305/iucn.uk.2016-2.rlts.t21878a21985717.en>. Accessed on: 2023-06-05.
- Solari S, Villada-Cadavid T** (2018) *Thyroptera discifera*. The IUCN Red List of Threatened Species 2018: e.T21877A21985811. <https://doi.org/10.2305/iucn.uk.2018-2.rlts.t21877a21985811.en>. Accessed on: 2023-06-05.
- Storz JF, Balasingh J, Bhat HR, Natan TP, Doss PDS, Prakash AA, Kunz TH** (2001) Clinal variation in body size and sexual dimorphism in an Indian fruit bat, *Cynopterus sphinx* (Chiroptera: Pteropodidae). *Biological Journal of the Linnean Society* 75 (1): 17–31.
- Tavares V, Mantilla H** (2015) *Thyroptera tricolor* (errata version published in 2016). The IUCN Red List of Threatened Species 2015: e.T21879A97207863. <https://doi.org/10.2305/iucn.uk.2015-4.rlts.t21879a21985559.en>. Accessed on: 2023-06-05.
- Tavares VC, Nobre CC, Palmuti CFS, Nogueira EPP, Gomes JD, Marcos MH, Silva RF, Farias SG, Bobrowiec PED** (2017) The bat fauna from southwestern Brazil and its affinities with the fauna of western Amazon. *Acta Chiropterologica* 19 (1): 93–106. <https://doi.org/10.3161/15081109acc2017.19.1.007>
- Turcios-Casco MA, Medina-Fitoria A** (2018) Occurrence of *Hylonycteris underwoodi* (Chiroptera, Phyllostomidae) and *Thyroptera tricolor* (Chiroptera, Thyropteridae) in Honduras. *Studies on Neotropical Fauna and Environment* 54 (1): 69–72. <https://doi.org/10.1080/01650521.2018.1544205>
- Velazco PM, Gregorin R, Voss RS, Simmons NB** (2014) Extraordinary local diversity of disk-winged bats (Thyropteridae: Thyroptera) in Northeastern Peru, with the description of a new species and comments on roosting behavior. *American Museum Novitates* 3795: 1–28. <http://repositorio.ufla.br/jspui/handle/1/39990>
- Vonhof MJ, Fenton MB** (2004) Roost availability and population size of *Thyroptera tricolor*, a leaf-roosting bat, in north-eastern Costa Rica. *Journal of Tropical Ecology* 20 (3): 291–305. <https://doi.org/10.1017/s0266467404001403>
- Vonhof MJ, Whitehead HAL, Fenton MB** (2004) Analysis of Spix's Disc-winged Bat association patterns and roosting home ranges reveal a novel social structure among bats. *Animal Behaviour* 68 (3): 507–521. <https://doi.org/10.1016/j.anbehav.2003.08.025>
- Wilson D** (2008) Family Thyropteridae Miller, 1907. In: Gardner AL (Ed.) *Mammals of South America*. University of Chicago Press, Chicago/London, 392–395.
- Wimsatt WA, Enders AC** (1980) Structure and morphogenesis of the uterus, placenta, and paraplacental organs of the neotropical disc-winged bat *Thyroptera tricolor* Spix (Microchiroptera: Thyropteridae). *American Journal of Anatomy* 159 (2): 209–243. <https://doi.org/10.1002/aja.1001590208>
- Yusefi GH, Kamran S, Tarroso P, Brito JC** (2021) The impacts of extreme climate change on mammals differ among functional groups at regional scale: the case of Iranian terrestrial mammals. *Diversity and Distributions* 27 (9): 1634–1647.