



First records of *Lepidophyma zongolicum* García-Vázquez, Canseco-Márquez & Aguilar-López, 2010 (Squamata, Xantusiidae) from Veracruz, Mexico

ANGEL I. CONTRERAS-CALVARIO^{1,2}, MEDARDO ARREORTÚA¹, ADÁN BAUTISTA-DEL MORAL^{3,4*}, JORGE L. CASTILLO-JUÁREZ⁵, RENE AVALOS-VELA⁵

1 Instituto Politécnico Nacional, Centro Interdisciplinario de Investigación para el Desarrollo Integral Regional (CIIDIR) Oaxaca, Laboratorio de Ecología de Anfibios, Santa Cruz Xoxocotlán, Oaxaca, México • AICC: acontrerascalvario@gmail.com  <https://orcid.org/0000-0001-9674-8571> • MA: medardo.am@hotmail.com  <https://orcid.org/0000-0003-3775-0369>

2 Herpetario de Veracruz, Predios e Instalaciones que Manejan Vida Silvestre (PIMVS), Veracruz, México

3 Laboratorio de Herpetología, Departamento de Zoología, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad de México, México • ABM: bautistadelmoral@gmail.com  <https://orcid.org/0000-0001-8461-0225>

4 Posgrado en Ciencias Biológicas, Instituto de Biología, Universidad Nacional Autónoma de México, Ciudad de México, México

5 Facultad de Ciencias Biológicas y Agropecuarias, Universidad Veracruzana, Amatlán de los Reyes, Veracruz, México • RAV: renevela@live.com.mx  <https://orcid.org/0000-0002-0452-7817>

* Corresponding author

Abstract. *Lepidophyma* Duméril, 1851 is the most diverse genus among the nocturnal lizards, with 23 species distributed from the Atlantic and Pacific slopes of Mexico southeast into Panama. Currently, the available information on the ecology and distribution of these species is limited mainly due to their saxicolous habits. Here we present five new state records of *Lepidophyma zongolicum* from Atoyac, a locality in the state of Veracruz, Mexico. These new data expand the geographic range of this species by 48 km from its nearest previously known occurrence.

Keywords. Altas Montañas region, biogeography, geographic range extension, Zongolica Night Lizard

Academic editor: Ricardo Palacios Aguilar

Received 26 May 2023, accepted 29 August 2023, published 18 September 2023

Contreras-Calvario AI, Arreortúa M, Bautista-del Moral A, Castillo-Juárez JL, Avalos-Vela R (2023) First records of *Lepidophyma zongolicum* García-Vázquez, Canseco-Márquez & Aguilar-López, 2010 (Squamata, Xantusiidae) from Veracruz, Mexico. Check List 19 (5): 635–638. <https://doi.org/10.15560/19.5.635>

Introduction

Lepidophyma Duméril, 1851 is the most diverse genus among the night lizard family (Lara-Tufiño and Nieto-Montes de Oca 2021; Grünwald et al. 2023). They are distributed from Nuevo León and Colima along the Atlantic and Pacific slopes, respectively, to southeastern Panama (Noonan et al. 2013; Palacios-Aguilar et al. 2018; Arenas-Moreno et al. 2021; Lara-Tufiño and Nieto-Montes de Oca 2021). These lizards are found in a wide variety of habitats, from xeric scrub to temperate and tropical forests (Bezy and Camarillo-Rangel 2002). Despite the great diversity of habitats and their wide distribution, little is known about *Lepidophyma*

lizards, mainly due to their saxicolous habits and their physiological and behavioral requirements (Arenas-Moreno et al. 2018; Muñoz-Nolasco et al. 2019).

Despite this, five new species of *Lepidophyma* have been described since 2010, most known only from their type locality (García-Vázquez et al. 2010; Palacios-Aguilar et al. 2018; Lara-Tufiño and Nieto-Montes de Oca 2021): *L. zongolicum* García-Vázquez, Canseco-Márquez & Aguilar-López, 2010; *L. inagoi* Palacios-Aguilar, Santos-Bibiano & Flores-Villela, 2018; *L. lusca* Arenas-Moreno, Muñoz-Nolasco, Bautista-del Moral, Rodríguez-Miranda, Domínguez-Guerrero & Méndez-de la Cruz, 2021; *L. ramírezi* Lara-Tufiño & Nieto-Montes de Oca, 2021; and *L. jasonjonesi* Grünwald,

Reyes-Velasco, Ahumada-Carrillo, Montaño-Ruvalcaba, Franz-Chávez, La Forest, Ramírez-Chaparro, Terán-Juárez & Borja-Jiménez, 2023. *Lepidophyma zongolicum* has only been recorded from its type locality on the Atlantic slopes of southeastern Puebla (Tepeyac, municipality of Eloxochitlán), regardless of having been described in 2010 (García-Vázquez et al. 2010). Here, we extend the geographic range of *L. zongolicum* to the east and present the first records of it from Veracruz, Mexico.

Methods

Field surveys were conducted in Atoyac municipality, Veracruz, Mexico, by three of us (AICC, JLCJ, and RAV). The vegetation in this area is made up of tropical semideciduous forests (Acevedo 1988; Rivera-Hernández 2015). The average annual temperature is 22.8 °C, the maximum mean temperature of the warmest month is 32 °C, the minimum mean temperature of the coldest month is 13.2 °C, and the annual precipitation is 1578 mm. These climatic data were obtained through WorldClim v. 2, with 2.5 minute resolution (Fick and Hijmans 2017).

Collected specimens of *Lepidophyma zongolicum* were euthanized with pentobarbital, fixed with 10% formalin, and preserved in 70% ethanol for permanent storage. The specimens were deposited at the Museo de

Zoología, Facultad de Estudios Superiores Zaragoza, Universidad Nacional Autónoma de México. Collecting was conducted under permit FAUT-0093 issued to Daniel Lara-Tufiño by the Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT).

The geodetic datum used is WGS84.

Results

Lepidophyma zongolicum García-Vázquez, Canseco-Márquez & Aguilar-López, 2010

Figure 1

New records. MEXICO – Veracruz • Atoyac, Antiguo Túnel Ferroviario; 18.9220, –096.7676; 540 m elev.; 06.V.2017; Jorge L. Castillo-Juarez obs.; active, at the exit of a rock tunnel, on limestone; 1 ♂ adult; CNAR-RF 925 (Fig. 1A) • same locality (Fig. 1B); 23.IV.2022; Angel I. Contreras-Calvario & Daniel Lara-Tufiño leg.; inside a stone tunnel, in cracks about 1 cm wide; 2 adults, 2 juveniles; MZFZ-4409–4412 (Fig. 1C).

Identification. The specimens were identified, following the original description (García-Vázquez et al. 2010), by the number of dorsal scales, total femoral pores, number of lateral tubercles, number of small granular tail scales separating the larger ones, number of paravertebral rows, number of foot lamellae, and number of gular scales.




Figure 1. *Lepidophyma zongolicum* and habitat in Atoyac, Veracruz, Mexico. **A.** Adult male. **B.** Habitat of *L. zongolicum* in Veracruz, Mexico. **C.** Adult female. Photographs by JLCJ and RAV.




Figure 2. New state records of *Lepidophyma zongolicum* from Veracruz (red star) and previously published locality (green circle) (García-Vázquez et al. 2010).

Discussion

Lepidophyma zongolicum was included by Torres-Hernández et al. (2021) in their list of the herpetofauna of Veracruz, indicating that this species is found in the Sierra Madre Oriental and in the Transverse Mexican Volcanic Axis. However, they did not mention any locality, specimens deposited in scientific collections, or photographs that confirm the presence of this species in Veracruz. Therefore, we consider that the type locality at El Tepeyac, Puebla state, is the only formally recorded locality for this species. Consequently, our record extends the geographic range of *L. zongolicum* by 48 km in a straight line from the nearest known record (Fig. 2).

Nevertheless, our new records suggest that *L. zongolicum* is found in a wide variety of habitats, as mentioned by García-Vázquez et al. (2010). Generating reliable information on the distribution of secretive species, such as *L. zongolicum*, brings with it positive effects such as the study of ecophysiological and demographic requirements, which will ultimately serve as an important tool to know and conserve enigmatic species such as the nocturnal lizards of the genus *Lepidophyma*.

Acknowledgements

We thank Daniel Lara-Tufiño for corroborating the identity of the specimens and for using his collection permit, Raquel Hernández-Austria for her help with

fieldwork, Karla I. Contreras-Calvario for linguistic revision, and Ricardo Palacios-Aguilar for his comments that improved the manuscript.

Author Contributions

Conceptualization: RAV, ABM, AICC. Data curation: AICC, MA, ABM. Formal analysis: ABM, AICC. Funding acquisition: AICC, ABM, JLCJ, MA, RAV. Investigation: ABM, MA, JLCJ, AICC, RAV. Methodology: RAV, AICC, JLCJ. Project administration: AICC. Resources: AICC, JLCJ. Supervision: AICC. Writing – original draft: ABM, AICC, MA. Writing – review and editing: ABM, AICC, MA.

References

- Acevedo R (1988) La vegetación de la Sierra de Atoyac, Veracruz, México. Batchelor's thesis, Facultad de Biología. Universidad Veracruzana. Xalapa, Mexico, 102 pp.
- Arenas-Moreno DM, Santos-Bibiano R, Muñoz-Nolasco FJ, Charruau P, Méndez-de la Cruz FR (2018) Thermal ecology and activity patterns of six species of tropical night lizards (Squamata: Xantusiidae: *Lepidophyma*) from Mexico. *Journal of Thermal Biology* 75: 97–105. <https://doi.org/10.1016/j.jtherbio.2018.06.001>
- Arenas-Moreno DM, Munoz-Nolasco FJ, del Moral-Bautista A, Rodriguez-Miranda LA, Dominguez-Guerreiro SF, Mendez de la Cruz FR (2021) A new species of *Lepidophyma* (Squamata: Xantusiidae) from San Luis

- Potosí, México, with notes on its physiological ecology. *Zootaxa* 4949: 4949. <https://doi.org/10.11646/zootaxa.4949.1.6>
- Bezy RL, Camarillo-Rangel JL** (2002) Systematics of xantusiid lizards of the genus *Lepidophyma*. *Contributions in Science* 493: 1–41. <https://doi.org/10.5962/p.226792>
- Bezy RL** (2019) Night lizards: field memoirs and a summary of the Xantusiidae. ECO Herpetological Wear & Publishing, Rodeo, NM, USA, 220 pp.
- Fick SE, Hijmans RJ** (2017) WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37: 4302–4315. <https://doi.org/10.1002/joc.5086>
- García-Vázquez UO, Canseco-Marquez L, Aguilar-Lopez JL** (2010) A new species of night lizard of the genus *Lepidophyma* (Squamata: Xantusiidae) from southern Puebla, Mexico. *Zootaxa* 2657: 47–54. <https://doi.org/10.11646/zootaxa.2657.1.4>
- Grünwald CI, Reyes-Velasco J, Ahumada-Carrillo IT, Montaña-Ruvalcaba C, Franz-Chávez H, La Forest BT, Ramírez-Chaparro R, Terán-Juárez S, Borja-Jiménez JM** (2023) A new species of saxicolous *Lepidophyma* (Squamata, Xantusiidae) from Tamaulipas, Mexico. *Herpetozoa* 36: 9–21. <https://doi.org/10.3897/herpetozoa.36.e96184>
- Lara-Tufiño JD, Nieto-Montes de Oca A** (2021) A new species of night lizard of the genus *Lepidophyma* (Xantusiidae) from southern Mexico. *Herpetologica* 77: 320–334. <https://doi.org/10.1655/herpetologica-d-21-00019.1>
- Muñoz-Nolasco FJ, Arenas-Moreno DM, Santos-Bibiano R, Bautista-del Moral A, Gandarilla-Aizpuro FJ, Brindis-Badillo DA, Méndez de La Cruz FR** (2019) Evaporative water loss of some habitat-restricted mexican lizard species. *Herpetological Conservation and Biology* 14: 51–66.
- Noonan BP, Pramuk JB, Bezy RL, Sinclair EA, Queiroz K de, Sites JW** (2013) Phylogenetic relationships within the lizard clade Xantusiidae: using trees and divergence times to address evolutionary questions at multiple levels. *Molecular Phylogenetics and Evolution* 69: 109–122. <https://doi.org/10.1016/j.ympev.2013.05.017>
- Palacios-Aguilar R, Santos-Bibiano R, Flores-Villela O** (2018) A new species of *Lepidophyma* (Squamata: Xantusiidae) from the pacific lowlands of Guerrero, Mexico. *Journal of Herpetology* 52: 327–331. <https://doi.org/10.1670/17-061>
- Rivera-Hernandez JE** (2015) Flora, vegetación y priorización de áreas de conservación del Parque Nacional Cañón del Río Blanco, Veracruz, México. Doctoral thesis, Instituto Tecnológico de Costa Rica, Universidad Estatal a Distancia, Provincia de Cartago, Costa Rica, 254 pp.
- Torres-Hernández LA, Ramírez-Bautista A, Cruz-Elizalde R, Hernández-Salinas U, Berriozabal-Islas C, Desantis DL, Johnson JD, Rocha A, García-Padilla E, Mata-Silva V, Fucsko LA, Wilson LD** (2021) The herpetofauna of Veracruz, Mexico: composition, distribution, and conservation status. *Amphibian & Reptile Conservation* 15: e285.