



Distribution of *Thereuonema tuberculata* (Wood, 1862) (Chilopoda, Scutigeromorpha, Scutigeridae) in Hokkaido, Japan

NATSUKI HIRAKIZAWA¹, HIJIRI NODA², RYOTARO WAKIMURA³, KOJI TOJO², TAKEO YAMAUCHI^{1*}

¹ Laboratory of Entomology, Obihiro University of Agriculture and Veterinary Medicine, Obihiro, Hokkaido, Japan • NH: s23180006@st.obihiro.ac.jp • TY: tyamauchi@obihiro.ac.jp

² Institute of Mountain Science, Shinshu University, Matsumoto, Nagano, Japan • HN: 23ss506j@shinshu-u.ac.jp • KT: ktojo@shinshu-u.ac.jp

³ Tokai University, Sapporo campus, Sapporo, Hokkaido, Japan • RW: wakimur9drpep@gmail.com

* Corresponding author

Abstract. *Thereuonema tuberculata* (Wood, 1862), a centipede, is distributed mainly in Asia. In Japan, *T. tuberculata* is found throughout Honshu Island and southwards, although less commonly to the north on Hokkaido. Our survey additionally showed that this species occurs commonly at altitudes of 0–540 m on Hokkaido. The northern limit of this species in Japan seems to be Nakagawa Town, Hokkaido. *Thereuonema tuberculata* was also collected on isolated islands off southwestern Hokkaido. Although previous studies have suggested distribution of *T. tuberculata* in Hokkaido is anthropogenic, it may be natural.

Keywords. Indoor invasion, Japanese house centipede, northern limit, scutigeromorph centipedes, subarctic zone, synanthropic area

Academic editor: Peter Decker

Received 1 June 2023, accepted 22 August 2023, published 30 August 2023

Hirakizawa N, Noda H, Wakimura R, Tojo K, Yamauchi T (2023) Distribution of *Thereuonema tuberculata* (Wood, 1862) (Chilopoda, Scutigeromorpha, Scutigeridae) in Hokkaido, Japan. Check List 19 (5): 599–604. <https://doi.org/10.15560/19.5.599>

Introduction

The house-centipede order Scutigeromorpha (class Chilopoda) has about 95 species worldwide (Edgecombe 2011). Two species are known from Japan: *Thereuopoda clunifera* (Wood, 1862) and *Thereuonema tuberculata* (Wood, 1862) (Shinohara et al. 2015). In Japan, *T. clunifera* is distributed in the Kanto region and southwards, and *T. tuberculata* is known to be commonly distributed throughout the main island of Honshu and southwards. *Thereuonema tuberculata* is less commonly found on the northern island of Hokkaido (Shinohara et al. 2015). *Thereuonema tuberculata* is also known to be distributed in China, Taiwan, mainland Korea, and Jeju Island, Korea (Würlmli 1975; Stoev 2002). Additionally, it has been reported as an introduced species in the United Kingdom and the United States (Barber 2011; Reeves 2017; Reeves and Miller 2022).

The western part of the Oshima Peninsula of Hokkaido is in a temperate zone, and the rest of Hokkaido

except for the western and southern parts of the Oshima Peninsula is in a subarctic zone (Miyamoto 2009). *Thereuonema tuberculata* has rarely been reported from Hokkaido, with records only from Otaru City, Yakumo Town, Sapporo City, and Rankoshi Town (Takakuwa 1943, 1955; Takakuwa and Shinohara 1968; Tanabe 2001; Ōomote 2021). For that reason, it has been suggested that this species might occur on Hokkaido for anthropogenic reasons (Takakuwa 1955; Takakuwa and Shinohara 1968; Shinohara 1974). Here, we investigate the distribution of *T. tuberculata* in Hokkaido, Japan.

Methods

We conducted a distribution survey of *Thereuonema tuberculata* throughout Hokkaido, Japan from 2018 to 2023. Specimens from the collection in the Higashitai-setsu Nature Center (HTMNH) were also examined. All specimens were preserved in 70% or 99.5% ethanol

and were identified under a stereomicroscope (Olympus SZ4045) according to Würmli (1975) and Barber (2011). The seventh tergite of one specimen was skinned with tweezers, and the female gonopods were excised. The whole specimen and the first tarsal segment of leg 6 were photographed with a digital camera (TG-6 Tough, Olympus), and photographs of the seventh tergite and female gonopods were taken with a single-lens reflex camera (Canon EOS Kiss X10).

Results

Order Scutigeromorpha Pocock, 1895

Family Scutigeridae Leach, 1814

Subfamily Thereuoneminae Verhoeff, 1905

Genus *Thereuonema* Verhoeff, 1904

Thereuonema tuberculata (Wood, 1862)

Figures 1, 2

New records. JAPAN, Hokkaido Prefecture – **Hokkaido Island – Okhotsk District** • Abashiri City, Minami 6 jō Higashi; 44°01'14"N, 144°16'31"E, ca. 10 m alt.; 28.VII.2021; Y. Uni leg.; inside a house; 1 ♂ • Abashiri City, Minami 6 jō Higashi; 44°01'14"N, 144°16'31"E, ca. 10 m alt.; 31.VII.2021; Y. Uni leg.; inside a house; 1 ♂ • Abashiri City, Minami 6 jō Higashi; 44°01'14"N, 144°16'31"E, ca. 10 m alt.; 7.VIII.2021; Y. Uni leg.; inside a house; 1 ♂ • Abashiri City, Minami 6 jō Higashi; 44°01'14"N, 144°16'31"E, ca. 10 m alt.; 25.VIII.2021; Y. Uni leg.; inside a house; 1 ♂ • Bihoro Town, Mitomi;




Figure 1. Adult female of *Thereuonema tuberculata* in Obihiro City, Hokkaido, Japan. **A.** Habitus, dorsal view. **B.** Part of tergite VII, dorsal view. **C.** First tarsal segment of leg 6. **D.** Gonopod of female, ventral view.




Figure 2. Distributional records of *Thereuonema tuberculata*. Closed circles - locality of specimens examined, open circles = previous records (Takakuwa 1943, 1955; Takakuwa and Shinohara 1968; Tanabe 2001; Ōmote 2021), closed triangle = eyewitness information, grey area = previous records (Würmli 1975).

43°48'40"N, 144°06'00"E, ca. 60 m alt.; 15.VII.2018; collector unknown; 2 ♂ 1 ♀ • Bihoro Town, Aoyamaminami; 43°49'11"N, 144°06'35"E, ca. 30 m alt.; 29.VIII.2021; K. Machida leg.; 1 ♂ • Shari Town, Honmachi; 43°54'51"N, 144°39'59"E, ca. 3 m alt.; 17.VII.2021; N. Hirakizawa leg.; 1 ♂ 1 ♀ • Shari Town, Honmachi; 43°54'40"N, 144°40'07"E, ca. 7 m alt.; 18.VII.2021; T. Murakami, N. Domoto, and N. Hirakizawa leg.; 1 ♂ 2 juv. • Shari Town, Utorokagawa; 44°04'11"N, 144°59'48"E, ca. 60 m alt.; 18.VII.2021; N. Domoto leg.; 1 ♂ 1 juv. • Shari Town, Utorokagawa; 44°04'11"N, 144°59'48"E, ca. 60 m alt.; 18.VII.2021; N. Hirakizawa leg.; 1 juv. • Koshimizu Town, Hamakoshimizu; 43°55'59"N, 144°27'13"E, ca. 10 m alt.; 15.IX.2022; R. Wakimura leg.; 1 ♂ 1 ♀ – **Kamikawa District** • Nakagawa Town, Homare; 44°48'05"N, 142°04'01"E, ca. 20 m alt.; 12.IX.2022; N. Hirakizawa leg.; 1 ♂ 2 ♀ 1 juv. • Shibetsu City, Nishishibetsu-chô (Shibetsu City Museum); 44°10'00"N, 142°22'32"E, ca. 130 m alt.; 22.X.2021; T. Honbe leg.; inside the museum; 1 ♀ • Shibetsu City, Nishishibetsu-chô; 44°10'00"N, 142°22'32"E, ca. 150 m alt.; 10.IX.2022; R. Wakimura and H. Arimatsu leg.; 1 ♂ • Nayoro City, Nishi 6 jô Minami; 44°20'35"N, 142°27'18"E; 8.VII.2018; collector unknown; 1 ♂ • Asahikawa City, Hanasaki-chô; 43°47'12"N, 142°21'59"E, ca. 110 m alt.; 5.VII.2018; collector unknown; 4 ♂ 5 ♀ 1 ex. • Asahikawa City, Shunkodai 4 jô; 43°49'19"N, 142°21'20"E, ca. 150 m alt.;

9.XI.2021; K. Saito leg.; 1 ♀ • Furano City, Higashitorinuma; 43°20'35"N, 142°26'05"E, ca. 170 m alt.; 11.VII.2018; collector unknown; 1 ♂ 3 ♀ – **Sorachi District** • Yubari City, Fukuzumi; 43°03'58"N, 141°59'08"E, ca. 340 m alt.; 9.VII.2018; collector unknown; 5 ♂ 1 ex. • Iwamizawa City, Kitamura (Kitamura elementary school); 43°15'28"N, 141°41'25"E, ca. 10 m alt.; 18.VIII.2022; R. Kaneda leg.; 1 ♂ – **Tokachi District** • Kamishihoro Town, Nukabiragensenkyô (Higashitai-setsu Nature Center); 43°21'55"N, 143°11'37"E, ca. 540 m alt.; 4.XI.2022; A. Tanaka leg.; 1 ♀, HTMNH-48672 • Shihoro Town, Nishi 2 sen; 43°09'54"N, 143°14'40"E, ca. 200 m alt.; 9.V.2022; T. Hosaka leg.; 1 juv. • Obihiro City, Higashi 4 jô Minami; 42°55'04"N, 143°12'44"E, ca. 40 m alt.; 23.VIII.2022; N. Ohi leg.; inside a house; 3 juv. • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 13.VIII.2020; F. Yamauchi leg.; 1 ♀ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 19.VIII.2020; R. Yamauchi leg.; inside a house; 1 ♀ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 23.VIII.2020; F. Yamauchi leg.; 1 ♂ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 27.VIII.2020; F. Yamauchi leg.; 1 ♂ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 26.IV.2021; T. Yamauchi leg.; inside a house; 1 ♀ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.;

- 27.X.2020; F. Yamauchi leg.; inside a house; 1 ♀ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 24.VI.2021; T. Yamauchi leg.; inside a house; 1 ♂ 1 ♀ • Obihiro City, Nishi 4 jô Minami; 42°54'19"N, 143°12'03"E, ca. 50 m alt.; 27.VII.2021; R. Yamauchi and F. Yamauchi leg.; 1 ♀ • Obihiro City, Nishi 5 jô Minami; 42°53'29"N, 143°12'06"E, ca. 50 m alt.; 19.IV.2022; F. Yamauchi leg.; 1 ♀ • Obihiro City, Nishi 5 jô Minami; 42°53'29"N, 143°12'06"E, ca. 50 m alt.; 26.IV.2022; F. Yamauchi leg.; 1 ♂ 2 exs. • Obihiro City, Nishi 18 jô Minami; 42°55'21"N, 143°09'39"E, ca. 40 m alt.; data unknown; collector unknown; 2 ♀ 1 ex. • Obihiro City, Nishi 18 jô Minami; 42°55'23"N, 143°09'37"E, ca. 50 m alt.; 12.V.2021; S. Takamiya leg.; 1 ♀ • Obihiro City, Nishi 19 jô Minami; 42°52'39"N, 143°10'33"E, ca. 80 m alt.; 4.VI.2021; H. Takese leg.; 1 ♀ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 6.IX.2020; M. Iwasa leg.; 1 ♂ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 7.IX.2020; M. Iwasa leg.; inside a house; 1 ♀ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 8.IX.2020; M. Iwasa leg.; inside a house; 1 ♀ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 28.IX.2020; M. Iwasa leg.; inside a house; 1 ♂ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 22.X.2020; M. Iwasa leg.; inside a house; 1 ♂ • Obihiro City, Nishi 22 jô Minami; 42°54'59"N, 143°08'13"E, ca. 60 m alt.; 21.IX.2021; M. Iwasa leg.; inside a house; 1 juv. • Obihiro City, Ôdôriminami, Ôdôri Park; 42°54'33"N, 143°12'22"E, ca. 40 m alt.; 30.VII.2021; F. Yamauchi leg.; 1 juv. • Obihiro City, Ôdôriminami, Ôdôri Park; 42°54'33"N, 143°12'22"E, ca. 40 m alt.; 30.VI.2022; F. Yamauchi leg.; 1 juv. • Obihiro City, Midorigaoka, Green and Flower Center; 42°54'36"N, 143°11'16"E, ca. 50 m alt.; 25.II.2022; K. Yano and K. Azuta leg.; inside a greenhouse; 1 ♂ • Obihiro City, Midorigaoka; 42°54'17"N, 143°10'52"E, ca. 60 m alt.; 28.III.2022; A. Sogawa leg.; inside a house; 1 ♀ • Obihiro City, Minami-nomori; 42°52'53"N, 143°10'18"E, ca. 70 m alt.; 1.VIII.2020; N. Hirakizawa leg.; 1 ♂ 1 ♀ • Obihiro City, Minami-nomori; 42°52'53"N, 143°10'18"E, ca. 70 m alt.; 24.VIII.2022; Y. Konno leg.; inside a house; 1 ♀ • Obihiro City, Ôzora-chô; 42°52'52"N, 143°09'09"E, ca. 80 m alt.; 25.IV.2021; K. Takahashi leg.; 1 ♀ • Obihiro City, Ôzora-chô; 42°52'52"N, 143°09'09"E, ca. 80 m alt.; 9.VII.2021; K. Takahashi leg.; 1 ♂ • Otofuke Town, Ryokuyôdai; 42°58'14"N, 143°11'52"E, ca. 70 m alt.; 10.IX.2021; M. Furuto leg.; 1 ♂ 4 ♀ • Otofuke Town, Ryokuyôdai; 42°58'14"N, 143°11'52"E, ca. 70 m alt.; 13.IX.2021; M. Furuto leg.; 1 ♂ 1 ♀ • Uraho Town, Kita-machi; 42°48'50"N, 143°39'48"E, ca. 30 m alt.; 14.VI.2021; N. Hirakizawa leg.; 2 juv. • Uraho Town, Shin-machi; 42°48'28"N, 143°39'27"E, ca. 20 m alt.; 5.X.2021; M. Mochida leg.; inside a house; 1 ♂ – **Ishikari District** • Sapporo City, Minami-ku, Monami Park; 42°59'12"N, 141°20'10"E, ca. 80 m alt.; 18.V.2021; R. Wakimura leg.; 2 ♀ • Sapporo City, Atsubetsu-ku, Atsubetsu-chô, Konopporo; 43°03'12"N, 141°29'46"E, ca. 50 m alt.; 11.VIII.2021; M. Mizushima leg.; 1 ♀ • Ebetsu City, Bunkyo-dai; 43°03'48"N, 141°29'28"E, ca. 40 m alt.; 15.VIII.2011; O. Suda leg.; 1 ex., HTMNH-44687 – **Hidaka District** • Erimo Town, Hon-machi; 42°00'51"N, 143°09'00"E, ca. 30 m alt.; 7.VIII.2021; N. Hirakizawa leg.; 2 juv. • Erimo Town, Hon-machi; 42°00'51"N, 143°09'00"E, ca. 30 m alt.; 7.VIII.2021; A. Sogawa leg.; 2 juv. • Erimo Town, Yamato; 42°01'17"N, 143°08'43"E, ca. 10 m alt.; 7.VIII.2021; N. Hirakizawa leg.; 3 juv. • Samani Town, Horoman; 42°04'41"N, 143°02'06"E, ca. 3 m alt.; 13.VII.2018; collector unknown; 4 ♂ 2 ♀ • Urakawa Town, Nishihorobetsu; 42°08'36"N, 142°50'24"E, ca. 2 m alt.; 16.VI.2022; N. Hirakizawa leg.; 1 ♂ 1 ♀ 3 juv. • Shinhidaka Town, Shizunaimidori-chô; 42°20'33"N, 142°22'25"E, ca. 5 m alt.; 16.IV.2023; N. Yoshikawa leg.; inside a house; 1 ♀ • Biratori Town, Nibutani; 42°38'13"N, 142°09'23"E, ca. 60 m alt.; 3.VIII.2021; Y. Osada leg.; inside a museum; 1 ♀ – **Shiribeshi District** • Otaru City, Hariusu-chô; 43°09'18"N, 141°07'09"E, ca. 40 m alt.; 4.VIII.2021; S. Tsuchiya leg.; inside a garage; 1 ♀ • Yoichi Town, Tomisawa-chô; 43°12'36"N, 140°46'16"E, ca. 4 m alt.; 3.VII.2018; collector unknown; 1 ♀ • Iwanai Town, Mandai; 42°59'02"N, 140°30'57"E, ca. 2 m alt.; 3.VII.2018; collector unknown; 2 ♀ 1 ex. – **Iburi District** • Tomakomai City, Sankou-chô; 42°39'29"N, 141°37'09"E, ca. 10 m alt.; 1.VIII.2021; A. Kodama leg.; inside a house; 1 ♂ • Noboribetsu City, Noboribetsuonsen-chô; 42°29'45"N, 141°08'29"E, ca. 200 m alt.; 18.IX.2021; T. Shibuya leg.; 1 juv. • Sobetsu Town, Nakajima Island; 42°35'44"N, 140°50'41"E, 0 m alt.; 20.VII.2021; R. Wakimura leg.; 1 ♂ – **Hiyama District** • Esashi Town, Ubagami-chô; 41°51'58"N, 140°07'06"E, ca. 2 m alt.; 13.VIII.2021; R. Wakimura leg.; 1 juv. • Esashi Town, Ubagami-chô; 41°51'58"N, 140°07'06"E, ca. 2 m alt.; 13.VIII.2021; N. Hirakizawa leg.; 2 juv. – **Oshima District** • Hakodate City, Syowa; 41°48'41"N, 140°43'59"E, ca. 9 m alt.; 18.VII.2018; collector unknown; 4 ♂ 2 ♀ • Hakodate City, Goryokaku-chô; 41°47'53"N, 140°45'24"E, ca. 20 m alt.; 19.IX.2022; R. Wakimura leg.; 1 juv. • Matsumae Town, Matsushiro; 41°25'52"N, 140°06'30"E, ca. 30 m alt.; 12.VIII.2021; R. Wakimura leg.; 1 juv. • Fukushima Town, Fukushima; 41°28'45"N, 140°15'20"E, ca. 5 m alt.; 19.IX.2022; R. Wakimura leg.; 1 juv. • Shiriuchi Town, Yunosato; 41°35'43"N, 140°17'41"E, ca. 50 m alt.; 19.IX.2022; R. Wakimura leg.; 2 ♂ • Shiriuchi Town, Yunosato; 41°35'43"N, 140°17'41"E, ca. 50 m alt.; 13.VIII.2021; N. Hirakizawa leg.; 2 juv. – **Okushiri Island** • Okushiri Town, Akaishi; 42°08'46"N, 139°31'06"E, ca. 50 m alt.; 24.VII.2021; S. Inagaki leg.; 1 ♀ • Okushiri Town, Akaishi; 42°08'45"N, 139°30'56"E, ca. 10 m alt.; 28.VII.2021; S. Inagaki leg.; 1 ♂ 1 ♀ • Okushiri Town, Akaishi; 42°08'46"N, 139°31'06"E, ca. 10 m alt.; 1.VIII.2021; S. Inagaki leg.; 1 ♂ • Okushiri Town, Okushiri or Akaishi; 28.VII.2021; S. Inagaki leg.; 1 ♂ 2 ♀ • Okushiri Town, Okushiri; 42°10'21"N, 139°30'14"E, ca. 60 m alt.; 31.VII.2021; F. Suzuki leg.; 1 ♀ • Okushiri Town, Aonae; 42°03'41"N, 139°26'53"E, ca. 20 m alt.; 2.VIII.2021; S. Watabe leg.; 1 juv. – **Oshima-ôshima**

Island • Oshima-ôshima; 41°30'34"N, 139°21'16"E; 30.VII.2021; S. Hori leg.; crack of rock; 1 ♀.

Eyewitness information. Japan, Hokkaido Prefecture–**Hokkaido Island** – **Okhotsk District** • Mombetsu City, Shiomi-chô; 44°21'26"N, 143°21'07"E; Y. Sudo obs. • Engaru Town; 44°03'37"N, 143°31'14"E; N. Taguchi obs. – **Kamikawa District** • Takasu Town; 43°51'22"N, 142°21'35"E; K. Saito obs. – **Kushiro District** • Kushiro City, Midorigaoka; 42°59'06"N, 144°24'14"E; S. Kitada obs. • Kushiro City, Chiyonoura; 42°58'04"N, 144°23'19"E; S. Kitada obs.

Identification. All specimens obtained were identified as *T. tuberculata* based on the appearance of numerous long, spike-like spiculae on the tergites (Fig. 1B), the absence of a pair of spine-bristles at the distal end of the first tarsal segment of the fifth or sixth to fourteenth legs (Fig. 1C), and the morphology of the female gonopod (Fig. 1D).

Discussion

Neither *Thereuonema clunifera* nor *Scutigera coleoptrata* (Linnaeus, 1758), widely distributed worldwide, was found in our survey. It is likely that *T. tuberculata* is the only scutigermorph centipede present on Hokkaido. *Thereuonema tuberculata* has been previously recorded from only two cities and two towns in southwestern Hokkaido (Fig. 2; Table 1), but our survey shows that it is distributed in almost all areas of Hokkaido at altitudes between 0 and 540 m. The species was found mainly in human-influenced environments such as houses, schools, roadside stations, and parks. It was also collected on the lacustrine Nakajima Island (Toya Lake, Iburi District), as well as Okushiri Island and Oshima ôshima Island (Fig. 2).

Although previous studies have suggested the distribution of *T. tuberculata* in Hokkaido is anthropogenic (Takakuwa 1955; Takakuwa and Shinohara 1968; Shinohara 1974), our data suggest it may also be naturally occurring. However, despite surveys and interviews, no *T. tuberculata* were found in Wakkanai City, Sarufutsu Village, Hamatombetsu Town, or Horonobe Town on the northern tip of Hokkaido 30–90 km north of Nakagawa Town, or in Akkeshi Town, and no records of *T. tuberculata* were found in the northern territories of Hokkaido (Iturup Island, Shikotan Island, Kunashir Island, Habomai Islands). Thus, the northern limit of this species in Japan seems to be Nakagawa Town (44°48'06"N).

Table 1. Previous distributional records of *Thereuonema tuberculata* in Hokkaido, Japan.

Locality	Reference
Otaru City	Takakuwa (1943, 1955); Takakuwa and Shinohara (1968)
Sapporo City	Tanabe (2001)
Rankoshi Town	Ôomote (2021)
Yakumo Town	Takakuwa and Shinohara (1968)

Murakami (1956) described the life history of laboratory-reared specimens of *T. tuberculata* (as *T. hilgen-dorfi*), which probably came from Ehime Prefecture, Japan (although not specified). He found that *T. tuberculata* stops molting around November and overwinters in that state, and that this species has a life span of 3–4 years (Murakami 1956). Most of Hokkaido has a subarctic climate (Miyamoto 2009); for example, the mean minimum temperature in Obihiro City (south-central Hokkaido) in February 1990–2020 reached –12 °C (Japan Meteorological Agency 2023). This species was collected indoors in Obihiro City in February, suggesting that adult *T. tuberculata* overwinter in Hokkaido to avoid the cold winter temperatures.

Acknowledgements

We would like to express our cordial thanks to all those individuals listed below (we have omitted honorific titles here): Takahiro Murakami, Masaki Yoshida, Yasuyuki Oppata, Airi Tanaka, Nanami Domoto, Mitsuhiko Iwasa, Tetsuya Honbe, Kimio Abe, Kazuki Miura, Shigehisa Hori, Miki Mizushima, Sinta Inagaki, Shingo Watabe, Fuma Suzuki, Aiko Kodama, Yoshiyasu Machida, Kyoko Machida, Makoto Mochida, Kazunori Saito, Nao Taguchi, Shuzo Tsuchiya, Yoshikazu Uni, Toshiyasu Nakaoka, Yoshihiro Osada, Yasuo Konno, Masayuki Furuto, Syota Kitada, Takuma Hosaka, Akie Sogawa, Kanori Takahashi, Sota Takamiya, Hiroki Arimatsu, Yuta Okuno, Takanobu Shibuya, Nagisa Ohi, Riichi Yamauchi, Fumi Yamauchi, Haruna Takese, Kaho Azuta, Koharu Yano, Sakura Asato, Rima Kaneda, Yuna Sudo, Moeko Toda, Norie Yoshikawa, Ryosuke Kuwahara, Glen Hill, and three anonymous reviewers. This work was partly supported by JSPS KAKENHI grant number JP23KJ0075.

Author Contributions

Conceptualization: NH, TY. Data curation: NH, HN. Funding acquisition: TY, KT, NH. Investigation: RW, NH, HN, TY. Project administration: TY, NH. Supervision: TY. Visualization: NH. Writing – original draft: NH. Writing – review and editing: TY, KT, NH, HN, RW.

References

- Barber AD (2011) *Thereuonema tuberculata*, a scutigermorph centipede from China, found in a warehouse at Swindon. Bulletin of the British Myriapod and Isopod Group 25: 49–50.
- Edgecombe GD (2011) Taxonomic overview. Order Scutigermorpha. In: Minelli A (Ed.) Treatise on zoology—anatomy, taxonomy, biology. The Myriapoda. Volume 1. Brill, Leiden, the Netherlands, 363–370. https://doi.org/10.1163/9789004188266_020
- Japan Meteorological Agency (2023) [Historical weather data]. <https://www.data.jma.go.jp/obd/stats/etrn/>. Japan

- Meteorological Agency, Tokyo, Japan. Accessed on: 2023-4-10. [In Japanese]
- Miyamoto M** (2009) A reexamination on Koeppen climate classification in northern Tohoku and Hokkaido. *Geographical Studies* 84: 111–117. [in Japanese]
- Murakami Y** (1956) The life history of *Thereuonema hilgendorfi* Verhoeff (Chilopoda, Scutigeridae). *Zoological Magazine* 65: 42–46. [in Japanese with English summary]
- Ôomote S** (2021) 蘭越の昆虫II, カメムシ目・コウチュウ目・その他の目・昆虫以外の虫 [Insects of Rankoshi II, Hemiptera, Coleoptera, other order, invertebrates except for insects]. Self-published, Rankoshi, Hokkaido, Japan, 89 pp. [in Japanese]
- Reeves WK** (2017) Discovery of an exotic population of *Thereuonema tuberculata* (Chilopoda: Scutigeromorpha), the Japanese house centipede, in Ohio, USA. *The American Midland Naturalist* 177: 162–164. <https://doi.org/10.1674/0003-0031-177.1.162>
- Reeves WK, Miller MM** (2022) *Thereuonema tuberculata* (Wood, 1862) (Chilopoda, Scutigeromorpha, Scutigeridae) from forested habitats in North America. *Check List* 18(2): 431–434. <https://doi.org/10.15560/18.2.431>
- Shinohara K** (1974) 多足類の採集と観察: 方法と実例 [Observation and collecting of myriapods: skill and practical guidebook]. New Science, Tokyo, Japan, 109 pp. [in Japanese]
- Shinohara K, Takano M, Ishii K** (2015) Chilopoda. In: Aoki J (Ed.) Pictorial keys to soil animals of Japan, second edition. Tokai University Press, Kanagawa, Japan, 873–910. [in Japanese]
- Stoev P** (2002) The scutigeromorphs (Chilopoda: Scutigeromorpha) in the collection of the National Museum of Natural History, Sofia. *Historia Naturalis Bulgarica* 15: 79–85.
- Takakuwa Y** (1943) 日本産ゲジの種 [Species of scutigeromorph centipedes in Japan]. *Acta Arachnologica* 8: 50–58. [in Japanese]
- Takakuwa Y** (1955) ゲジの内外の解剖及分類 附 恩師故理學博士丘先生の追憶 [Morphology and classification of the scutigeromorph centipedes, with a memory of the late Dr. Asajiro Oka]. Gakufu-Shoin Publisher, Tokyo, Japan, 60 pp. [in Japanese]
- Takakuwa Y, Shinohara K** (1968) 多足類研究夜話 [The dawn of myriapod research: Yosioki Takakuwa's notes with annotations by Keizaburo Shinohara]. *Atypus* 46147: 39–42. [in Japanese]
- Tanabe T** (2001) The biology of centipedes and millipedes. Tokai University Press. Tokyo, Japan, 178 pp. [in Japanese]
- Würmli M** (1975) Revision der Hundertfüßer-Gattung *Thereuonema* (Chilopoda: Scutigeridae). *Entomologica Germanica* 2: 189–196.