



Local extinction of *Melanophrynniscus montevidensis* (Anura: Bufonidae) in the Argentine Pampas

Federico L. Agnolin,^{1,2} Elián L. Guerrero³

1 Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Av. Ángel Gallardo 470, C1405DJR, Buenos Aires, Argentina. **2** Fundación de Historia Natural “Félix de Azara”, Universidad Maimónides, Hidalgo 775, C1405BDB, Buenos Aires, Argentina. **3** División Plantas Vasculares. Museo de La Plata. Paseo del Bosque s/n, La Plata, 1900, Buenos Aires, Argentina. CONICET.

Corresponding author: Federico L. Agnolin, fedeagnolin@yahoo.com.ar

Abstract

Melanophrynniscus montevidensis inhabits sandy dunes along the Rio de la Plata shoreline and Atlantic coast in Uruguay and southern Brazil. Here we report 2 specimens from different localities at eastern Buenos Aires province, Argentina, that were collected prior to the 1970s. *Melanophrynniscus montevidensis* probably inhabited patchy sand dune environments in Buenos Aires. It is possible that the disappearance of natural environments (including dunes) in the Argentine Pampas resulted in the local extinction of *M. montevidensis*, a species sensitive to habitat fragmentation and anthropic modifications.

Key words

Sand dunes; *Melanophrynniscus montevidensis*; local extinction; Buenos Aires; Argentina.

Academic editor: Ross MacCulloch | Received 12 January 2017 | Accepted 1 May 2017 | Published 10 July 2017

Citation: Agnolin FL, Guerrero EL (2017) Local extinction of *Melanophrynniscus montevidensis* (Anura: Bufonidae) in the Argentine Pampas. Check List 13 (4): 11–15. <https://doi.org/10.15560/13.4.11>

Introduction

Melanophrynniscus is an endemic Neotropical bufonid genus composed of 29 species (Frost 2017) distributed in Brazil, Paraguay, Bolivia, Uruguay and Argentina (Cruz and Caramaschi 2003, Kwet et al. 2005, Di Bernardo et al. 2006, Peloso et al. 2012). In Argentina 12 species have been recorded, mainly distributed in forested areas in the northern half of the country, and in patches in the southern and central hills (Baldo and Krauczuk 1999, Céspedes et al. 2000, Prigioni and Langone 2000, Céspedes and Motte 2001, Céspedes and Motte 2007, Baldo 2001, Cabrera 2001, Caramaschi and Cruz 2002, Céspedes 2003, Céspedes 2008, Céspedes 2009, Baldo and Basso 2004, Agnolin and Bogan 2014).

Currently, the genus is represented by 3 phenotypic

groups (Céspedes and Motte 2001, Cruz and Caramaschi 2003) that are poorly known with regard to specific content and most behavioral aspects (Céspedes et al. 2000). Most of the living species are included within the phenotypic group *M. stelzneri* (Prigioni and Langone 2000, Caramaschi and Cruz 2002, Céspedes 2008, Baldo and Basso 2004). *Melanophrynniscus. montevidensis* belongs to the *M. stelzneri* group. It inhabits a narrow corridor of sandy dunes along the Rio de la Plata shoreline and Atlantic coast in Uruguay (Langone 1995, Núñez et al. 2004, Maneyro and Kwet 2008) and Alvorada (Barra do Chui) at the southern extreme of Rio Grande do Sul, Brazil (Tedros et al. 2001, Bernardo-Silva et al. 2012) (Fig. 1). A species informally known as *M. aff. montevidensis* is known from the Ventania and Tandilia Hill




Figure 1. Map showing new locality records for *Melanophrynniscus montevidensis*. Black shading indicates current distribution of the species (based on Zank et al. 2014), black triangles constitute Buenos Aires records of the species (see Vaira et al. 2012), and black stars include localities here reported.

systems and the South Coastal Dunes in Buenos Aires province, Argentina (Cairo et al. 2008, Cortelezzi et al. 2015, Friedman et al. 2016, Kacoliris et al. 2017). In older works these populations were assigned to the Uruguayan

species *M. montevidensis* based on coloration similarities (Cei 1980, Gallardo 1987, Gallardo and Varela de Olmedo 1992), but Klappenbach and Langone (1992), based on geographical distribution, concluded that they could be different species, a hypothesis supported by Céspedes et al. (2000). These specimens are still under study; however, genetic information indicates that they belong to a single species (Vaira et al. 2012), a criterion that is followed here.

Because of its restricted geographical distribution, as well as the strong anthropic modifications of the area, the populations of *M. montevidensis* in South America are decreasing, and thus this species is treated as “Vulnerable” at the global level (Langone 2004).

Here we report 2 specimens of *Melanophrynniscus montevidensis* from different locations in eastern Buenos Aires province, Argentina. These constitute one of the few available records for the country, and the only ones from the Eastern Argentine Pampas.

Methods

Specimens here reported are housed at the Colección Herpetología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia”, Buenos Aires, Argentina (MACN He).

MACN He 04513 comes from Zelaya city, Pilar, Buenos Aires province, Argentina. It was collected by A.G. Freers in December 1924 (Fig. 2A, B).

MACN He 18139 comes from Banfield city, Lomas de Zamora, Buenos Aires province, Argentina. It was collected by Juan R. Deriu on 6 June 1963 (Fig. 2C, D).

There is a possibility that these 2 specimens of *Melanophrynniscus montevidensis* may be mislabeled. However, it is unlikely that specimens collected at different times by different collectors, and coming from neighboring localities in northern Buenos Aires, represent 2 parallel and independent mistakes. Furthermore, the specimens have been labeled 3 times: one label is associated with the preserved specimen, and the other 2 labels are housed in different catalogues of the MACN-He collection, reinforcing the possibility that a label mistake for these individuals is unlikely.

Results

The specimens here reported exhibit a unique combination of characters that support identification as *M. montevidensis*: dorsal surface black, with pale lateral blotches, sparse blotches on the dorsal surface of head and shoulders, small scattered pale blotches ventrally and weakly granular dorsal skin. The presence of a large cream-colored patch along the posterior portion of the abdomen and the thigh indicates the presence of a wide red blotch in life (Langone 2002, Cruz and Caramaschi 2003, Maneyro and Kwet 2008). Furthermore, this combination of characters distinguishes *M. montevidensis* from other species of the genus (see Céspedes 2003, Céspedes 2008, Céspedes 2009, Céspedes and Motte 2001,




Figure 2. *Melanophrynniscus montevidensis* reported here. **A, B.** MACN He 04513, dorsal and ventral views, respectively. **C, D.** MACN He 18139, dorsal and ventral views, respectively. Scale bar: 1 cm.

Céspedes and Motte 2007). In general aspect *M. atroluteus* is similar to *M. montevidensis* (Maneyro and Kwet 2008) but differs from the latter in lacking (or having only scattered) yellow blotches along the lateral surface of the body and dorsal surface of the head (Klappenbach and Langone 1992, Cruz and Caramaschi 2003, Maneyro and Kwet 2008). Furthermore, *M. atroluteus* exhibits notably rugose and spiny dorsal skin (Kwet et al. 2005, Maneyro and Kwet 2008).

Discussion

The anuran fauna of central Argentina (Buenos Aires and La Pampa provinces) shows a species composition similar to that of faunas from Uruguay and Rio Grande do Sul (Gallardo 1965, Maneyro et al. 1995). Among these, a particular group has been named as “Atlantic shore-fauna” (Maneyro et al. 1995, Maneyro and Carreira 2006). This includes sand-dune adapted amphibians such as *Melanophryniscus montevidensis*, among other taxa.

As indicated above, in the Ventania and Tandilia hill systems of southern Buenos Aires province there is a *M. montevidensis* population that is disjunct from the main distributional range of the species. In historical times this population was even more geographically widespread, as demonstrated by the finding of specimens among sand dunes near Bahía Blanca city by Charles Darwin (Bell 1843, Cabrera 2001). It is worthy to mention that recent reports are based on old museum specimens, and that the species has not been found in the last 100 years, in spite of fieldwork and sampling in the coastal dunes of the province (Kacoliris et al. 2017).

Currently, sand dunes, which are the most suitable habitat for *M. montevidensis*, are totally absent in the northern tip of Buenos Aires province. However, the presence of sand dune environments in northeastern Buenos Aires province can be inferred from old botanical collections that indicate the existence of plants associated with sand dunes in Ensenada and Punta Indio Districts (Hicken 1910, Parodi 1940). Parodi (1940) published photographs showing sand dunes in Monte Veloz (Punta Indio District). This locality has provided some rare zoological records, such as the only Solifugae known from northern Buenos Aires (Maury 1977), and the rodent *Necromys benefactus* (Galliari and Pardiñas 2000), both recorded there prior to 1950, in semiarid and open environments. Furthermore, specimen MACN He 04513 of *M. montevidensis* comes from Zeloya, a town that in the early 20th century yielded specimens of *Ctenomys talarum*, a rodent of open sandy environments that is extirpated from most of its original area (Agnolin and Lucero 2013). *Melanophryniscus montevidensis* was probably an inhabitant of those patchy sand dune environments.

The disappearance of undisturbed environments (including dunes) from part of the Argentine Pampas affected a large number of amphibians (Lavilla 2001). This probably resulted in the local extinction of *M. montevidensis*, a species sensitive to habitat fragmentation

due to climatic change and anthropic modifications (Langone 1995, Langone 2003, Langone 2004; see also Zank et al. 2014). Uruguayan populations of *M. montevidensis* are considered as Endangered (Maneyro and Langone 2001, Carreira and Maneyro 2015), and several local populations from Montevideo and Canelones departments have been extirpated (Langone 2003). A probable cause for this decline is the urbanization and consequent habitat fragmentation along the Uruguayan coast (Langone 1995). It is expected that climatic change would affect negatively this species in the future (Toranza and Maneyro 2013, Toranza et al. 2016).

In summary, based on these data, we infer that in the past, *M. montevidensis* occurred in the eastern Buenos Aires Pampean grasslands, and became locally extinct by the mid 20th century, probably due to degradation of sandy soil habitats.

Acknowledgements

We especially thank S. Nenda and S. Bogan for their help during the study of the material, as well as for useful comments. We thank J. Faivovich for allowing us to study the specimens under his care. We acknowledge F. Suazo Lara for his help during the revision of materials at MACN. Special thanks to 2 anonymous reviewers and editors L. Langone and R. MacCulloch, who made insightful and helpful comments on an early version of the manuscript.

Authors' Contributions

FLA and EG collected the data, wrote the text, and made the analysis.

Literature Cited

- Agnolin FL, Bogan S (2014) Algunos comentarios sobre la posible presencia de *Melanophryniscus dorsalis* (Anura, Bufonidae) en Argentina y sus implicancias biogeográficas. Historia Natural 3: 107–114.
- Agnolin FL, Lucero SO (2013) Sobre la presencia de *Ctenomys talarum* (Rodentia Ctenomyidae) en el Noreste de la provincia de Buenos Aires, Argentina. Historia Natural 3: 77–85.
- Agostini MG, Saibene PE, Roesler I, Bilenca D (2016) Amphibians of northwestern Buenos Aires province, Argentina: checklist, range extensions and comments on conservation. Check List 12 (6): 1998. <https://doi.org/10.15560/12.6.1998>
- Agostini MG, Corteleazzi A, Berkunsky I, Soler G, Burrowes P (2015) First record of *Batrachochytrium dendrobatidis* infecting threatened populations of Tandilean Red-belly toad (*Melanophryniscus* aff. *montevidensis*) in Argentina. Revista Mexicana de Biodiversidad 86: 826–828. <https://doi.org/10.1016/j.rmb.2015.07.007>
- Baldo JD (2001) Acerca de la localidad tipo y la distribución geográfica de *Melanophryniscus klappenbachi* Prigioni and Langone, 2000 (Anura: Bufonidae). Cuadernos de Herpetología 15: 141–142.
- Baldo JD, Basso NG (2004) A new species of *Melanophryniscus* Galhardo, 1961 (Anura: Bufonidae), with comments on the species of the genus reported for Misiones, northeastern Argentina. Journal of Herpetology 38: 140–150.
- Baldo JD, Krauczuk ER (1999) *Melanophryniscus devincenzi* Klappenbach, 1968 (Anura: Bufonidae). Primer registro para la República Argentina. Cuadernos de Herpetología 13: 101.

- Bell T (1843) Zoology of the voyage of the HMS Beagle, under the command of Captain Fitzroy, RN, during the years 1832 to 1836. Edited and superintended by Charles Darwin, naturalist to the expedition. Part 5. Reptiles. 51 pp. <https://doi.org/10.5962/bhl.title.14216>
- Bernardo-Silva J, Martins-Ferreira C, Maneyro R, Freitas TRO (2012) Identification of priority areas for conservation of two endangered parapatric species of red-bellied toads, using ecological niche models and hotspot analysis. *Natureza a Conservação* 10: 207–213. <https://doi.org/10.4322/natcon.2012.026>
- Cabrera MR (2001) Sobre la distribución geográfica de *Melanophryniscus s. stelzneri* (Weyenberg) (Anura, Bufonidae). *Cuadernos de Herpetología* 15: 137–139.
- Cairo SL, Zalba SM, Úbeda CA (2008) Reproductive behaviour of *Melanophryniscus* sp. From Sierra de la Ventana (Buenos Aires, Argentina). *South American Journal of Herpetology* 3: 10–14. [https://doi.org/10.2994/1808-9798\(2008\)3\[10:RBOMSF\]2.0.CO;2](https://doi.org/10.2994/1808-9798(2008)3[10:RBOMSF]2.0.CO;2)
- Caramaschi U, Cruz CAG (2002) Taxonomic status of *Atelopus pachyrhynchos* Miranda-Ribeiro, 1920, redescription of *Melanophryniscus tumifrons* (Boulenger, 1905), and descriptions of two new species of *Melanophryniscus* from the state of Santa Catarina, Brazil (Amphibia, Anura, Bufonidae). *Arquivos do Museu Nacional* 60: 303–314.
- Carreira S, Maneyro R (2015) Lista Roja de los Anfibios y Reptiles del Uruguay. Una evaluación del estado de conservación de la herpetofauna de Uruguay sobre la base de los criterios de la Unión Internacional para la Conservación de la Naturaleza. 64 pp.
- Cei JM (1993) Reptiles del centro, centro-oeste y sur de la Argentina. Herpetofauna de las zonas áridas y semiáridas. Museo Regionale di Science Naturali Torino, Monografia 4: 1–527.
- Céspedes JA (2003) Una nueva especie de *Melanophryniscus* Gallardo, 1961 de Argentina (Amphibia: Anura: Bufonidae). *Actas de Reuniones de Comunicaciones Científicas y Tecnológicas UNNE*: 1–4.
- Céspedes JA (2008) Una nueva especie de *Melanophryniscus* Gallardo, 1961 de Argentina (Amphibia: Anura: Bufonidae). *Facena* 24: 35–48.
- Céspedes JA (2009) Primer registro de *Melanophryniscus krauczukii* (Basso y Baldo, 2000) para Corrientes, Argentina. *Comunicaciones Científicas y Tecnológicas*, UNNE: 1.
- Céspedes JA, Motte M (2001) Distribución de sapos del género *Melanophryniscus* (Gallardo, 1961) en Argentina y Paraguay (Anura: Bufonidae). *Boletín de la Asociación Herpetológica Española* 12: 71–76.
- Céspedes JA, Motte M (2007) Una nueva especie de *Melanophryniscus* Gallardo 1961 (Amphibia: Anura: Bufonidae) de Paraguay. *Facena* 23: 29–40.
- Céspedes JA, Golobiski F, Alvarez B (2000) Distribución del género *Melanophryniscus* (Gallardo, 1961) en Argentina y Paraguay (Anura: Bufonidae). *Comunicaciones Científicas y Tecnológicas*, UNNE: 1–4.
- Cortelezzi A, Berkunsky I, Simoy MV, Cepeda R, Marinelli C, Kacoliris F (2015) Are breeding sites a limiting factor for the Tandilean red-belly toad (Bufonidae) in Pampean highland grasslands? *Neotropical Biology and Conservation* 10: 182–186.
- Cruz CAG, Caramaschi U (2003) Taxonomic status of *Melanophryniscus stelzneri dorsalis* (Mertens, 1933) and *Melanophryniscus stelzneri fulvoguttatus* (Mertens, 1937) (Amphibia, Anura, Bufonidae). *Boletim do Museu Nacional* 500: 1–11.
- Di-Bernardo M, Maneyro R, Grillo H (2006) A new species of *Melanophryniscus* (Anura: Bufonidae) from Rio Grande do Sul, southern Brazil. *Journal of Herpetology* 40: 261–266. <https://doi.org/10.1670/05-008.1>
- Friedman M, Cepeda RE, Cortelezzi A, Simoy MV, Marinelli CB, Kacoliris F, Dopazo J, Berkunsky I (2016) Searching for an elusive anuran: a detection model based on weather forecasting for the Tandilean Red-belly Toad. *Herpetological Conservation and Biology* 11: 476–485.
- Frost DR (2017) Amphibian Species of the World: an Online Reference. Version 6.0. American Museum of Natural History, New York. <http://research.amnh.org/herpetology/amphibia/index.html>. Accessed on: 2017-3-11.
- Gallardo JM (1965) Consideraciones zoogeográficas y ecológicas sobre los anfibios de la provincia de La Pampa, Argentina. *Revista del Museo argentino de Ciencias Naturales “Bernardino Rivadavia”* 1: 57–77.
- Gallardo JM (1987) Anfibios argentinos, Guía para su identificación. Librería Agropecuaria, Buenos Aires, 99 pp.
- Gallardo JM, Varela de Olmedo E (1992) Anfibios de la República Argentina: ecología y comportamiento. *Fauna de Agua Dulce de la República Argentina* 41: 1–116.
- Galliari CA, Pardinas UFJ (2000) Taxonomy and distribution of the sigmodontine rodents of genus *Necromys* in central Argentina and Uruguay. *Acta Theriologica* 45: 211–232.
- Hicken CM (1910) *Chloris Platensis* Argentina. *Apuntes de Historia Natural* 2: 3–292.
- Kacoliris F, Williams J, di Pietro D (2017) Herpetofauna de las dunas costeras bonaerenses. In: Athor J, Celsi CE (Eds) *La Costa Atlántica de Buenos Aires: Naturaleza y Patrimonio Cultural*. Fundación de Historia Natural Félix de Azara, Buenos Aires, 234–251.
- Klappenbach MA, Langone JA (1992) Lista sistemática y sinonímica de los anfibios del Uruguay. Con comentarios y notas sobre su distribución. *Anales del Museo Nacional de Historia Natural de Montevideo* 8: 163–222.
- Kwet A, Maneyro R, Mebs D, Zillikens A (2005) Advertisement calls of *Melanophryniscus dorsalis* (Mertens, 1933) and *M. montevidensis* (Philippi, 1902), two parapatric species from southern Brazil and Uruguay, with comments on the *Melanophryniscus stelzneri* group (Anura: Bufonidae). *Salamandra* 41: 3–20.
- Langone JA (1995) Ranas y sapos del Uruguay (Reconocimiento y aspectos biológicos). Museo D. A. Larrañaga, Serie de Divulgación 5: 1–123.
- Langone JA (2003) Amphibia. *Anales del Museo Nacional de Historia Natural y Antropología* 10: 1–12.
- Langone JA (2004) *Melanophryniscus montevidensis*. In: IUCN 2014 . IUCN Red List of Threatened Species. Version 2014.1 . <http://www.iucnredlist.org>. Accessed on 2017-3-11.
- Langone JA (2002) *Melanophryniscus*, interesantes sapitos uruguayos. Documentos de divulgación Museo Nacional de Historia Natural y Antropología 4: 1–11.
- Lavilla EO (2001) Amenazas, declinaciones poblacionales y extinciones en anfibios argentinos. *Cuadernos de Herpetología* 15: 59–82.
- Maneyro R, Forni F, Santos M (1995) Anfibios del Departamento de Rocha. Probides, Rocha, Uruguay, 21 pp.
- Maneyro R, Carreira S (2006) La herpetofauna de la costa uruguaya. In: Menafra R, Rodriguez L, Scarabino F, Conde D (Eds) *Bases Para la Conservación y Manejo de la Costa Uruguaya*. Graphis, Montevideo, 233–246.
- Maneyro R, Kwet A (2008) Amphibians in the border region between Uruguay and Brazil: updated species list with comments on taxonomy and natural history (Part I: Bufonidae). *Stuttgarter Beiträge zur Naturkunde* 1: 95–121.
- Maneyro R, Langone J (2001) Categorización de los anfibios del Uruguay. *Cuadernos de Herpetología* 15: 107–118.
- Maury E (1977) Notas sobre la sistemática y distribución geográfica de *Procleobis patagonicus* (Holmberg 1876) (Solifugae, Ammotrechidae, Saronominae). *Physis* 36: 283–293.
- Núñez D, Maneyro R, Langone JA, de Sá RO (2004) Distribución geográfica de la fauna de anfibios del Uruguay. *Smithsonian Herpetological Information Service* 134: 1–34. <https://doi.org/10.5479/si.23317515.134.1>
- Parodi LR (1940) La distribución geográfica de los talares en la Provincia de Buenos Aires. *Darwiniana* 4: 33–56.
- Peloso PLV, Faivovich J, Grant T, Gasparini JL, Haddad CFB (2012) An extraordinary new species of *Melanophryniscus* (Anura, Bufonidae) from southeastern Brazil. *American Museum Novitates*

- 3762: 1–31. <https://doi.org/10.1206/3762.2>
- Prigione CM, Langone JA (2000) Una nueva especie de *Melanophryniscus*-*cus* Gallardo, 1961, de Argentina y Paraguay (Amphibia, Anura, Bufonidae). Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 195: 1–12.
- Tedros M, Kolenc F, Borteiro C (2001) *Melanophryniscus montevidensis* (Philippi, 1902) (Anura, Bufonidae). Novedades zoogeográficas. Cuadernos de Herpetología 15: 143.
- Toranza C, Maneyro R (2013) Potential effects of climate change on the distribution of an endangered species: *Melanophryniscus montevidensis* (Anura, Bufonidae). Phylomedusa 12: 97–106.
- Toranza C, Brazeiro A, Maneyro R (2016) Anfibios amenazados de Uruguay: efectividad de las áreas protegidas ante el cambio climático. Ecología Austral 26: 138–149.
- Vaira M, Akmentins MS, Attademo M, Baldo D, Barrasso D, Barriónuevo S, Basso N, Blotto B, Cairo S, Cajade R, Céspedes J, Corbalán V, Chilote P, Duré M, Falcione C, Ferraro D Gutierrez, FR, Ingaramo M, Junges C, Lajmanovich R, Lescano JN, Marangoni F, Martinazzo L, Martí R, Moreno L, Natale GS, Pérez Iglesias JM, Peltzer P, Quiroga L, Rosset S, Sanabria E, Sanchez L, Schaefer E, Úbeda C, Zaracho V. 2012. Categorización del estado de conservación de los anfibios de la República Argentina. Cuadernos de Herpetología 26: 131–159.
- Zank, C., F. Gertum, M. Abadie, D. Baldo, R. Maneyro and M. Borges-Martins (2014) Climate change and the distribution of Neotropical red bellied toads (*Melanophryniscus*, Anura, Amphibia): How to prioritize species and populations? Plos ONE 9: e94625. <https://doi.org/10.1371/journal.pone.0094625>