

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Cnidaria, Anthozoa, Octocorallia, Anthelidae, *Tripalea clavaria* (Studer, 1878): Distribution extension, first record for sub-tropical waters, Bahia, Brazil.

Carlos Daniel Pérez  
Bárbara de Moura Neves

GPA – Anthozoan Research Group, Núcleo de Biologia – Centro Acadêmico de Vitória, UFPE, rua do Alto do Reservatório s/n, Bela Vista, CEP 55608-680, Vitória de Santo Antão, PE, Brazil.  
E-mail: gpa@ufpe.br

*Tripalea clavaria* (Studer, 1878) is a colonial scleraxonian octocoral that presents a continuous distribution from Rio de Janeiro to the patagonics coasts in Argentine. This species has records in the Brazilian coast from Rio de Janeiro to Rio Grande do Sul, Uruguay and Argentine extending to the Strait of Magellan (Studer 1879; Küenthal 1919; Deichmann 1936; Bayer 1961; Verseveldt 1967; Tixier-Durivault 1969-1970; Tommasi 1971; Capitoli and Bemvenuti 2004; Excoffon et al. 2004) (Figure 1).




Figure 1. Distribution map of *Tripalea clavaria*.

Jacques Laborel collaborating with the scientific vessel Calypso during in 1962 and carried out samples of benthic cnidarians along the Brazilian coast between Santos and Recife. Afterwards, in the years 1966 and 1967, he finished his studies in Todos os Santos bay (Bahia) (Laborel 1969). Part of the material collected by Laborel was deposited in the cnidarian collection of the Departament of Oceanography at the Universidade Federal de Pernambuco (DOCEAN-CNI). One sample with four colonies of *T. clavaria* was analized (DOCEAN-CNI 101) proceeding from the Bahia state ( $12^{\circ}58'15''$  S,  $38^{\circ}30'39''$  W) collected by the scientific vessel Calypso in 1962 (Figure 2).


Bayer (1961) had associated this species to cold waters with temperatures lesser than  $20^{\circ}$  C, fixing the northern limit of its distribution in the proximities of the Guanabara Bay in Rio de Janeiro, coincidently with the north limit of the influence of the Malvinas current. However, our results contrast with Bayer's assumption, indicating that *T. clavaria* is not a cold waters exclusive species, appearing to be a potential eurythermic species. This emphasizes the great plasticity of adaptation to different environments. Based upon this new report from the litoral of Bahia, *T. clavaria* presents a continuous distribution along the sudamerican coast between Argentine, Uruguay and Brazil, crossing natural zoogeographic barriers, such as the estuary of Rio de la Plata with a low salinity of 32 ‰ (Pérez and Zamponi 2004).

Contrarily to the endemic status for the subantarctic region, assigned by Bayer (1961), this octocoral is also a faunistic component of the subtropical fauna in the South Western Atlantic Ocean.

### Acknowledgments

We are grateful to Dr. André Mauricio Santos (Núcleo de Biologia, CAV-UFPE) for carrying out the map.

NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** *Tripalea clavaria* (Studer, 1878) (DOCEAN-CNI 101): a, colony; b, cross section with longitudinal canals; c, detail of lacunar inner cortex; d, longitudinal section with retracting polyps.

**Literature cited**

- Bayer, F. M. 1961. The shallow-water Octocorallia of the West Indian region. A manual for marine biologists. Studies Fauna of Curaçao and other Caribbean Islands 12: 1-373.
- Capitoli, R. C. and C. Bemvenuti. 2004. Distribuição batimétrica e variações de diversidade dos macroinvertebrados bentônicos da plataforma continental e talude superior no extremo sul do Brasil. Atlântica 26(1): 27-43.
- Deichmann, E. 1936. The Alcyonaria of the Western part of the Atlantic Ocean. Memoirs

- of the Museum of Comparative Zoölogy 53: 253-308.
- Excoffon, A. C., F. H. Acuña, M. O. Zamponi, and G. N. Genzano. 2004. Reproduction of the temperate octocoral *Tripalea clavaria* (Octocorallia, Anthothelidae) from sublittoral outcrops of Mar del Plata, Argentina. Journal of Marine Biological Association of United Kingdom 84(4658): 1-5.
- Kükenthal, W. 1919. Gorgonaria. Wissenschaftliche Ergebnisse der Deutschen Tiefsee-Expedition auf dem Dampfer "Valdivia" 13(2): 1-946, pls. 30-89.

## NOTES ON GEOGRAPHIC DISTRIBUTION

- Laborel, J. 1969. Les peuplements de Madréporaires des côtes tropicales du Brésil. Annales de l'Université D'Abidjan Série E II (3): 1-261.
- Pérez, C. D. and M. O. Zamponi, 2004. New records of octocorals (Cnidaria, Anthozoa) from Southwestern Atlantic Ocean. Zoogeographic considerations. Zootaxa 630: 1-12.
- Studer, T. 1879. Übersicht der Anthozoa Alcyonaria, welche während der Reise S.M.S. Gazelle um die Erde gesammelt wurden. Monatsbericht der K. preussischen Akademie der Wissenschaften Berlin 1878: 632-688.
- Tixier-Durivault A. 1969-1970. Octocoralliaires. Campagne de la Calypso au large des côtes Atlantiques de l'Amerique du Sud (1961-1962). I. Annales de L'Institut Oceanographic 47: 145-169.
- Tommasi, L. R. 1971. Ocorrência de *Tripalea clavaria* (Studer, 1878) na plataforma continental do Rio Grande do Sul. Boletim do Instituto oceanográfico de São Paulo 20: 27-32.
- Verseveldt, J. 1967. The Octocorallia collected by R/V "Vema" in the Atlantic Ocean. American Museum Novitates 2282: 1-19.

Received December 2006

Accepted March 2007

Published online April 2007