

## NOTES ON GEOGRAPHIC DISTRIBUTION

### **Amphibia, Anura, Hylidae, *Phyllomedusa azurea*: Distribution extension.**

Vitor Hugo M. do Prado

Rinneu E. Borges

Fernando R. da Silva

Thiago T. Tognolo

Denise de C. Rossa-Feres

Departamento de Zoologia e Botânica, Universidade Estadual Paulista (UNESP), Campus de São José do Rio Preto.  
Rua Cristóvão Colombo 2265, CEP 15054-000, São José do Rio Preto, SP, Brazil.  
E-mail: vitorbioibilce@yahoo.com.br

*Phyllomedusa azurea* Cope, 1862 (Figure 1), was recently revalidated and included in the species group of *Phyllomedusa hypochondrialis* (Daudin, 1800) by Caramaschi (2006). The geographic range of *P. azurea* was then published as the open formations of the Chacoan regions of Bolivia, Paraguay, and northern Argentina, and in the Pantanal and Cerrado regions of central Brazil (Caramaschi 2006; Frost 2007). Herein, we present the first record of *P. azurea* for the state of São Paulo, Brazil.




**Figure 1.** *Phyllomedusa azurea*. Arrows show three of the diagnostic characters of this species according to Caramaschi (2006). A, narrow white stripe on the upper lip, not reaching the lower eyelid; B, inguinal region and sides of members with black transversal bars over an orange or red background; C, wide green stripe along the dorsal surface of thighs.

Two calling males were heard and one was collected in the municipality of União Paulista ( $20^{\circ}55'54''$  S,  $49^{\circ}56'41''$  W) by R. E. Borges and

F. R. Silva in 25 October 2006. The specimen was found perching on vegetation (Poaceae: *Brachiaria* sp.) within a permanent pond in a pasture area, ca. 350 m far from a semideciduous forest fragment. The pond has  $3,118 \text{ m}^2$  of total area, 90 cm of maximum depth, and 70 % of plant cover (Poaceae). This record now represents the southeastern limit of the distribution of *P. azurea* (Figure 2), which is ca. 360 km from the nearest locality previously known for this species (Serranópolis, in the state of Goiás, Brazil  $18^{\circ}17'37''$  S,  $51^{\circ}58'10''$  W).

The voucher specimen was identified according to the diagnosis presented by Caramaschi (2006) and deposited in the Coleção do Departamento de Zoologia e Botânica (DZSJR-16604) housed at UNESP, São José do Rio Preto, São Paulo, Brazil. According to Caramaschi (2006), the revalidation of *P. azurea* and the recognition of *P. nordestina* Caramaschi, 2006, both previously identified under the name *P. hypochondrialis*, demonstrate that three morphologically similar species are associated to three of the great morphoclimatic domains of South America (*sensu* Ab'Sáber 1977): *P. hypochondrialis* with Amazonian distribution, *P. nordestina* associated to the Caatinga region, and *P. azurea* occurring in the Cerrado-Caatinga-Chaco complex (Cei 1980; Caramaschi 2006; Frost 2007). The record of *P. azurea* in the northwestern region of São Paulo evidences the necessity of additional anuran surveys, even in a well studied region (i.e., previously investigated in Vizotto 1967; Bernarde and Kokubum 1999; Rossa-Feres and Jim 2001; Vasconcelos and Rossa-Feres, 2005; Santos et al. 2007; Santos and Rossa-Feres 2007).

## NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** Distribution map for *Phyllomedusa azurea* with the new record for the municipality of União Paulista, State of São Paulo, Brazil (red dot).

### Acknowledgements

The authors are grateful to Célio F. B. Haddad for the identification of the specimen collected, and to FAPESP (grant 06/51534-1) for financial support.

### Literature cited

- Ab'Sáber, A. N. 1977. Os domínios morfoclimáticos na América do Sul, primeira aproximação. *Geomorfologia* 52: 1-21.
- Bernarde, P. S. and M. N. C. Kokubum. 1999. Anurofauna do município de Guararapes, estado de São Paulo, Brasil (Amphibia, Anura). *Acta Biologica Leopoldensia* 21: 89-97.
- Caramaschi, U. 2006. Redefinição do grupo de *Phyllomedusa hypochondrialis*, com redescrição de *P. megacephala* (Miranda-Ribeiro, 1926), revalidação de *P. azurea* Cope, 1862, e descrição de uma nova espécie (Amphibia, Anura, Hylidae). *Arquivos do Museu Nacional* 64(2): 159-179.
- Cei, J. M. 1980. Amphibians of Argentina. *Monitore Zoologico Italiano* (n. s.) Monograph 2: 1-609.
- Frost, D. R. 2007. Amphibian Species of the World: an online reference. Version 5.0. Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.php>. American Museum of Natural History, New York. Captured on August 2007.
- Rossa-Feres, D. C. and J. Jim. 2001. Similaridade no sítio de vocalização em uma comunidade de anfíbios anuros na região noroeste do Estado de São Paulo, Brasil. *Revista Brasileira de Zoologia* 18(2): 439-454.
- Santos, T. G. and D. C. Rossa-Feres. 2007. Similarities in calling site and advertisement call among anuran amphibians in Southeastern Brazil. *South American Journal of Herpetology* 2(1): 17-30.
- Santos, T. G., D. C. Rossa-Feres, and L. Casatti. 2007. Diversidade e distribuição espaço-temporal de anuros em região com pronunciada estação seca no sudeste do Brasil. *Iheringia (Zoologia)* 97(1): 37-49.
- Vasconcelos, T. S. and D. C. Rossa-Feres. 2005. Diversidade, distribuição espacial e temporal de anfíbios anuros (Amphibia, Anura) na região noroeste do estado de São Paulo, Brasil. *Biota Neotropica* 5(2): 1-14.
- Vizotto, L. D. 1967. Desenvolvimento de anuros da região norte-occidental do estado de São Paulo. Faculdade de Filosofia, Ciências e Letras, Zoologia nº especial: 1-161.

Received August 2007

Accepted January 2008

Published online March 2008

## NOTES ON GEOGRAPHIC DISTRIBUTION

### ERRATUM

#### **Amphibia, Anura, Hylidae, *Phyllomedusa azurea*: Distribution extension.**

Vitor Hugo M. do Prado  
Rinneu E. Borges  
Fernando R. da Silva  
Thiago T. Tognolo  
Denise de C. Rossa-Feres

*Departamento de Zoologia e Botânica, Universidade Estadual Paulista (UNESP), Campus de São José do Rio Preto.  
Rua Cristóvão Colombo 2265, CEP 15054-000, São José do Rio Preto, SP, Brazil.  
E-mail: vitorbioibilce@yahoo.com.br*

#### **Acknowledgements**

The authors are grateful to Célio F. B. Haddad for the identification of the specimen collected, and to FAPESP (grants 04/04820-3 and 06/51534-1) for financial support.

We apologize and are grateful for your comprehension.

The authors.

*Erratum published online April 2008*

## NOTES ON GEOGRAPHIC DISTRIBUTION

### ERRATUM

#### **Amphibia, Anura, Hylidae, *Phyllomedusa azurea*: Distribution extension.**

Vitor Hugo M. do Prado  
Rinneu E. Borges  
Fernando R. da Silva  
Thiago T. Tognolo  
Denise de C. Rossa-Feres

*Departamento de Zoologia e Botânica, Universidade Estadual Paulista (UNESP), Campus de São José do Rio Preto.  
Rua Cristóvão Colombo 2265, CEP 15054-000, São José do Rio Preto, SP, Brazil.  
E-mail: vitorbioibilce@yahoo.com.br*

#### **Acknowledgements**

The authors are grateful to Célio F. B. Haddad for the identification of the specimen collected, and to FAPESP (grants 04/04820-3 and 06/51534-1) for financial support.

We apologize and are grateful for your comprehension.

The authors.

*Erratum published online April 2008*