

NOTES ON GEOGRAPHIC DISTRIBUTION

**Pisces, Melamphaidae, *Poromitra capito* Goode & Bean, 1883 and *Melamphaes simus* Ebeling, 1962: Distribution extension to the western South Atlantic Ocean.**

**Paula N. Campos  
Ana Cristina T. Bonecker  
Márcia S. de Castro**

*Universidade Federal do Rio de Janeiro, Instituto de Biologia, Departamento de Zoologia.  
CCS, Bloco A, Ilha do Fundão. CEP 21941-590. Rio de Janeiro, RJ, Brazil. E-mail: polinhacampos@yahoo.com.br*

The family Melamphaidae includes 36 species in five genera (Nelson 2006) distributed throughout the meso and bathypelagic oceanic zones of the world (Garcia and Morgan 2002). According to Kotlyar (2004) it is the largest and the most diverse family of the stephanoberycoid fishes. This family has three possible pigmentation patterns (Frias-Torres 2006) and different ornaments in the head (spines and ridges), that distinguish the genera (Sandknop and Watson 1996). *Poromitra* has crestlike head ridges and many spines internally and on the operculum, while *Melamphaes* is recognized by the pigment characters (Sandknop and Watson 1996).

Only seven Melamphaidae species were recorded from the western South Atlantic Ocean (Kotlyar 2004) and among them only adults of *Poromitra crassiceps* (Günther, 1878) were recorded from

Brazil (Menezes et al. 2003). Therefore, this study attempted to identify the larval stages of melamphoids occurring along the Brazilian coast (12° S – 23° S).

The specimens included in this study were collected in the oceanic regions of the Brazilian coast, between Bahia and Rio de Janeiro (12° S – 23° S). All stations were located beyond the 1,000 m isobath. Larvae were collected using a bongo net (330- and 500 µm mesh size) that was towed obliquely from the maximum depth of 200 m to the surface; and by vertical hauls from the maximum depth of 1,000 m, using a cylindrical-conical net (500 µm mesh size). Samples were immediately preserved in a borax-buffered solution of 4 % formalin. More information about the study area and sampling methodology are described in Table 1.

**Table 1.** Study area data, sampling methodology and Melamphaidae larvae standard length (SL).

Species	DZUFRJ	N	SL (mm)	Latitude / Longitude	Date	Local depth (m)	Sampling depth (m)	Net	Mesh size (µm)
<i>P. capito</i>	7512	1	20.8	21°54'36.5" / 39°45'20.0"	Oct/10/2001	1,700	0-1,000	cylindrical-conical	500
<i>M. simus</i>	15355	2	3.3-5.6	22°12'40.0" / 38°40'00.0"	Oct/31/1998	3,200	0-200	bongo	330
<i>M. simus</i>	5499	4	4.9-11.0	22°32'49.0" / 40°04'20.9"	Nov/07/2001	1,054	0-1,000	cylindrical-conical	500
<i>M. simus</i>	5500	1	3.6	22°31'58.9" / 40°02'53.4"	Nov/07/2001	1,066	0-1,000	cylindrical-conical	500
<i>M. simus</i>	5502	8	3.1-4.8	22°32'50.0" / 40°04'09.9"	Nov/06/2001	1,070	0-1,000	cylindrical-conical	500
<i>M. simus</i>	5501	4	3.6-5.4	22°32'50.0" / 40°04'09.9"	Nov/06/2001	1,100	0-1,000	cylindrical-conical	500
<i>M. simus</i>	118	1	3.3	21°14'52.7" / 40°03'10.7"	Feb/28/2002	1,252	0-1,100	cylindrical-conical	500
<i>M. simus</i>	119	2	4.6-7.3	21°14'52.7" / 40°03'10.7"	Feb/28/2002	1,252	0-1,100	cylindrical-conical	500

DZUFRJ, number of voucher species in the larval fish collection; N, number of specimens identified.


## NOTES ON GEOGRAPHIC DISTRIBUTION

Melamphaid larvae were totally sorted from the samples and measurements (standard length – SL) were made with a micrometer ruler (precision of 0.1 mm). Figures were made from the collected material and all specimens were deposited in the larval fish collection of the Zooplankton and Ichthyoplankton Integrated Laboratory of Federal University of Rio de Janeiro – Brazil (DZUFRJ).

A total of 23 Melamphaidae larvae were identified including two species: *Poromitra capito* Goode & Bean, 1883 and *Melamphaes simus* Ebeling, 1962. This was the first occurrence of these species from the Brazilian coast. According to published reports (Ebeling and Weed 1973; Keene et al. 1987) *Melamphaes simus* had been found previously in Atlantic, Indo-Pacific, and Central

North Pacific oceans; in the western North Atlantic, from south of Grand Bank to Bahamas and Gulf of México. The species *Poromitra capito* occurs in North Atlantic Ocean and in the western North Atlantic, from Flemish Cap to the Bahamas. Therefore, in this study both species had their distribution extended to the western South Atlantic Ocean and this new finding contributes to future zoogeographical studies.

In the study area *Poromitra capito* was collected only over the 1,000 m isobath (Table 1). The unique larvae identified is 20.8 mm long (SL) having 3 spines and 11 dorsal fin rays; and 10+22 gillrakers. It also has well developed preopercule, rostral spines, and small eyes in relation to head length (Figure 1).



**Figure 1.** Postflexion larvae of *Poromitra capito* larvae (DZUFRJ 7512).


The genus *Poromitra* was already represented along the northeastern Brazilian coast by the species *Poromitra crassiceps* (Günther, 1878) (McEachran and Fechhelm 1998; Menezes et al. 2003). According Frias-Torres (2006) the differences between these species are the number of gillrakers, pigmentation pattern, and the presence of many spines in the top of the head that are more developed in *P. capito*, while *P. crassiceps* has internarial spines well developed.

Twenty two larvae of *Melamphaes simus* Ebeling, 1962 were identified and all of them were collected between 21° and 22° (Table 1). According to Sandknop and Watson (1996) the

pigmentation pattern and the number of gillrakers are the main characteristics that separate the *Melamphaes* species.

In this study *Melamphaes* larvae varied from of 3.1 mm (preflexion) to 11 mm (postflexion) (Table 1). Larvae of this species have 3 spines and 15-16 rays in the dorsal fin and 4+11-12 gillrakers (Figure 2). The pigmentation pattern comprises peritoneal pigment and internal pigment over the brain during flexion stage. In this stage there is also a line of melanophores in the dorsal margin of the body from the beginning of the dorsal fin to the caudal peduncle and a small cluster of spots on caudal peduncle (Figure 2).

NOTES ON GEOGRAPHIC DISTRIBUTION



**Figure 2.** Preflexion larvae of *Melamphaes simus*; A, DZUFRJ 5502, flexion larvae; B and C, DZUFRJ 5499, 119, postflexion larvae; D, DZUFRJ 5499.

## NOTES ON GEOGRAPHIC DISTRIBUTION

All Melamphaidae larvae identified in this study were collected at stations located beyond the 1,000 m and 3,000 m isobaths (Table 1). According to the literature this family is a typical

member of the deep water fish fauna which often occur in waters over the continental slope (Sandknop and Watson 1996; Moore 2002; Frias-Torres 2006; Fahay 2007).

### Acknowledgments

We thank the team of Zooplankton and Ichthyoplankton Integrated Laboratory of Federal University of Rio de Janeiro for assistance in field surveys. Special thanks to Sérgio Bonecker for comments on manuscript. We are grateful to CENPES/PETROBRAS for financial support and for allowing publishing this study.

### Literature cited

- Ebeling, A. W. and W. H. Weed. 1973: Order Xenoberyces (Sthenoberyciformes); p. 397-478 *In* D. M. Cohen (ed.). Fishes of the Western North Atlantic. Memoir Sears Foundation for Marine Research 1(6): 698 p.
- Fahay, M. P. 2007. Early Stages of Fishes in the Western North Atlantic Ocean (Davis Strait, Southern Greenland and Flemish Cap to Cape Hatteras) Volume 2. Nova Scotia: Northwest Atlantic Fisheries Organization. 1696 p.
- Frias-Torres, S. 2006. Melamphaidae: Bigscale fishes or ridgeheads; p. 1067-1074 *In*: W.J. Richards (ed.). Early Stages of Atlantic Fishes: An Identification Guide for the Western Central North Atlantic. London: CRC Marine Biology Series.
- Garcia, M. L. and C. C. Morgan. 2002. *Poromitra crassiceps* (Teleostei, Melamphaidae) associated with the 500 fathoms fauna off Argentina. *Journal of Applied Ichthyology* 18: 216–218.
- Keene, M. J., R. H. Gibbs, Jr., and W. H. Krueger. 1987. Family Melamphaidae, Bigscales; p. 169-185 *In* R. H. Gibbs, Jr. and W. H. Krueger (ed.). Biology of midwater fishes of the Bermuda Ocean Acre. Smithsonian Contributions to Zoology 452.
- Kotlyar, A. N. 2004. Family Melamphaidae Gill 1893 - bigscales. Bulletin of the Southern California Academy of Sciences Annotated Checklists of Fishes 29.
- Menezes, N. A., P. A. Buckup, J. L. Figueiredo, and R. L. Moura. 2003. Catálogo das espécies de peixes marinhos do Brasil. São Paulo. Museu de Zoologia USP. 160 p.
- McEachran, J. D. and J. D. Fechhelm. 1998. Fishes of the Gulf of México. Myxiniiformes to Gasterosteiformes. Austin: University of Texas Press. viii + 1112 p.
- Moore, J. A. 2002. Order Stephanoberyciformes. Melamphaidae Bigscale fishes (ridgeheads); p 1162-1163 *In* K. E. Carpenter. The living marine resources of the Western Central Atlantic. Rome: FAO.
- Nelson, J. R. 2006. Fishes of the World. 4<sup>th</sup> edition. Hoboken, New Jersey: John Wiley & Sons. 600 p.
- Sandknop, E. M., and W. Watson. 1996. Melamphaidae: Bigscales; p. 692-711. *In* H. G. Moser (ed.). The early stages of fishes in the California Current region. CalCOFI Atlas 33. Lawrence: Allen Press.

Received May 2008

Accepted July 2008

Published online August 2008