

NOTES ON GEOGRAPHIC DISTRIBUTION

Reptilia, Squamata, *Polychrus* spp.: New record, range extensions, and distribution map in the state of Mato Grosso, Brazil.

Ricardo A. Kawashita-Ribeiro¹
Robson W. Ávila²

¹ Universidade Federal de Mato Grosso, Instituto de Biociências, Coleção Zoológica de Vertebrados. Avenida Fernando Corrêa da Costa s/n. CEP 78060-900. Cuiabá, MT, Brazil. E-mail: serpentesbr@gmail.com

² Universidade Estadual Paulista, Instituto de Biociências, Departamento de Parasitologia. Programa de Pós-Graduação em Biologia Geral e Aplicada. Distrito de Rubião Júnior s/n. CEP 18618-000. Botucatu, SP, Brazil.

The arboreal medium-sized lizard genus *Polychrus* consists of six species distributed throughout South and Central America (Peters and Donoso-Barros 1970). Three species occurs in Brazil: *Polychrus acutirostris*, *P. liogaster*, and *P. marmoratus*. The first is distributed in tropical open vegetation formations from the Chaco through central Brazilian Cerrado and the Caatinga (Vanzolini et al. 1980). The other two species inhabits forested areas, in which *P. marmoratus* is a widespread species, ranging from the Atlantic Forest in São Paulo and Pernambuco states to the Amazon, in the Amazonas, Amapá, Maranhão, Pará, and Roraima states, while *Polychrus liogaster* is restricted to southwestern Amazon and neighboring areas, in the states of Acre, Amazonas, and Rondônia (Ávila-Pires 1995).

The state of Mato Grosso is covered with three major Brazilian biomes, including the Pantanal, Cerrado, and Amazon. Surprisingly, there are published records only for *P. acutirostris*, and for a single locality: Chapada dos Guimarães (Cope 1887; Strüssmann 2000). *Polychrus marmoratus* was previously found in the state, in the municipality of Aripuanã (Mato Grosso 2000). However, this record was published only in a government technical report with limited access to the scientific community.

Herein we report for the first time the presence of *P. liogaster* in the state of Mato Grosso. Also, we present the only record for *P. marmoratus* and a distribution map for *P. acutirostris* in the state. Specimens were collected mainly during faunal rescues and monitoring plans for water dams in Mato Grosso. Voucher specimens were housed at

the Coleção Zoológica de Vertebrados do Instituto de Biociências da Universidade Federal de Mato Grosso (UFMT). *P. acutirostris*: Cáceres (UFMT 3443; 16°04' S, 57°41' W); Chapada dos Guimarães (UFMT 616-22, 718-9, 2288, 2307, 2410-1, 3442, 5754-73, and 6116-7; 14°52' S, 55°48' W); Jaciara (UFMT 2211; 15°57' S, 54°58' W); Nossa Senhora do Livramento (UFMT 016; 15°45' S, 56°20' W); Parque Nacional do Pantanal UFMT 6202; 17°26' S, 57°10' W); Poconé, Fazenda Santa Inês (UFMT 264 and 4935; 16°30' S, 56°45' W); Primavera do Leste (UFMT 6756-7; 15°19' S, 54°09' W); Rodovia Transpantaneira km 112 (UFMT 043; 17°16' S, 56°59' W); Santo Antônio do Leverger (UFMT 130 and 302; 15°52' S, 56°04' W). *P. liogaster*: Araputanga, PCH Ombreiras (UFMT 2888; 15°08' S, 58°54' W); Jauru, PCH Ombreiras (UFMT 2885-7 and 2889; 15°08' S, 58°54' W); Vale de São Domingos, UHE Guaporé (UFMT 896, 5021-2, 5448, 5884-94, and 6115; 15°00' S, 58°58' W). *P. marmoratus*: Aripuanã (UFMT 2992, 10°09' S, 59°27' W) (Figure 1).

The records here reported extend the known geographical distribution of *P. marmoratus* (Figure 2) ca. 1,144 km southeast from the nearest record, the municipality of Carajás, in the state of Pará. For *P. liogaster* (Figure 3), the geographical distribution is extended ca. 467 km east from the province of Sara, in Bolívia, and ca. 874 km southeast from the UHE Samuel, in the state of Rondônia (see Ávila-Pires for a taxonomic discussion). For *P. acutirostris* (Figure 4), the distribution in Mato Grosso presented in Figure 1 fills the gaps in the western Cerrado region of central Brazil.

NOTES ON GEOGRAPHIC DISTRIBUTION




Figure 1. Known geographic distribution of *Polychrus* spp. in the state of Mato Grosso, Brazil; triangles, *P. acutirostris*; square, *P. marmoratus*; circles, *P. liogaster*. Colors represent biomes: green, Amazon; gray, Cerrado; rose, transition zones; blue, Pantanal.



Figure 2. Adult male *P. marmoratus* collected in the municipality of Aripuanã. Photo by R. A. K. Ribeiro.

NOTES ON GEOGRAPHIC DISTRIBUTION



Figure 3. Adult male *P. liogaster* collected at the municipality of Jauru. Photo by R. W. Ávila.



Figure 4. Adult male *P. acutirostris* collected at the municipality of Chapada dos Guimarães. Photo by C. Strüssmann.

NOTES ON GEOGRAPHIC DISTRIBUTION

This study reaffirms the lack of basic herpetofaunal knowledge for some areas in central Brazil (see Colli et al. 2002). Mato Grosso is one of the largest states of the country and harbors a considerable biodiversity. However, there is a huge lack of information for parts of the state. For

example, the eastern region of Mato Grosso, which encompasses the Araguaia River basin, is poorly represented in zoological collections. Probably, the distribution of *Polychrus* species will be further enlarged with new herpetological surveys in those areas.

Acknowledgements

We thank to Reuber A. Brandão and one anonymous reviewer for critical reading of the manuscript and Marcos André de Carvalho for the access to the UFMT Zoological Collection. We also thank to Tami Mott for help with construct the distribution map in ArcView.

Literature cited

- Ávila-Pires, T. C. S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). Leiden: Zoolochische Verhandelingen. 706 p.
- Cope, E. D. 1887. Synopsis of the Batrachia and Reptilia obtained by H. H. Smith, in the province of Mato Grosso, Brazil. Proceedings of the American Philosophical Society 24: 44-60.
- Colli, G. R., R. P. Bastos, and A. F. B. Araújo. 2002. The Character and Dynamics of the Cerrado Herpetofauna; p. 223-241 In P. S. Oliveira and J. Marquis (ed.). The Cerrados of Brazil: ecology and natural history of a Neotropical Savanna. New York: Columbia University Press.
- Mato Grosso. Secretaria do Estado de Planejamento e Coordenação Geral do Estado de Mato Grosso (SEPLAN). 2000. Zoneamento Sócio-Econômico-Ecológico do Estado de Mato Grosso: diagnóstico sócio-econômico-ecológico e assistência técnica na formulação da 2^a aproximação. Projeto de Desenvolvimento Agroambiental do Estado de Mato Grosso (PRODEAGRO). Fauna. Parte 2: sistematização das informações temáticas, nível compilatório. Cuiabá: SEPLAN and BIRD. 154 p.
- Peters, J. A. and R. Donoso-Barros. 1970. Catalogue of the Neotropical Squamata. Part II. Lizards and Amphisbaenians. United States National Museum Bulletin 297: viii + 293 p.
- Strüessmann, C. 2000. Herpetofauna; p. 153-189 In C. J. R. Alho, P. N. Conceição, R. Constantino, T. Schlemmermeyer, C. Strüessmann, L. A. S. Vasconcellos, D. M. M. Oliveira, and M. Schneider (ed.). Fauna silvestre da região de Manso - MT. Brasília: Ministério do Meio Ambiente, Edições IBAMA and Centrais Elétricas do Norte do Brasil.
- Vanzolini, P. E., A. M. Ramos-Costa, and L. J. Vitt. 1980. Répteis das Caatingas. Rio de Janeiro: Academia Brasileira de Ciências. 161 p.

Received January 2008

Accepted August 2008

Published online September 2008