

NOTES ON GEOGRAPHIC DISTRIBUTION

New distribution and host records of Chalcidoidea (Insecta: Hymenoptera) from various parts of India

Ankita Gupta
J. Poorani

Project Directorate of Biological Control (ICAR)
P. B. No.2491, H.A. Farm post, Bellary Road, Hebbal, Bangalore 560 024, Karnataka, India.
080-23511982/98 -Ext.335. E-mail: ankitagupta9@rediffmail.com

New distribution and host records of 14 species from five families of the superfamily Chalcidoidea from various parts of southern and western India are presented below, based on the studies carried out at Project Directorate of Biological Control (PDBC), Bangalore. The specimens are deposited in the collections of PDBC which is a designated National Repository under the National Biodiversity Authority.

Ooencyrtus musa is reported for the first time from lepidopterous eggs on *Cassia auriculata*, *Pachyneuron groenlandicum* from indet. hemerobiid on *Chromolaena odorata* and *Pachyneuron solitarium* from *Lipaphis erysimi*. From Andhra Pradesh, *Anagyrus chrysos*, *Blepyrus insularis*, *Homalotylus indicus* and *Oricoruna arcotensis* are recorded. From Karnataka, *Metaphycus bolangerae*, *Prochiloneurus aegyptiacus*, *Neanastatus trochantericus*, *Mymar taprobanicum*, *Mangostigmus amraeus* and *Megastigmus viggianii* are recorded. From Maharashtra, *Sphegigaster brunneicornis* are recorded. The present study deals with six species of Encyrtidae, four species of Pteromalidae, two species of Torymidae and one species each of Mymaridae and Eupelmidae.

The study area comprises the states Andhra Pradesh, Karnataka, Tamil Nadu and Maharashtra. The annual mean temperature in these regions is between 16-28 °C. The studies were carried out using S8APO stereozoom microscope and identification was done with the aid of recent literature as well as data obtained from Universal Chalcidoidea database (Noyes 2007). Specimen collection was done by using sweep-net, aspirator,

and yellow pan traps set up in the collection sites. Along with this various mealybugs infested plant parts, parasitized eggs on pods, parasitized pupae, scale infested leaves, leaves with galls were also collected and kept in jars for emergence of parasitoids. The collected specimens were then dry mounted and labelled according to the standard procedure (Noyes 1982). Collection site, date, host and collector's name and other details are provided in the subhead for each species.

Family Encyrtidae

1. *Anagyrus chrysos*

Anagyrus chrysos Noyes & Hayat, 1994: 80, 176-178, F, M. Holotype F: India, Lucknow.

Material studied: 7F, 5M, Andhra Pradesh, Guntoor, ex. indet. mealybugs on cotton, 02 December 2007, Naveen.

Distribution: India: Andhra Pradesh (New record), Bihar, Karnataka, Madhya Pradesh, Maharashtra, Tamil Nadu and Uttar Pradesh; Malaysia.

2. *Blepyrus insularis*

Encyrtus? insularis Cameron, in Blackburn & Cameron, 1886: 243-245, M. Holotype F: Hawaii (Type lost).

Blepyrus insularis (Cameron): Timberlake, 1922: 167.

Blepyrus longiscapus Kaul & Agarwal, 1986: 50. Lectotype F: India, Pantnagar. Synonymy by Hayat, 1999: 360.

Blepyrus annulobliquus Kaul & Agarwal, 1986: 13, 50-52. Holotype F: India, Madhya Pradesh.

Material studied: 1F, Andhra Pradesh: Rajamundry, ex. indet. mealybugs on bean, 30 November 2007, Lalitha.

Distribution: India: Andhra Pradesh (new record), Delhi, Karnataka, Kerala, Goa, Madhya Pradesh,

NOTES ON GEOGRAPHIC DISTRIBUTION

Maharashtra, Orissa, Tamil Nadu, Uttaranchal, Uttar Pradesh and West Bengal; Sri Lanka; Malaysia.

3. *Homalotylus indicus* (Figure 1)

Neoenasioidea indica Agarwal, 1966: 73-74, F, M. Holotype F: India, Aligarh.

Prochiloneurus nipaecocci Subba Rao, 1967: 1, F. Holotype F: India, Delhi. Synonymy by Anis & Hayat, 1998: 208.

Homalotylus indicus (Agarwal): Hayat, Alam & Agarwal, 1975: 65, 67, F, M. India, Aligarh & Shoranur.

Echthrolepis shafii Fatima & Agarwal, 1994: 281, F. Holotype F: India, Aligarh. Synonymy by Anis & Hayat, 1998: 208.

Material studied: 1F, 1M, Andhra Pradesh: Guntoor, ex. indet. mealybugs on cotton, 02 December 2007, Ankita.

Distribution: Andhra Pradesh (New record), Delhi, Kerala, Rajasthan, Uttar Pradesh and Tamil Nadu.



Figure 1: *Homalotylus indicus*

4. *Metaphycus bolangerae*

Metaphycus bolangerae Hayat, in Hayat *et al.*, 2003: 313-314, F, M. Holotype F: India, Bangalore.

Material studied: 2F, Karnataka: Bangalore, ex. *Saissetia* sp., 14 October 2007, Ankita.

Distribution: India: Karnataka (New host record on *Saissetia* sp.). It was recorded earlier from *Ceroplastes actiniformis* (Homoptera: Coccidae) on sandalwood.

5. *Ooencyrtus musa*

Ooencyrtus musa Huang & Noyes, 1994: 15, 77-78, Holotype F: India, Karnataka, Bannerghatta.

Material studied: 2F, 1M, Karnataka, ex. indet. lepidopterous eggs on *Cassia auriculata*, 20 October 2007, Ankita.

Host: For the first time this species is recorded from any known host *viz.* indet. lepidopterous eggs on *Cassia auriculata*.

Distribution: India: Karnataka.

6. *Prochiloneurus aegyptiacus* (Mercet)

Achrysopophagus aegyptiacus Mercet, 1929: 5, 360.

Prochiloneurus aegyptiacus (Mercet): Niyazov, 1968: 899.

Prochiloneurus valparianus Mani & Kaul, 1974: 66-68, F. Holotype F: India, Anamalai Hills. Synonymy by Fatima & Shafee, 1994: 24.

Prochiloneurus indicus Shafee, Alam & Agarwal, 1975: 49-50, F, M. Holotype F: India, Coimbatore. Synonymy by Fatima & Shafee, 1994: 24.

Material studied: 1F, Karnataka: Hessarghatta, ex. indet. mealy bugs on pomelo, 18 July 2007, Ankita.

Distribution: India: Karnataka (new record), Kerala, Tamil Nadu and Andhra Pradesh.

Family Eupelmidae

7. *Neanastatus trochantericus*

Neanastatus trochantericus Gahan, 1919: 56, 520. F, India, Tamil Nadu.

Neanastatus trochantericus Gahan: Narendran, 1996: 81.

Material studied: 1M, Karnataka: Hessarghatta, sweeping on grasses, 19 September 2007, Ankita.

Distribution: India: Karnataka (new record) and Tamil Nadu.

Family Mymaridae

8. *Mymar taprobanicum*

Mymar taprobanicum Ward, 1875: 11, 197. F, Srilanka.

Mymar tyndalli Girault, 1912: 1, 168. Holotype F, Australia, Queensland.

Mymar antillanum Dozier, 1937: 21, 130. Puerto Rico.

Mymar indica Mani, 1942: 4, 160. India.

Oglobliniella aegyptiaca Soyka, 1950: 34, 131. Egypt.

NOTES ON GEOGRAPHIC DISTRIBUTION

Material studied: 1F on slide, Karnataka: Bangalore, yellow pan trap, 05 December 2007, Ankita.

Distribution: India: Karnataka (new record), Delhi, Madhya Pradesh and Rajasthan.

Family Pteromalidae

9. *Oricoruna arcotensis*

Pachycrepoides arcotensis Mani & Kurian, 1953: 15, 11. F, India, N. Arcot. (Lost).

Pachycrepoides coorgensis Mani & Kurian, 1953: 15, 10. M, F, India, Bangalore (lost).

Material studied: 1F, Andhra Pradesh: Tirupathi vet. College, Resting on bamboo leaf, 16 January 2007, Ankita.

Distribution: Andhra Pradesh (new record), Bihar, Jharkhand, Karnataka and Tamil Nadu.

10. *Pachyneuron groenlandicum*

Pteromalus groenlandicus Holmgren, 1872: 29, 100, M, F. Lectotype F: Greenland.

Pachyneuron karnalensis Mani, 1939: 1, 85. F, India, Karnal.

Pachyneuron umbratum Delucchi, 1955: 38, 132-133. F, Switzerland.

Pachyneuron bakrotu, Mani & Saraswat, 1974: 3, 102. F, India, Dalhousie.

Host: Hemerobiidae is a new host record for this species. Earlier records were from *Ischiodon* sp. on *Cassia* sp. and syrphids associated with *Brachycaudus* sp.

Material studied: 1F; Karnataka: Bangalore, ex. Indet. Hemerobiid on *Chromolaena odorata*, 15 September 2006, Ankita.

Distribution: India: Delhi, Haryana, Himachal Pradesh, Kerala, Karnataka, Orissa, Tamil Nadu, Jammu & Kashmir; Europe.

11. *Pachyneuron solitarium*

Pachyneuron coccorum Linnaeus, 1758. Syst. Nat. 10th Ed.

Pachyneuron solitarius Hartig, 1838: 1, 250. Lectotype F: Germany.

Material studied: 1F, Maharashtra: Akola, ex. *Lipaphis erysimi* Kalt, 23 January 2006, Joshi.

Host: This species has a wide range of hosts, hyperparasitising aphids, coccids, etc. but no host data was available from India. For the first time it is recorded from *Lipaphis erysimi* in India.

Distribution: India: Maharashtra (new record) and Kerala; Japan; Europe.

12. *Sphegigaster brunneicornis*

Trigonogastra brunneicornis Ferrière, 1930: 21, 356-357. Lectotype F: Sri Lanka, Peradeniya.

Material studied: 1F, Tamil Nadu, ex. Pupae of stem fly, 10 October 2007, Tamil Nadu Agricultural University collection.

Distribution: India: Tamil Nadu (new record) and Kerala; Sri Lanka.

Family Torymidae

13. *Mangostigmus amraeus*

Neomegastigmus amraeus Chandy Kurian, 1953: 2, 241. Holotype F: India.

Mangostigmus amraeus (Chandy Kurian): Boucek, 1986.

Mangostigmus amraeus (Chandy Kurian): Narendran, 1994: 32-33.

Material studied: 6F, 1M, Karnataka: Thalaghatapura, ex. Galls on *Mangifera indica*, Ankita.

Distribution: India: Karnataka (new record), Uttar Pradesh; Malaysia; Thailand.

14. *Megastigmus viggianii* (Figure 2)

Megastigmus viggianii Narendran & Sureshan, 1988: 45, 38-41, M, F. Holotype F: India.

Megastigmus viggianii Narendran & Sureshan: Narendran, 1994: 29-30.



Figure 2: *Megastigmus viggianii*

Material studied: 2F, 2M, Karnataka: Yana, ex. Leaf galls on indet tree, 05 November 2007, Ankita. 5F, Karnataka: Hessarghatta, ex. Leaf

NOTES ON GEOGRAPHIC DISTRIBUTION

galls on *Eucalyptus* sp. infested with *Leptocybe invasa* Fisher and La Salle, 09 June 2008, Ankita.
Distribution: Karnataka (new record) and Kerala.

Metaphycus bolangerae is reported from *Saissetia* sp. which is a new host record. Earlier record was from *Ceroplastes actiniformis* (Homoptera: Coccidae) on sandalwood. The host of *Ooencyrtus musa*, which is endemic to Karnataka, was not known. For the first time, this species has been found parasitizing lepidopterous eggs on *Cassia auriculata*. The known host range of *Pachyneuron groenlandicum* includes dipterans, viz., *Ischiodon* sp. on *Cassia* sp. and syrphids associated with *Brachycaudus* sp. We

recorded it from an indeterminate hemerobiid (Neuroptera) on *Chromolaena odorata*. This species occurs throughout the country. *Pachyneuron solitarium* was found to be associated with various aphids and coccids from various parts of the world, but for the first time it is recorded from *Lipaphis erysimi* in India. This species was prevalent in south India but our study reports that its range extends to western India also. *Mangostigmus amraeus* prevalent in North India has been found to exist in south India also. *Megastigmus viggianii* was found to be associated with eucalyptus galls infested by the gall wasp, *Leptocybe invasa* for the first time.

Acknowledgements

This work was carried out under the Network Project on Insect Biosystematics funded by the Indian Council of Agricultural Research, New Delhi.

Literature cited

- Agarwal, M. M. 1966. Three undescribed genera and species of Encyrtidae (Hymenoptera-Chalcidoidea) parasitic on coccids. Proceedings of the Indian Academy of Sciences (B) 63(2): 73.
- Anis, S. B. and M. Hayat. 1998. The Indian species of *Homalotylus* (Hymenoptera: Encyrtidae). Oriental Insects 32: 191-218.
- Boucek, Z. 1986. Taxonomic study of chalcidoid wasps (Hymenoptera) associated with gall midges (Diptera: Cecidomyiidae) on mango trees. Bulletin of Entomological Research 76 (3): 393-407.
- Blackburn, T. and P. Cameron. 1886. On the Hymenoptera of the Hawaiian Islands. Memoirs of the Manchester Literary and Philosophical Society 10(3): 243.
- Chandy, K. 1953. Description of a new chalcid parasite of midge of echinate gall on mango leaf. Agra University Journal of Research (Science) 2: 241.
- Delucchi, V. 1955. Beiträge zur Kenntnis der Pteromaliden (Hym., Chalcidoidea). I. Zeitschrift für Angewandte Entomologie 38: 132.
- Dozier, H. L. 1937. Descriptions of miscellaneous chalcidoid parasites from Puerto Rico (Hymenoptera). Journal of Agriculture of the University of Puerto Rico 21: 130.
- Fatima, A. and M. M. Agarwal. 1994. Two new species of Encyrtidae (Hymenoptera: Chalcidoidea) from Aligarh (India). Journal of Entomological Research 17(4): 279-282.
- Fatima, A. and S. A. Shafee. 1994. Studies on the taxonomy of the Indian encyrtids (Hymenoptera: Encyrtinae). Aligarh Muslim University Publications (Zoological series) on Indian Insect types 15: 141.
- Ferrière, C. 1930. Notes on Asiatic Chalcidoidea. Bulletin of Entomological Research 21(3): 356.
- Girault, A. A. 1912. Australian Hymenoptera Chalcidoidea. II. The family Mymaridae with descriptions of new species. Memoirs of the Queensland Museum 1: 168.
- Gahan, A.B. 1919. Report on a small collection of Indian parasitic Hymenoptera. Proceedings of the United States National Museum 56: 520.
- Hartig, T. 1838. Über den Raupenfrass im Königlichen Charlottenburger Forste unfern Berlin, während des Sommers 1837. Jahresberichte über die Fortschritte der Forstwissenschaft und Forstlichen Naturkunde im Jahre 1836 und 1837 nebst Original-Abhandlungen aus dem Gebiete und Cameralisten 1(2): 250.
- Hayat, M. 1999. Taxonomic notes on Indian Encyrtidae (Hymenoptera: Chalcidoidea) - V. Oriental Insects 33: 349-407.
- Hayat, M., S. M. Alam and M. M. Agarwal. 1975. Taxonomic survey of encyrtid parasites (Hymenoptera: Encyrtidae) in India. Aligarh Muslim University Publications (Zoological series) on Indian Insect Types 9 (1-3): 1-112.

NOTES ON GEOGRAPHIC DISTRIBUTION

- Hayat, M., T. C. Narendran, O. K. Remadevi and S. Manikandan. 2003. Parasitoids (Hymenoptera: Chalcidoidea; Ceraphronoidea) reared mainly from Coccoidea (Homoptera) attacking sandalwood, *Santalum album* L. *Oriental Insects* 37: 309-334.
- Huang, D. W. and J. S. Noyes. 1994. A revision of the Indo-Pacific species of *Ooencyrtus* (Hymenoptera: Encyrtidae), parasitoids of the immature stages of economically important insect species (mainly Hemiptera and Lepidoptera). *Bulletin of the Natural History Museum (Entomology Series)* 63(1): 77-78.
- Holmgren, A.E. 1872. Insekter från Nordgrönland, samlade af Prof. A.E. Nordensköld år a870. Granskade Öfversigt af Kongl. Vetenskaps-Akademiens Förhandlingar 1872 (6): 100.
- Kaul, K. and M. M. Agarwal. 1986. Taxonomic studies on encyrtid parasitoids (Hymenoptera: Chalcidoidea) of India (with XVII plates). Aligarh Muslim University Publication, Zoological Series on Indian Insect Types 13: 50-52.
- Linnaeus, C. 1758. *Systema Naturae, sive regna tria naturae systematice proposita per classes, ordines, genera et species*. Holmiae (Stockholm). 824 p.
- Mani, M.S. 1939. Descriptions of new and new records of some known chalcidoid and other hymenopterous parasites from India. *Indian Journal of Entomology* 1:85.
- Mani, M. S., O. P. Dubey, B.K. Kaul and G. G. Saraswat. 1974. Descriptions of some new and new records of some known Chalcidoidea (Hymenoptera) from India. *Memoirs of the School of Entomology, St. John's College, Agra* 3: 103-104.
- Mani, M.S. 1942. Studies on Indian parasitic Hymenoptera II. *Indian Journal of Entomology* 4(2): 160.
- Mani, M.S. and C. Kurian. 1953. Descriptions and records of chalcids from India. *Indian Journal of Entomology* 15(1): 10-11.
- Mercet, R.G. 1929. Los generos *Prochiloneurus* y *Achrysoophagus*. *Eos, Revista Española de Entomología, Madrid* 5: 360.
- Narendran, T. C. and P. M. Sureshan. 1988. A contribution to our knowledge of Torymidae of India (Hymenoptera: Chalcidoidea). *Bollettino del Laboratorio di Entomologia Agraria 'Filippo Silvestri', Portici* 45: 38-41.
- Narendran, T. C. 1994. Torymidae and Eurytomidae of Indian subcontinent (Hymenoptera: Chalcidoidea) *Zoological Monograph, Department of Zoology, University of Calicut, Kerala, India*, 500pp.
- Narendran, T.C. 1996. Alpha systematics of some Eupelmidae (Hymenoptera: Chalcidoidea) from India. *Entomon* 21(1): 80-81.
- Niyazov, O.D. 1968. The role of the temperature factor in the parasitic activity of *Anagyrus pseudococci* (Gir.) (Hymenoptera, Encyrtidae). *Izvestiya Akademii Nauk Turkmenskoy SSR (Seriya Biologicheskikh Nauk)* (1): 79-82.
- Noyes J. S. 1982. Collecting and preserving chalcid wasps (Hymenoptera: Chalcidoidea). *Journal of Natural History* 16: 315-334.
- Noyes, J. S. 2007. Universal Chalcidoidea database. Electronic Database accessible at <http://internt.nhm.ac.uk/jdsml/perth/chalcidoids>. Captured on 05 July 2007.
- Noyes, J. S. and M. Hayat. 1994. Oriental mealybug parasitoids of the Anagyrini (Hymenoptera: Encyrtidae). CAB International, Oxon, UK. 554p.
- Soyka, W. 1950. New and little known alaptids and mymarids from Egypt. *Bulletin de la Société Fouad Ier d'Entomologie, Le Caire* 34:131.
- Shafee, S. A., S. M. Alam and M. M. Agarwal. 1975. Taxonomic surveys of encyrtid parasites (Hymenoptera: Encyrtidae). Aligarh Muslim University Publications (Zoological series) on Indian Insect types 10 (3): 1-125.
- Subba Rao, B.R. 1967. Description of some new species of encyrtids from India. *Bulletin of Entomology* 8(1): 1.
- Timberlake, P.H. 1922. Notes on the identity and habits of *Blepyrus insularis* Cameron (Hymenoptera, Chalcidoidea). *Proceedings of the Hawaiian Entomological Society* 5(1): 167-173.
- Ward, A.O. 1875. Description of a new species of Proctotrypidae from Ceylon. *Entomologist's Monthly Magazine* 11:197.

Received May 2008

Accepted September 2008

Published online October 2008