

NOTES ON GEOGRAPHIC DISTRIBUTION

Amphibia, Anura, Bufonidae, *Melanophrynniscus devincenzi*: First record for Paraguay and geographic distribution map

Katia Airaldi ¹
Diego Baldo ^{2,3}
Esteban Orlando Lavilla ²

¹ Universidad Nacional de Asunción, Facultad de Ciencias Exactas y Naturales, Departamento de Biología.
Ruta Mariscal Estigarribia km.11 (Campus Universitario), San Lorenzo, Paraguay.

² Instituto de Herpetología, Fundación Miguel Lillo – CONICET.
Miguel Lillo 251. 4000 – San Miguel de Tucumán, Argentina.

³ Universidad Nacional de Misiones, Departamento de Genética, Facultad de Ciencias Exactas, Químicas y Naturales.
Félix de Azara 1552, (3300) Posadas, Misiones, Argentina. E-mail: diegobaldo@gmail.com

The red-belly toad *Melanophrynniscus devincenzi* Klappenbach, 1968 belongs to the *M. tumifrons* group, occurring in Misiones province (Argentina), state of Rio Grande do Sul (Brazil), and northern Uruguay (Zanella et al. 2007). Brusquetti and Lavilla (2006) speculated about its presence in Paraguay, due to the proximity of populations in Misiones province and the continuity of available habitats. Herein, we present the first record of *M. devincenzi* for Paraguay, extending its known distribution range ca. 180 km northwards from Puerto San Juan, Misiones, Argentina (Baldo and Basso 2004), and

we provide a geographic distribution map with all known localities for this species.

The new record of *M. devincenzi* corresponds to a male collected near to the Pirapo'i stream ($25^{\circ}55' S$, $56^{\circ}09' W$, 260 m a.s.l.), Finca 2400 of Naturaleza Pura S.A., Reserva de Recursos Manejados del Ybyturuzú, Distrito Independencia, Guairá Department, Paraguay, on October 24 2008. The specimen is deposited in the Zoology Collection of the Facultad de Ciencias Exactas y Naturales, Universidad Nacional de Asunción, Asunción, Paraguay (CZCEN049) (Figure 1).




Figure 1. Dorsal (A) and ventral (B) views of *Melanophrynniscus devincenzi*, adult male (CZCEN 049), from Pirapo'i stream, Reserva de Recursos Manejados del Ybyturuzú, Guairá Department, Paraguay.

NOTES ON GEOGRAPHIC DISTRIBUTION

The Pirapo'i stream is a tributary of the Tebicuary-mi River that originates on the slopes of Tres Candú hill (842 m a.s.l.), in the Ybyturuzú mountain systems (Figure 2). This region belongs to the Atlantic Forest ecoregion (Dinerstein et al. 1995; Olson et al. 2001).



Figure 2. View of the Tres Candú hill, Ybyturuzú mountain systems, Paraguay.

Melanophrynniscus devincenzi is considered as “Endangered” by the IUCN, because its area of occupancy is less than 500 km², its distribution is severely fragmented, and there is continuing decline in the extent and quality of its habitat in Argentina and Uruguay (Lavilla et al. 2004). However, the recent record of this species in Rio Grande do Sul, Brazil (Zanella et al. 2006), and the present one for Paraguay largely increases its

known and potential distribution range (Figure 3), which would allow a reassessment of its conservation status.



Figure 3. Geographic distribution of *Melanophrynniscus devincenzi*. New record from Paraguay (asterisk); historical records (black circles) (Klappenbach 1968; Baldo and Basso 2004; Zanella et al. 2007), type locality (white circle) (Klappenbach 1968).

Acknowledgements

KA acknowledges *Naturaleza Pura S.A.* by the authorization and logistic support during her permanency in the field, and *Dirección de Investigaciones de la Universidad Nacional de Asunción* for financial support. We thank Claudio Bortero for his comments and assistance with the English version. DB acknowledges ANPCyT for Grants PICT-O (37035), PICT (16-35045), and PICT (06-233).

Literature cited

Baldo, D. and N. G. Basso. 2004. A new species of *Melanophrynniscus* Gallardo, 1961 (Anura: Bufonidae), with comments on the species of the genus reported for Misiones, northeastern Argentina. Journal of Herpetology 38(3): 393-403.

Brusquetti, F. and E. O. Lavilla. 2006. Lista comentada de los anfibios de Paraguay. Cuadernos de Herpetología 20(2): 3-79.
Dinerstein, E., D. M. Olson, D. J. Graham, A. L. Webster, S. A. Primm, M. P. Bookbinder, and G.

NOTES ON GEOGRAPHIC DISTRIBUTION

- Ledec. 1995. A conservation assessment of the terrestrial ecoregions of Latin America and the Caribbean. Washington: The World Bank and World Wildlife Foundation. 129 p.
- Klappenbach, M. A. 1968. Notas herpetológicas IV. El género *Melanophryniscus* (Amphibia, Salientia) en el Uruguay, con la descripción de dos nuevas especies. Comunicaciones Zoológicas del Museo de Historia Natural de Montevideo 9(118): 1-12.
- Lavilla, E., D. Baldo, and J. Langone 2004. *Melanophryniscus devincenzi*. In IUCN 2008. 2008 IUCN Red List of Threatened Species. Electronic Database accessible at www.iucnredlist.org. Captured on 15 February 2009.
- Olson, D. M., E. Dinerstein, E. D. Wikramanayake, N. D. Burgess, G. V. N. Powell, E. C. Underwood, J. A. D'amico, I. Itoua, H. E. Strand, J. C. Morrison, C. J. Loucks, T. F. Allnutt, T. H. Ricketts, Y. Kura, J. F. Lamoreux, W. W. Wettenberg, P. Hedao, and K. R. Kassem. 2001. Terrestrial ecoregions of the world: a new map of life on earth. Bioscience 51(11): 933-938.
- Zanella, N., C. S. Busin, A. Giusti, L. Crestani, and R. S. Oliveira. 2007. Amphibia, Anura, Bufonidae, *Melanophryniscus devincenzi*: First record for Brazil. Check List 3(2): 104.

Received March 2009

Accepted May 2009

Published online July 2009