

NOTES ON GEOGRAPHIC DISTRIBUTION

Mammalia, Chiroptera, Phyllostomidae, *Vampyrum spectrum* (Linnaeus, 1758): First record for the state of Rondônia, Brazil, and new prey records

Danayre Santana Discher ¹

Paulo Sérgio Bernarde ²

Kátia Gomes Facure ³

¹ Faculdade de Ciências Biomédicas de Cacoal - Facimed.
Av. Cuiabá, 3087, Jardim Clodoaldo. CEP 78976-005. Cacoal, Rondônia, Brazil.

² Universidade Federal do Acre - UFAC, Centro Multidisciplinar.
Campus Floresta, CEP 69980-000. Cruzeiro do Sul, Acre, Brazil. E-mail: SnakeBernarde@hotmail.com

³ União Educacional Minas Gerais - UNIMINAS.
Avenida dos Vinhedos, 1200, Morada da Colina. CEP 38411-159. Uberlândia, Minas Gerais, Brazil.

Vampyrum spectrum (Linnaeus, 1758) is the largest bat (adults 150–200 g) in the New World with up to one meter wingspan (Navarro and Wilson 1982). Geographic distribution of this species extends throughout the forested regions from Central Brazil, Bolivia, and Peru until Southern Mexico, also occurring in Guyana, Suriname, and Trinidad (Navarro and Wilson 1982; Emmons and Feer 1997; Nowak 1999). In Brazil, this species has been reported to Acre, Amazonas, Amapá, Mato Grosso, Pará, Piauí, Roraima, and Tocantins states (Gregorin et al. 2008; Nogueira et al. 2007).

In spite of its wide distribution, this species is poorly represented in zoological collections (Nogueira et al. 2007) and available information on its life history is based on a few specimens (Bonato et al. 2004; Vargas-Spinosa et al. 2004; Acosta and Azurduy 2006). Similarly to other large Microchiropteran species, *V. spectrum* is known to be predominantly carnivorous, feeding on a wide range of prey, such as birds, mice, and other bats (Vehrencamp et al. 1977; Bonato et al. 2004). In this note we present the first record of *V. spectrum* for the state of Rondônia, Northern Brazil, and report two new prey items consumed by this species.

On 22 October 2003 at 6 a.m., an adult male (forearm length, 111 mm) of *V. spectrum* was found electrocuted (resulting in a power outage) in the wiring of a lamppost of 10 m height at Santana ranch (line 9, lot 89, glebe 8) (11°24'64" S, 61°22'75" W), municipality of Cacoal,

Rondônia, Brazil. The specimen was collected and deposited at the mammalian collection of the Museu de História Natural da Universidade Estadual de Campinas (Campinas, SP), as a voucher (ZUEC 2457). The property where the animal was found is located in deforested areas open for agriculture and farming where fragments of the original Amazon Forest can be found scattered across the landscape. When found, this specimen had a smooth-billed ani (*Crotophaga ani*) (Cuculiformes, Cuculidae) in its mouth. In addition, we found remains of a partially digested house mouse (*Mus musculus*) (Rodentia, Muridae) inside its digestive tract. None of these prey items were reported to be consumed by this species to date. The smooth-billed ani is a relatively large bird (body weight about 120 g) typical from open areas (Sick 2001), and another species of ani (*Crotophaga sulcirostris*) had already been reported as a prey of *V. spectrum* in Costa Rica (Vehrencamp et al. 1977). The house mouse (20 g) is an invasive species, rare in the rainforest and generally restricted to the vicinity of buildings (Emmons and Feer 1997).

Absence of previous records of *Vampyrum spectrum* in Rondônia may be due to the scarcity of bat inventories in this state (Lewinsohn and Prado 2002) and the small abundance of this species in the nature (Emmons and Feer 1997). Considering the usual low densities of this bat species, local *V. spectrum* populations might be under threat due to site deforestation and frequent fires (Oliveira 2002).

NOTES ON GEOGRAPHIC DISTRIBUTION

Acknowledgements

To Dr. Ivan Sazima, Fátima M. de Souza and Elizabeth R. Bilo for providing the specimen registration number and to Drs. Vinícius Bonato and Wilson Uieda for their valuable suggestions. Thiago Carvalho performed the linguistic revision and Ariovaldo Giaretta critically read the draft.

Literature cited

- Acosta, L. and H. Azurduy. 2006. Primeras colectas del Falso Vampiro *Vampyrum spectrum* (Phyllostomidae, Chiroptera) en el sector sur del Bosque Seco Chiquitano, Santa Cruz, Bolivia. *Kempffiana* 2(1): 119-126.
- Bonato V., K. G. Facure, and W. Uieda. 2004. Food habits of bats of subfamily Vampyrinae in Brazil. *Journal of Mammalogy* 85(4): 708–713.
- Emmons, L. H. and F. Feer. 1997. Neotropical rainforest mammals: a field guide. 2nd ed. Chicago: University of Chicago Press. 307pp.
- Gregorin, R., A. P. Carmignotto, and A. R. Percequillo. 2008. Quirópteros do Parque Nacional da Serra das Confusões, Piauí, nordeste do Brasil. Chiroptera Neotropical 14(1): 366-383.
- Lewinsohn, T. M. and P. I. Prado. 2002. Biodiversidade brasileira: síntese do estado atual do conhecimento. São Paulo: Editora Contexto. 176 pp.
- Navarro, D. and D. E. Wilson. 1982. *Vampyrum spectrum*. *Mammalian Species* 185:1-4.
- Nogueira, M. R., A. L. Peracchi, and R. Moratelli. 2007. Subfamília Phyllostominae; p. 61-97 In N. R. dos Reis, A. L. Perachi, W. A. Pedro and I. P. Lima (ed.). Morcegos do Brasil. Londrina: Nélio Roberto dos Reis.
- Nowak, R. M. 1999. *Walker's Mammals of the World*. Baltimore: Johns Hopkins University Press. 836 pp.
- Oliveira, O. A. 2002. Geografia de Rondônia: Espaço e Produção. Porto Velho: Dinâmica Editora. 160 pp.
- Sick, H. 2001. *Ornitologia Brasileira*. 4th edn. Revista e ampliada por José Fernando Pacheco. Rio de Janeiro: Nova Fronteira. 862 pp.
- Vargas-Espinoza, A., L. F. Aguirre, M. S. Warner, L. Emmons, and M. Teran. 2004. Distribución de *Vampyrum spectrum* en Bolivia y comentarios sobre su estado de conservación. *Ecología en Bolivia* 39(2): 46-51.
- Vehrencamp, S. L., F. G. Stiles, and J. W. Bradbury. 1977. Observations on the foraging behavior and avian prey of the neotropical carnivorous bat, *Vampyrum spectrum*. *Journal of Mammalogy* 58(4): 469-477.

Received May 2009

Accepted July 2009

Published online August 2009