

Reptilia, Iguania, Liolaemidae, Liolaemus goetschi Müller and Hellmich, 1938: Distribution extension

Javier Nori 1,2*, Cristian S. Abdala 3 and Gustavo J. Scrocchi 3

- 1 Universidad Nacional de Córdoba, Centro de Zoología Aplicada, Facultad de Ciencias Exactas, Físicas y Naturales, Laboratorio de herpetología y animales venenosos. Rondeau 798. Córdoba (5000), Argentina.
- 2 Museo Patagónico de Ciencias Naturales. Av. Roca 1250, General Roca (8332). Río Negro, Argentina.
- 3 Fundación Miguel Lillo-CONICET y Facultad de Ciencias Naturales e IML, Instituto de Herpetología. Miguel Lillo 251. Tucumán (4000), Argentina.
- * Corresponding author. E-mail: javiernori@gmail.com

ABSTRACT: Liolaemus goetschi Müller and Hellmich, 1938 is a poorly known Patagonian lizard. Two specimens were collected in the central-western portion of Río Negro Province, Argentina, located 82 km west and 110 km south of the known distribution for the species in its type locality.

The genus Liolaemus is a particularly diverse group of iguanian lizards, with more than 200 species described to date, and other species waiting for description (Abdala and Lobo 2006; Abdala 2007; Pincheira-Donoso et al. 2007; Quinteros et al. 2008). Many of these species occur in a variety of habitats in Patagonia (Argentina) (Scolaro 2005; Abdala 2007; Morando et al. 2007). The knowledge of the geographic distribution of Liolaemus lizards is still very scarce. It is necessary to increase the information available on these lizards to improve our understanding of this large genus. Here we present new geographic distribution data on a poorly known Patagonian species, Liolaemus goetschi Müller and Hellmich 1938.

Whithin the boulengeri group, Liolaemus goetschi was described in 1938 by Müller and Hellmich on the basis of six specimens from Laguna Playa (province of Río Negro) stored at the Museum of Munich and Breslau. Unfortunately, this material is no longer available and the specimens could not be found by the curators of these collections. During a recent field trip to the central-western



FIGURE 1. Dorsal view of adult male of Liolaemus goetschi from the new locality, town of Cerro Policia, departament of El Cuy, province of Río Negro, in the Patagonian mount region of Argentina




FIGURE 2. Geographic distribution of *Liolaemus goetschi* in the province of Rio Negro. Star = type-locality; closed circle = new locality.

portion of Río Negro, we collected two specimens which were morphologically similar to the species described as Liolaemus goetschi (Figure 1). The individuals were collected by noose, euthanased with pericardic injection of tiopental sódico (Abbot®), fixed in 20 % formalin, and preserved in 70 % ethanol. Latitude, longitude, and elevation were determined with a Garmin[™] GPS 12.

The new site for Liolaemus goetschi is presented in Figure 2. It is situated 82 km west and 110 km south of the known distribution for the species (Laguna Playa, 20 km Northern of General Roca, Rio Negro, Argentina). Voucher specimens were deposited in the herpetological collection of the Museo Patagónico de Ciencias Naturales (MPCN-H: 0047-0048) housed in General Roca, Río Negro, Argentina.

Specimens were collected at the town of Cerro Policia, near the entrance to a private residence (39°43'40" S, 68°32'59" W, 640 m above sea level), department of El Cuy, province of Río Negro in the Patagonian mount region of Argentina, by J. Nori, M. J. Arellano, F. Fernadez, D. Fernandez, 5 January, 2008.

Vegetation is composed of shrubs (Larrea divaricata, L. cuneifolia, L. nitida, Cercidiurn praecox, Atriplex lampa), grasses (Stipa humilis, Bromus brevis brevis, Psila spartoides), and some trees (Salix humboldtiana, Maytenus boariacan). Lizards were located under vegetation, probably hunting insects.

To confirm the species identity we compared the specimens collected in the present study with those of *Liolaemus goetschi* from its type locality, deposited at the collections of the Museo Patagónico de Ciencias Naturales (MPCN) (MPCN 057-065) and Fundación Miguel Lillo (FML), (FML 21501-21505; 21504-21507; 21511) as well as with the original description (Müller and Hellmich 1938).

LITERATURE CITED

- Abdala, C.S. 2007. Phylogeny of the boulengeri group (Iguania: Liolaemidae, Liolaemus) based on morphological and molecular characters. Zootaxa 1538: 1-84.
- Abdala, C.S. and F. Lobo. 2006 Description of a new Patagonian lizard species of the Liolaemus silvanae group (Iguania: Liolaemidae). South American Journal of Herpetology 1: 1-8.

- Morando M., L.J. Avila, C.R. Turner and J.W. Sites Jr. 2007. Molecular evidence for a species complex in the patagonian lizard Liolaemus bibronii and phylogeography of the closely related Liolaemus gracilis (Squamata: Liolaemini). Molecular Phylogenetics and Evolution 43: 952-973
- Müller, L. and W. Hellmich. 1938. Liolaemus -Arten aus dem Westlichen Argentinien (Ergebnisse der Argentinienreise von Prof. Dr. W. Goetsch, Breslau). I. Liolaemus darwini (Bell) und Liolaemus goetschi n. sp. Zoologischer Anzeiger 123 (5-6): 130-142.
- Pincheira-Donoso, D.J., A. Scolaro and A. Schulte. 2007. The limits of polymorphism in *Liolaemus rothi*: Molecular and phenotypic evidence for a new species of the Liolaemus boulengeri clade (Iguanidae, Liolaemini) from boreal Patagonia of Chile. Zootaxa 1452: 25-42.
- Quinteros, A.S., C.S. Abdala, J.M. Diaz Gomez and G.J. Scrocchi. 2008. Two new species of Liolaemus (Iguania: Liolaemidae) of central west Argentina. South American Journal of Herpetology 3:101-111.
- Scolaro, J.A. 2005. Reptiles Patagónicos: Sur. Guía de campo. Trelew: Ediciones Universidad Nacional de la Patagonia. 80 p.

RECEIVED: August 2009 REVISED: November 2009 Accepted: December 2009 Published online: February 2010

EDITORIAL RESPONSIBILITY: Julio Cesar de Moura-Leite