

Amphibia, Anura, Hemiphractidae, Hemiphractus helioi Sheil and Mendelson, 2001: Distribution extension in the state of Acre and second record for Brazil

Paulo Sérgio Bernarde 1*, Daniele Bazzo Miranda 2, Saymon de Albuquerque 2 and Luiz Carlos Batista Turci²

- 1 Universidade Federal do Acre UFAC, Campus Floresta, Centro Multidisciplinar, Laboratório de Herpetologia. CEP 69980-000. Cruzeiro do Sul,
- 2 Universidade Federal do Acre, Programa de Pós Graduação em Ecologia e Manejo de Recursos Naturais. CEP 69915-900. Rio Branco, AC, Brazil
- * Corresponding author. E-mail: SnakeBernarde@hotmail.com

ABSTRACT: We report the second records of Hemiphractus helioi from Brazil based on specimens collected during field studies in forest of Môa River, Cruzeiro do Sul, Acre. These new records extended the range of this species about 100 km to the east from the nearest collection site known in Parque Nacional da Serra do Divisor.

The Alto Juruá region is located at the northwestern limits of Brazil and it is known for exhibiting high amphibian species richness; 125 species of frogs occur only in Parque Nacional da Serra do Divisor (Souza 2009). Despite the region's high diversity, the knowledge on amphibians from Alto Juruá is based only on a few studies (e.g. Martins and Cardoso 1987; Cardoso and Vielliard 1990; Gascon 1996; Bernarde et al. 1999; Caldwell 2005; Souza 2009).

The horned treefrog Hemiphractus helioi was described by Sheil and Mendelson III (2001), and has been wrongly identified for almost a century (Noble, 1917) as H. johnsoni, an endangered species according to the IUCN Red List of Threatened Species (Silvano and Segalla 2005).

The current geographic range of *H. helioi* comprises the upper Amazon Basin and the lower Amazon slopes of the Andes, from Ecuador to northern Bolivia (Sheil and Mendelson 2001) and Brazil (as H. johnsoni) (Silvano and Segalla 2005). Köhler (in litt. apud Frost 2009) suggested that the record by Sheil and Mendelson III (2001) in



FIGURE 1. Specimen of Hemiphractus helioi found on the vegetation (1 m above ground) at night on 13 November 2008 in Môa River forest, municipality of Cruzeiro do Sul, Acre, Brazil.



FIGURE 2. Specimen of Hemiphractus helioi found on the vegetation (40 cm above ground) at night on 15 December 2008 in Môa River forest, municipality of Cruzeiro do Sul, Acre, Brazil.

northern Bolivia was wrong. Recently, this species was reported in Brazil (as H. helioi), in Parque Nacional da Serra do Divisor (Souza 2009), near Peru. Based on specimens collected during field studies in the lowland forest of Môa River in Acre, we extended the range of this species and made its second record in Brazil.

On November 13th 2008 and on December 15th 2008 two individuals of *H. helioi* were collected in the lowland forest of Môa River (7°38'23.9" S, 72°47'43.1" W), municipality of Cruzeiro do Sul, state of Acre, northern Brazil (collecting permits granted by Ibama (SISBIO) #12178-1 and #12178-2). They were found in a primary forest at night, perching on the vegetation at 40 and 100 cm above the ground (Figures 1 and 2). The specimens were deposited in the herpetological collection of Universidade Federal do Acre -Campus Floresta (UFACF 1191, 1257).

These new records from the lowland forest of Môa River in Cruzeiro do Sul extended the range of this species about 100 km to the east (Figure 3) from the nearest collection site known in Parque Nacional da Serra do Divisor (Souza 2009).




FIGURE 3. Map of Brazil showing the state of Acre and the known distribution of Hemiphractus helioi. Red circles indicate the three localities of the records: 1 = Rio Môa, casa do Piroca, P. N. Serra do Divisor; 2 = Trilha do Anil, P. N. Serra do Divisor; 3 = Igarapé Ramon, P. N. Serra do Divisor; 4 = Môa River lowland forest, Cruzeiro do Sul.

ACKNOWLEDGMENTS: We thank Moisés Barbosa de Souza for providing us with information on *Hemiphractus helioi* specimens of the herpetological collection of Universidade Federal do Acre (UFAC) - Campus Rio Branco. Field work was funded by Conselho Nacional de Desenvolvimento Científico (CNPq) (Project: 479225/2007-9). CAPES granted fellowships to D. B. Miranda and S. Albuquerque, and CNPq granted a fellowship to P. S. Bernarde (501927/2009-3). We also thank Reginaldo Machado for reviewing our manuscript.

LITERATURE CITED

Bernarde, P.S., M.B. Souza and M.C.N. Kokubum. 1999. Predation on Hyla minuta Peters, 1872 (Anura, Hylidae) by Ancylometes spp. (Araneae, Pisauridae). Biociências 7: 199-203.

Caldwell J.P. 2005. A new Amazonian species of Cryptophyllobates (Anura: Dendrobatidae). Herpetologica 61(4): 449-461.

Cardoso, A.J. and J. Vielliard. 1990. Vocalizações de anfíbios anuros de um ambiente aberto, em Cruzeiro do Sul, Estado do Acre. Revista Brasileira de Biologia 50: 229-242.

Frost D.R. 2009. Amphibian Species of the World: an Online Reference. Version 5.3 Electronic Database accessible at http://research.amnh. org/herpetology/amphibia/. American Museum of Natural History, New York, USA. Captured on 12 February 2009.

Gascon C. 1996. Amphibian litter fauna and river barriers in flooded and non-flooded Amazonian Rain Forest. Biotropica 28: 136-140.

Martins, M. and A.J. Cardoso. 1987. Novas espécies de hilídeos do Estado do Acre (Amphibia: Anura). Revista Brasileira de Biologia 47: 549-

Sheil, C.A. and J.R. Mendelson III. 2001. A new species of Hemiphractus (Anura: Hylidae: Hemiphractinae), and a redescription of *H. johnsoni*. Herpetologica 57(2): 189-202.

Silvano, D.L. and M.V. Segalla. 2005. Conservation of Brazilian amphibians. Conservation Biology 19(3): 653-658.

Souza, M.B. 2009. Anfíbios - Reserva Extrativista do Alto Juruá e Parque Nacional da Serra do Divisor, Acre. Série Pesquisa e Monitoramento Participativo em Áreas de Conservação Gerenciadas por Populações Tradicionais. Volume 2. Campinas: Instituto de Filosofia e Ciências Humanas (IFCH) Unicamp. 77 p.

RECEIVED: November 2009 REVISED: August 2010 ACCEPTED: August 2010

Published online: October 2010

Editorial responsibility: Marcelo N. de C. Kokubum