

Reptilia, Squamata, Gymnophthalmidae, *Arthrosaura reticulata* (O'Shaughnessy, 1881): Distribution extension and new state record

Tamí Mott^{1*}, Luana Aparecida Gomes de Arruda², Anelise Figueiredo Silva², Jacqueline Pimentel Silva², Ricardo Alexandre Kawashita-Ribeiro² and Marcos André de Carvalho¹

- 1 Universidade Federal de Mato Grosso, Instituto de Biociências, Departamento de Biologia e Zoologia, Avenida Fernando Corrêa da Costa 2367, CEP 78060-900 Cuiabá, MT, Brasil.
 - 2 Universidade Federal de Mato Grosso, Instituto de Biociências, Coleção Zoológica de Vertebrados, Avenida Fernando Corrêa da Costa 2367, CEP 78060-900 Cuiabá, MT, Brasil.
- * Corresponding author: tamimott@hotmail.com

ABSTRACT: *Arthrosaura reticulata* (O'Shaughnessy, 1881) is reported from two localities in Mato Grosso, Brazil. It represents the first state record for the species, extending its known distribution in 830 km southeastern of the southernmost record in the state of Amazonas and 470 km eastern of the state of Rondônia, the southernmost record of the species.

Arthrosaura reticulata (O'Shaughnessy, 1881) is a small gymnophthalmid lizard of subfamily Ecleopinae (Castoe *et al.* 2004), with three supraoculars, 6-7 supralabials, no vertebral stripe, cylindrical body, tail round in cross section, with relatively short limbs, and elongate, keeled dorsal scales (Hoogmoed and Avila-Pires 1992; Avila-Pires 1995). The countries with records for this species are Ecuador (the type-locality of the species, Canelos, Pastaza), Peru, Colombia, Guyana, and Suriname (review in Hoogmoed and Avila-Pires 1992). In Brazil, the species has been reported from the states of Maranhão, Pará, Amapá, Amazonas, and Rondônia (Vanzolini 1986; Hoogmoed and Avila-Pires 1992; Avila-Pires 1995; Avila-Pires *et al.* 2009).

Here we report a new Brazilian state record for *A. reticulata*, in the state of Mato Grosso, extending its geographic distribution in 830 km southeastern of the southernmost record in the state of Amazonas, and 470 km eastern of the state of Rondônia, the southernmost record of the species (Figure 1). The two new records are located in the Amazonian rainforest. Voucher specimens are deposited in the Coleção Zoológica de Vertebrados do Instituto de Biociências da Universidade Federal de Mato Grosso (UFMT).

On June 1, 2008, one juvenile of *Arthrosaura reticulata* (UFMT 7124) was collected in the Parque Estadual Igarapés do Juruena, Colniza municipality (08°54' S, 59°06' W), at 12:20 h, active on the leaf litter near a creek. The specimen has a snout-vent length of 27.2 mm and a mutilated tail with 13.5 mm of remaining length. On December 9, 2009, a male (UFMT 8371, Figure 2) was collected during active search in Fazenda São Nicolau, Cotriguaçu municipality (09°48' S, 58°16' W), at 09:30 h, active on the leaf litter near the Juruena river. The specimen has a snout-vent length of 41.8 mm and a tail length of 80.9 mm.




FIGURE 1. New state record for *Arthrosaura reticulata* in Mato Grosso, Brazil. Triangle: Southernmost record of the species in the state of Amazonas (Fazenda Scheffer, Ituxi river, Avila-Pires *et al.* 2009); Square: Southernmost record, the state of Rondônia (Santa Cruz da Serra, Vanzolini 1986). Stars: Colniza and Cotriguaçu municipalities, state of Mato Grosso (new records), extending the distribution of the species on 725 km (Colniza) and 830 km (Cotriguaçu) southeastern of Fazenda Scheffer, state of Amazonas, 425 km (Colniza) and 470 km (Cotriguaçu) southeastern of Santa Cruz da Serra, state of Rondônia. Inset map represents South America where Amazonas and Rondônia states are shaded and Mato Grosso state is solid color.



FIGURE 2. Adult *Arthrosaura reticulata* (UFMT 8371) collected in Cotriguaçu municipality, state of Mato Grosso, Brazil. Photo by R. A. Kawashita-Ribeiro.

The number of notes about geographical extensions or even new state records of Squamata reptiles in the state of Mato Grosso has increased considerably during the last years (*e.g.* Costa *et al.* 2008; Kawashita-Ribeiro and Ávila 2008; Mott *et al.* 2008; Costa *et al.* 2010; Morais *et al.* 2010; Silva *et al.* 2010). It reinforces the need of biological surveys in many unexplored regions of this state. This action is urgent because natural landscapes are being converted into agricultural and pasture lands at an accelerated and alarming rate and they may disappear even before we have a good knowledge of the biodiversity in the state.

ACKNOWLEDGMENTS: Tami Mott, Ricardo A. Kawashita-Ribeiro and Marcos A. de Carvalho thanks NIEFA (Núcleo Interdisciplinar de Estudos Faunísticos, FAPEMAT 447441/2009).

LITERATURE CITED

- Avila-Pires, T.C.S. 1995. Lizards of Brazilian Amazonia (Reptilia: Squamata). *Zoologische Verhandelingen, Leiden*, 299: 1-706.
- Avila-Pires, T.C.S., L.J. Vitt, S.S. Sartorius and P.A. Zani. 2009. Squamata (Reptilia) from four sites in southern Amazonia, with a biogeographic analysis of Amazonian lizards. *Boletim do Museu Paraense Emílio Goeldi, Ciências Naturais* 4(2): 99-118.
- Castoe, T.A., T.M. Doan and C.L. Parkinson. 2004. Data partitions and complex models in Bayesian analysis: the phylogeny of Gymnophthalmid lizards. *Systematics Biology* 53(3): 448-469.
- Costa, H.C., V.A. São-Pedro, A.K. Peres-Jr. and R.N. Feio. 2008. Reptilia, Squamata, Teiidae, *Tupinambis longilineus*: Distribution extension. *Check List* 4(3): 267-268.
- Costa, H.C., V.A. São-Pedro and R.N. Feio. 2010. A new record of the poorly known Amazonian snake *Siphlophis worontzowi*. *Herpetology Notes* 3: 97-100.
- Hoogmoed, M.S. and T.C.S. Avila-Pires. 1992. Studies on the species of the South American lizard genus *Arthrosaura* Boulenger (Reptilia: Sauria: Teiidae), with the resurrection of two species. *Zoologische Mededelingen, Leiden* 66(35): 453-484.
- Kawashita-Ribeiro, R.A. and R. W. Ávila. 2008. Reptilia, Squamata, *Polychrus* spp.: New record, range extensions, and distribution map in the state of Mato Grosso, Brazil. *Check List* 4(3): 362-365.
- Morais, D.H., T. Mott, R.A. Kawashita-Ribeiro and A.P. Santos-Jr. 2010. Reptilia, Squamata, Dipsadidae, Xenodontinae, *Taeniophallus brevirostris* (Peters, 1863): Distribution extension and new state record. *Check List* 6(3): 456-457.
- Mott, T., D.H. Morais and R. A. Kawashita-Ribeiro. 2008. Reptilia, Squamata, Amphisbaenidae, *Anops bilabialatus*: Distribution extension, meristic data, and conservation. *Check List* 4(2): 146-150.
- Silva, P.C., T. Mott and R.A. Kawashita-Ribeiro. 2010. Reptilia, Squamata, Amphisbaenia, *Amphisbaena cuiabana* (Strussmann and Carvalho, 2001): Range extension in the state of Mato Grosso, Brazil. *Check List* 6(4): 644-645.
- Vanzolini, P.E. 1986. *Levantamento herpetológico da área do Estado de Rondônia sob a influência da rodovia Br-364*. Polonoreste/Ecologia Animal. Brasília: Relatório de Pesquisa nº1, CNPq. 50 p.

RECEIVED: March 2010

LAST REVISED: August 2010

ACCEPTED: December 2010

PUBLISHED ONLINE: January 2011

EDITORIAL RESPONSIBILITY: Daniel Loebmann