

Mammalia, Chiroptera, Phyllostomidae, *Platyrrhinus recifinus* (O. Thomas, 1901): First confirmed record in the state of Santa Catarina, southern Brazil

Fernando Carvalho* e Marta Elena Fabián

Programa de Pós-Graduação em Biologia Animal, Universidade Federal do Rio Grande do Sul, Departamento de Zoologia, Instituto de Biociências Avenida Bento Gonçalves. CEP 91501-970. Porto Alegre, RS, Brasil.

* Corresponding author. E-mail: fernando_bats@yahoo.com.br

ABSTRACT: We report the first confirmed record of *Platyrrhinus recifinus* to Santa Catarina state, south of Brazil. The distribution of this species was expanded approximately 350 km south of its previous known limit.

The genus *Platyrrhinus* Saussure, 1860 occurs from southern Mexico to northern Argentina (Gardner 2008). The diagnostic characters for the genus are: presence of two accessory cusps on the posterior face of P4, presence of three upper molars, and presence of a fringe of fur at the edge of uropatagium (Velazco 2005).

Nowadays *Platyrrhinus* is formed by at least 18 species (Velazco *et al.* 2010), but this complex of species presents difficulties in their identification (Velazco 2005). According to Zortéa (2007) and Velazco *et al.* (2010), seven species occur in Brazil: *P. brachycephalus* (Rouk and Carter, 1972), *P. fusciventris* Velazco, Gardner and Patterson, 2010, *P. helleri* (Peters, 1866), *P. incarum* (Thomas, 1912), *P. infuscus* (Peters, 1880), *P. lineatus* (É. Geoffroy, 1810), and *P. recifinus* (O. Thomas, 1901).

Platyrrhinus recifinus has confirmed records in northeast (Paraíba, Bahia, Espírito Santo, Ceará and Pernambuco states; Peracchi and Albuquerque 1993; Sousa *et al.* 2004; Faria *et al.* 2006; Zortéa 2007; Piccinini 1974 apud Gardner 2008) and southeast Brazilian states (Minas Gerais, Rio de Janeiro and São Paulo; Pedro *et al.* 2001; Esbérard and Bergallo 2005; Tavares *et al.* 2007), and recently the species was recorded in the state of Paraná, southern Brazil (Scultori *et al.* 2009). *Platyrrhinus recifinus* was considered endemic of Brazil, but Tavares and Velazco (2010) reported its occurrence also for Guyana and Suriname.

The biology of *P. recifinus* is little known (Gardner 2008). In part the lack of information may be associated to the low capture rates (*e.g.* Dias *et al.* 2002; Passos *et al.* 2003; Faria 2006; Tavares *et al.* 2007; Nobre *et al.* 2009). In Brazil this species has been sampled in several environments, such as preserved forests and impacted areas (*e.g.* Faria 2006; Gardner 2008; Tavares *et al.* 2007; Dias and Peracchi 2008). Like the other species of the genus, *P. recifinus* probably has a frugivorous diet (Zortéa 2007). The species is included in the List of Brazilian Fauna Threatened of Extinction, in the "Vulnerable" category,

due to habitat destruction, deforestation and fires in the forest (Machado *et al.* 2008). Globally *P. recifinus* is listed in the category "least concern" because it occurs in areas with different levels of disturbance Sampaio *et al.* (2008). We report the first confirmed record of *P. recifinus* in Santa Catarina state, expanding the southern limit of its distribution.

The record reported here was obtained during a study of the vertical structure of the bat community in a forest remnant in southern Brazil, between July 2005 and September 2010. Bats were captured with mist nets installed in different strata (understory, sub-canopy and canopy) opened for six hours after the sunset. A total of 74 nights were sampled, 39 in the understory and 35 in sub-canopy and canopy. The total sampling effort, calculated according to Straube and Bianconi (2002), was 26.910 m².h in the understory and 26.460 m².h in the sub-canopy and canopy. The permit to carry out the sampling was provided by the Instituto Chico Mendes de Conservação da Biodiversidade – ICMBio under number 22648-1 SISBIO.

The area where the record occurred is situated in Rancho dos Bugres, locality of the Pedras Grandes city (28°29'40" S, 49°15'24" W) in the southern state of Santa Catarina (Figure 1). The area is a Tropical Rain Forest remnant (Veloso *et al.* 1992) of about 7 ha with different succession stages, connected to larger forest remnants. According to the classification of Köppen the climate is Cfa (temperate subtropical), with average annual temperatures ranging between 22 and 24 °C in the warmer months and 13 and 15 °C in winter ones. The mean of annual rainfall is between 1250 and 1500 mm (Nimer 1990).

On April 17th, 2010, at 23:16 h one adult male *P. recifinus* was captured in a mist net installed at 7.5 m above the ground (canopy). The specimen was collected and incorporated into the scientific collection of the Departamento de Zoologia, Instituto de Biociências da Universidade Federal do Rio Grande do Sul (DZMAM 1230). Measurements of this specimen (Table 1) were obtained

using a caliper to the nearest 0.01 mm and following the protocol described by Taddei *et al.* (1998) and Scultori *et al.* (2009). Species identification was based on Velazco (2005), Gardner (2008), Dias and Peracchi (2008), Tavares and Velazco (2010) and Velazco *et al.* (2010). The diagnostic characters observed were: interramal vibrissae absent; forearm length less than 45 mm; lists facial and dorsal evident; a deep fossa in the hypoconal basin of the second upper premolar (P4) and a stylid cusp absent between the metaconid and the protoconid of first lower molar (m2); upper incisors completely separated and lower incisors with few distinct lobes and margins nearly straight.

Studies on the bat fauna of Santa Catarina include 46 species (Passos *et al.* 2010), with no reference to *P. recifinus*. Zortéa (2007) and Tavares and Velazco (2010) have reported the occurrence of this species in Santa Catarina, but they did not specify the site nor the scientific collection where specimens were deposited. Our study presents the first confirmed record of *P. recifinus* for Santa Catarina, extending in 350 km the southern limit of its distribution as described by Scultori *et al.* (2009).

Tavares and Velazco (2010) suggest that *P. recifinus* presents clinal variation, in which the specimens from southern Brazil would be morphologically larger. This variation was verified by Scultori *et al.* (2009) comparing specimens captured in Paraná state, with others from the southeast and north of Brazil. Nevertheless, for the specimen sampled by us, only the measurements of body length, length of the series of jaw teeth, and




FIGURE 1. Locality records of *Platyrrhinus recifinus* in the states of Paraná (Scultori *et al.* 2009) (●) and Santa Catarina (present study) (▲), in southern Brazil.

width between the upper canines were above the range described for this species in other regions (Table 1). According to Scultori *et al.* (2009) it is necessary to study a larger number of individuals in order to obtain more reliable data about the possible morphological variations occurring along the latitudinal gradient of the species distribution.

TABLE 1. Measurements of *P. recifinus* in southern Santa Catarina and from studies conducted in southeastern and southern Brazil. Min = minimum and Max = maximum. Total body length (TL); hind foot length (HL); length of forearm (FA); greatest length of skull, including incisors (GLS); condyloincisive length (CIL); basal length, including incisors (BL); palatal length, including incisors (PL); length of maxillary toothrow (MAXT); breadth across upper canines (BAC); breadth across upper molars (BAM); postorbital breadth (PB); zygomatic breadth (ZB); breadth of braincase (BB); mastoid breadth (MB); mandibular length (ML); length of mandibular toothrow (MANT).

Character	Dias <i>et al.</i> (2002) Min - Max (n = 2)	Dias and Peracchi (2008) Min - Max (n = 9)	Scultori <i>et al.</i> (2009) Min - Max (n = 5)	This study (n = 1)
TL	50.20 - 58.30	54.14 - 59.12	57.92 - 64.38	61.15
HL	9.20 - 10.50	10.50 - 13.54	11.84 - 13.44	12.66
FA	41.70 - 42.00	41.66 - 43.54	42.64 - 45.96	42.60
GLS	23.90 - 24.20	23.50 - 25.24	24.56 - 25.50	24.50
CIL	21.60 - 22.00	22.04 - 22.98	22.36 - 23.42	22.31
BL	19.40 - 19.80	19.84 - 20.66	19.82 - 20.84	20.17
PL	11.30 - 11.60	11.74 - 12.40	11.88 - 12.76	11.97
MAXT	8.70 - 8.80	9.00 - 9.56	9.20 - 9.72	9.77
BAM	10.20 - 10.40	10.28 - 10.98	10.74 - 11.44	11.22
BAC	6.00 - 6.10	6.00 - 6.44	6.28 - 6.70	6.72
PB	5.50 - 5.70	5.54 - 6.24	5.84 - 6.02	5.63
ZB	13.80 - 14.10	13.74 - 14.64	14.58 - 14.98	14.36
BB	10.30 - 10.50	10.20 - 10.74	10.50 - 11.22	10.39
MB	11.70 - 12.30	11.76 - 12.12	12.08 - 12.60	12.14
ML	15.60 - 16.00	16.20 - 16.96	16.18 - 17.20	15.80
MANT	9.50 - 9.50	9.54 - 10.26	9.62 - 10.48	9.67

ACKNOWLEDGMENTS: To Daniel S. de Souza for assistance in the measurement and preparation of material; to Karolina S. Souza, Poliana B. Peres, Rodrigo Á. Mendonça, André Tschiedel, Flávia V. Fonceca and Rildo V. Gonçalves, for their assistance during fieldwork; and to CAPES for Msc. Scholarship concessioned to F. Carvalho.

LITERATURE CITED

- Dias, D. and A.L. Peracchi. 2008. Quirópteros da Reserva Biológica do Tinguá, estado do Rio de Janeiro, sudeste do Brasil (Mammalia: Chiroptera). *Revista Brasileira de Zoologia* 25(3): 333-369.
- Dias, D., A.L. Peracchi and S.S.P. Silva. 2002. Quirópteros do Parque Estadual da Pedra Branca, Rio de Janeiro, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 19(1-2): 113-140.
- Esbérard, C.E.L. and H.G. Bergallo. 2005. Research on bats in the state of Rio de Janeiro, Southeastern Brazil. *Mastozoología Neotropical* 12(2): 237-243.
- Faria, D. 2006. Phyllostomid bats of a fragmented landscape in the northeastern Atlantic forest, Brazil. *Journal of Tropical Ecology* 22: 531-542.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from Atlantic Forest of Southern Bahia, Brazil. *Biota Neotropica* 6(2): 1-13.
- Gardner, A.L. 2008. Genus *Platyrrhinus* Saussure, 1860; p. 329-342 In A.L. Gardner (ed.). *Mammals of South America. Volumen 1: Marsupials, xenarthrans, shrews and bats*. Chicago: The University of Chicago (dated 2007, but published in 2008).
- Machado, A.B.M., G.M. Drummond and A.P. Paglia. 2008. *Livro vermelho da Fauna Brasileira Ameaçada de Extinção*. Brasília / Belo Horizonte: Ministério do Meio Ambiente / Fundação Biodiversitas. 998 p.
- Nimer, E. 1990. Clima; p. 151-187 In Instituto Brasileiro de Geografia e Estatística (ed.). *Geografia do Brasil: região Sul*. Rio de Janeiro, Instituto Brasileiro de Geografia e Estatística.
- Nobre, P.H., A.S. Rodrigues, A.I. Costa, A.E.S. Moreira and H.H. Moreira. 2009. Similaridade da fauna de Chiroptera (Mammalia), da Serra Negra, municípios de Rio Preto e Santa Bárbara do Monte Verde, Minas Gerais, com outras localidades da Mata Atlântica. *Biota Neotropica* 9(3): 151-156.
- Passos, F.C., R.W. Silva, W.A. Pedro and R.M. Bonin. 2003. Frugivoria em morcegos (Mammalia, Chiroptera) no Parque Estadual Intervales, sudeste do Brasil. *Revista Brasileira de Zoologia* 20(3): 511-517.
- Passos, F.C., J.M.D. Miranda, I.P. Bernardi, N.Y. Kaku-Oliveira and L.C. Munster. 2010. Morcegos da Região Sul do Brasil: análise comparativa da riqueza de espécies, novos registros e atualizações nomenclaturais (Mammalia, Chiroptera). *Iheringia* 100(1): 25-34.
- Pedro, W.A., F.C. Passos and B.K. Lim. 2001. Morcegos (Mammalia, Chiroptera) da Estação Ecológica de Caetetus, Estado de São Paulo. *Chiroptera Neotropical* 7(1-2): 136-140.
- Peracchi, A.L. and S.T. Albuquerque. 1993. Quirópteros do município de Linhares, Estado do Espírito Santo, Brasil (Mammalia, Chiroptera). *Revista Brasileira de Biologia* 53 (4): 575-581.
- Sampaio, E., B. Lim and S. Peters. 2008. *Platyrrhinus recifinus*. In IUCN. *IUCN Red List of Threatened Species. Version 2010.4*. Electronic Database Accessible at www.iucnredlist.org. Captured on 21 February 2011.
- Scultrori, C., D. Dias and A.L. Peracchi. 2009. Mammalia, Chiroptera, Phyllostomidae, *Platyrrhinus recifinus*: first record in the state of Paraná, southern Brazil. *Check List* 5(2): 238-242.
- Sousa, M.A.N., A. Langguth and E.A. Gimenez. 2004. Mamíferos dos brejos de altitude Paraíba e Pernambuco. p.229-249. In K.C. Porto, J.J.P. Cabral and M. Tabarelli (ed.). *Brejos de altitude em Pernambuco e Paraíba, história natural, ecologia e conservação*. Série Biodiversidade 9. Brasília: Ministério do Meio Ambiente.
- Straube, F.C. and G.V. Bianconi. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com a utilização de redes de neblina. *Chiroptera Neotropical* 8(1-2): 150-152.
- Taddei, V.A., C.A. Nobile and E. Moreille-Versute. 1998. Distribuição geográfica e análise morfométrica comparativa em *Artibeus obscurus* (Schinz, 1821) e *Artibeus fimbriatus* Gray, 1838 (Mammalia, Chiroptera, Phyllostomidae). *Ensaios e Ciência* 2(2): 71-127.
- Tavares, V.C. and P.M. Velazco. 2010. *Platyrrhinus recifinus* (Chiroptera: Phyllostomidae). *Mammalian Species* 42(859):119-123.
- Tavares, V.C., F.A. Perini and J. Lombardi. 2007. The bat communities (Chiroptera) of the Parque Estadual do Rio Doce, a large remnant of Atlantic Forest in southeastern Brazil. *Lundiana* 8(1): 35-47.
- Velazco, P.M. 2005. Morphological Phylogeny of the Bat Genus *Platyrrhinus* Saussure, 1860 (Chiroptera: Phyllostomidae) with the Description of Four New Species. *Fieldiana* 105: 1-54.
- Velazco, P.M., A.L. Gardner and B.D. Patterson. 2010. Systematic of the *Platyrrhinus helleri* species complex (Chiroptera, Phyllostomidae), with descriptions of two new species. *Zoological Journal of the Linnean Society* 159: 785-812.
- Veloso, H.P., L.C. Oliveira-Filho, A.M.S.F. Vaz, M.P.M. Lima, R. Marquette and J.E.M. Brazao. 1992. *Manual técnico da vegetação brasileira*. Manuais técnicos em geociências. Rio de Janeiro: Secretaria de Planejamento, Fundação Instituto Brasileiro de Geografia e Estatística, Diretoria de Geociências, Departamento de Recursos Naturais e Estudos Ambientais. 92 p.
- Zortéa, M. 2007. Subfamília Stenodermatinae; p.117-128 In N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.). *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina.

RECEIVED: October 2010

LAST REVISED: March 2011

ACCEPTED: March 2011

PUBLISHED ONLINE: March 2011

EDITORIAL RESPONSIBILITY: Marcelo R. Nogueira