

Distribution extension and distribution map of *Chiasmocleis jimi* Caramaschi and Cruz, 2001 (Amphibia: Anura: Microhylidae) and *Proceratophrys concavitympanum* Giaretta, Bernarde and Kokubum, 2000 (Amphibia: Anura: Cycloramphidae)

Pedro H. Bernardo^{1,2*}, William Matiazzi² and Ricardo A. Guerra-Fuentes²

1 Royal Ontario Museum, Centre for Biodiversity and Conservation Biology, 100 Queen's Park, Toronto, ON, M5S 2C6, Canada.

2 Museu de Zoologia da Universidade de São Paulo, Avenida Nazaré, 481, CEP 04263-000, São Paulo, SP, Brazil.

* Corresponding author. E-mail: bernardoph@gmail.com

ABSTRACT: We provide a review of the geographic range of *Chiasmocleis jimi* and *Proceratophrys concavitympanum* and include new records for these rare species in the Reserva Biológica do Tapirapé, southern Pará, Brazil. Our records extend the known distribution of *C. jimi* and *P. concavitympanum* in Pará state.

Chiasmocleis jimi Caramaschi and Cruz, 2001 is a small microhylid that occurs in the Amazon Rainforest with just a few published records. It was originally described from two localities in Amazonas and Pará states, Brazil (Caramaschi and Cruz 2001). Peloso and Sturaro (2008) collected *C. jimi* from two other localities in Pará. This species is also recorded in Pará state at the Floresta Nacional de Caxiuanã (Ribeiro-Júnior *et al.* 2011) and Reserva Biológica do Rio Trombetas (Funk and Cannatella 2009). Regardless of these records, Frost (2011) and Caramaschi (2004) listed *C. jimi* as known only from the type locality.

Proceratophrys concavitympanum Giaretta, Bernarde, and Kokubum, 2000 is a species without eyelid appendages from the *P. cristiceps* group (*sensu* Prado and Pombal-Jr 2008). It is the only species of *Proceratophrys* known in the Amazon Rainforest (Giaretta *et al.* 2000), originally described from specimens collected in Espigão D'Oeste in Rondônia state. The first record outside the type locality was made by Amaro *et al.* (2009) in UHE Lajeado in Tocantins and the first record for Pará state was made by Barros *et al.* (2011). To this date, some other records for this species were made in other Amazonian areas (Pavan and Dixo 2004; Santana *et al.* 2010; Ávila *et al.* 2011).

Both species were collected at the Reserva Biológica do Tapirapé (REBIOTA) (05°30'13" S, 50°16'44" W) a 150,000 ha reserve located in the municipalities of Marabá and Parauapebas, Pará, Brazil. They were collected during the dry season, from August 3rd to 12th, 2008 and during the rainy season from November 29th to December 9th, 2008. In total, 12 specimens of *C. jimi* and eight specimens of *P. concavitympanum* were collected in pitfall traps, while no specimens were found during active search.

The specimens were collected under permit SISBIO number 16826-1 issued by the Brazilian Ministry of

Environment (MMA- ICMBio) and deposited in the herpetological collection at the Museu de Zoologia da Universidade de São Paulo under the numbers: *C. jimi*: MZUSP 139635-139642, 140085, 140139-140141; *P. concavitympanum*: MZUSP 139426-139430, 140118, 140119.

REBIOTA is located in the Mosaico de Carajás, an area that includes five Federal Conservation Units (REBIOTA, Floresta Nacional do Tapirapé-Aquiri, Floresta Nacional de Carajás, Floresta Nacional do Itacaiunas and Área de Proteção Ambiental do Igarapé Gelado) totaling over 800,000 ha of protected areas in southeastern Pará. The predominant vegetation in the region is Floresta Ombrófila Aberta and Floresta Ombrófila Densa (IBGE, 1992) with various palms, bamboos and canga. According to the National Protected Areas Plan, REBIOTA is classified as a protected area of extremely high biological importance (MMA 2007) due to the great human pressure that surrounds the reserve.

The specimens of *C. jimi* (Figure 1A-C) conform to the original description and the paratypes (MZUSP 53980-53988). *Chiasmocleis jimi* can be separated from the other *Chiasmocleis* in the Amazon by the characteristic coloration and by exhibiting swollen tips of fingers III and IV and of toes II-V (tips of all digits not expanded in the other species) (Caramaschi and Cruz 2001). The specimens from REBIOTA extend the range 283 km to the southeast (Figure 2).

The specimens of *P. concavitympanum* collected at REBIOTA (Figure 1D-F) conform to the original description (Giaretta *et al.* 2000), in particular the following characteristics: eyelids bordered by large, irregularly arranged, warts; postocular swellings absent; snout obtuse with flaring lip in profile; tympanum defined as a

depression in the skin (visible in Figure 1E) and adult SVL: mean = 48.92 mm; N= 3. The specimens from REBIOTA extend the range 510 km, constituting the northernmost record for this species and the first record in Pará state (Figure 2).




FIGURE 1. A, B, C: *Chiasmocleis jimi* MZUSP 140140 from REBIOTA; D, E, F: *Proceratophrys concavitympanum* from REBIOTA (D and F: MZUSP 140119; E: MZUSP 139428).

According to the IUCN, *C. jimi* and *P. concavitympanum* are listed as “Data Deficient” due to the lack of information regarding natural history and geographic distribution (Caramaschi 2004; Rodrigues and Caramaschi 2004). This reinforces the importance of these new records and the importance of field surveys in unexplored areas, especially in the very threatened tropical forests of Brazil.




FIGURE 2. Geographic distribution of *Chiasmocleis jimi*. 1: Type locality: Humaitá, AM (Caramaschi and Cruz 2001); 2: Paratype: Uruá, PA (Caramaschi and Cruz 2001); 3: Itaituba, PA (Peloso and Sturaro 2008); 4: Anapu, PA (Peloso and Sturaro 2008); 5: FLONA Caxiuanã, PA (Ribeiro-Jr et al. 2011); 6: Reserva Biológica do Rio Trombetas (Funk and Cannatella 2009); Red triangle: new record in the REBIOTA.




FIGURE 3. Geographic distribution of *Proceratophrys concavitympanum*. 1: Type Locality: Espigão D'Oeste, RO (Giaretta et al. 2000); 2: UHE Lajeado, TO (Amaro et al. 2009); 3: UHE Luis Eduardo Magalhães, TO (Pavan and Dixo, 2004); 4: RESEX Riozinho do Anfrízio, Altamira, PA (Barros et al. 2011); 5: Cacoal, RO (Santana et al. 2010); 6: Ministro Andreazza, RO (Santana et al. 2010); 7: Rondolândia, RO (Santana et al. 2010); 8: Aripuanã, MT (Santana et al. 2010); 9: Colniza, MT (Ávila et al. 2011); 10: Juína, MT (Ávila et al. 2011); 11: Paranaíta, MT (Ávila et al. 2011); 12: Jacareacanga, PA (Ávila et al. 2011). Red triangle: new record in the REBIOTA.

ACKNOWLEDGMENTS: This work was part of the Management Plan of the Reserva Biológica do Tapirapé elaborated by Ambiental Consulting and Instituto Chico Mendes para Biodiversidade (ICMBio) and funded by ARPA. We thank the managers of REBIOTA R. Façanha and P. Farina for all their support during and after fieldwork. We also thank S. Steinmetz, N. Pedroso Jr. and the Ambiental Consulting team for their support during fieldwork. We thank Desivaldo, Tatuquira, Darlan, Adair, Borges, Taironi, Gilberto, and all the park rangers for their invaluable help during fieldwork. We also thank to C. Blair (Royal Ontario Museum and University of Toronto, Canada), R. MacCulloch (Royal Ontario Museum) and D. Baêta (Museu Nacional do Rio de Janeiro) for the review and suggestions for improving this paper.

LITERATURE CITED

- Amaro, R.C, D. Pavan and M.T. Rodrigues. 2009. On the generic identity of *Odontophrynus moratoi* Jim & Caramaschi, 1980 (Anura, Cycloramphidae) *Zootaxa* 2071:61-68.
- Avila, R.W., R.A. Kawashita-Ribeiro and D.H. Morais. 2011. A new species of *Proceratophrys* (Anura: Cycloramphidae) from western Brazil. *Zootaxa* 2890: 20-28.
- Barros, F.B., H.M. Pereira and L. Vicente. 2011. Amphibia, Anura, Cycloramphidae, *Proceratophrys concavitympanum* Giaretta, Bernarde and Kokubum, 2000: Distribution extension for Brazilian Amazonia and first record in the state of Pará. *Check List* 7(2): 110-111.
- Caramaschi, U. 2004. *Chiasmocleis jimi*. In IUCN. *IUCN Red List of Threatened Species. Version 2011.1*. <www.iucnredlist.org>. Captured on 01 November 2011.
- Caramaschi, U. and C.A.G. Cruz. 2001. A New species of *Chiasmocleis* Méhely, 1904 from Brazilian Amazonia (AMPHIBIA, ANURA, MICROHYLIDAE). *Boletim do Museu Nacional, nova série, Zoologia* 469: 1-8.
- Frost, D.R. 2011. *Amphibian Species of the World: an Online Reference. Version 5.5 (31 January, 2011)*. Electronic Database accessible at <http://research.amnh.org/vz/herpetology/amphibia/>. American Museum of Natural History, New York, USA. Captured on 1 November 2011.
- Funk, W.C. and D. Cannatella, 2009. A new, large species of *Chiasmocleis* Méhely 1904 (Anura: Microhylidae) from the Iquitos region, Amazonian Peru. *Zootaxa* 2247: 37-50.
- Giaretta, A.A., P.S. Bernarde and M.C.N. Kokubum. 2000. A new species of *Proceratophrys* (Anura: Leptodactylidae) from the Amazon Rain Forest. *Journal of Herpetology* 34(2):173-178.
- IBGE. 1992. *Manual Técnico da Vegetação Brasileira*. Rio de Janeiro: Instituto Brasileiro de Geografia e Estatística. 92p.
- MMA. 2007. *Áreas Prioritárias para Conservação, Uso Sustentável e Repartição de Benefícios da Biodiversidade Brasileira: Atualização - Portaria MMA nº9, de 23 de janeiro de 2007*. Série Biodiversidade, v. 31. Brasília: Ministério do Meio Ambiente, Secretaria de Biodiversidade e Florestas. 300 p.
- Pavan, D. and M. Dixo. 2004. A Herpetofauna da área de influência do reservatório da Usina Hidrelétrica Luís Eduardo Magalhães, Palmas, TO. *Humanitas* 4/6: 13-30.
- Peloso, P.L.V. and M.J. Sturaro. 2008. A new species of narrow-mouthed frog of the genus *Chiasmocleis* Méhely 1904 (Anura, Microhylidae) from the Amazonian rainforest of Brazil. *Zootaxa* 1947:39-52.
- Prado, G.M. and J.P. Pombal Jr. 2008. Espécies de *Proceratophrys* Miranda-Ribeiro 1920 com apêndices palpebrais (Anura; Cycloramphidae). *Arquivos de Zoologia* 39(1):1-85.
- Ribeiro-Júnior M.A., R.V. Rossi, C.L. Miranda and T.C.S. Ávila-Pires. 2011. Influence of pitfall trap size and design on herpetofauna and small mammal studies in Neotropical Forest. *Zoologia* 28(1): 80-91.
- Rodrigues, M.T. and U. Caramaschi. 2004. *Proceratophrys concavitympanum*. In IUCN 2011. *IUCN Red List of Threatened Species. Version 2011.1*. <www.iucnredlist.org>. Captured on 01 November 2011.
- Santana, D.J., V.A. São-Pedro, P.S. Bernarde and R.N. Feio. 2010. Descrição do canto de anúncio e dimorfismo sexual em *Proceratophrys concavitympanum* Giaretta, Bernarde & Kokubum, 2000. *Papéis Avulsos de Zoologia* 50(11): 167-174.

RECEIVED: December 2011

ACCEPTED: January 2012

PUBLISHED ONLINE: February 2012

EDITORIAL RESPONSIBILITY: Ross McCulloch