

# Amphibia, Anura, Hylidae, *Phyllomedusa tarsius* Cope, 1868: Distribution extension, new country record and geographic distribution map

Maurício C. Forlani<sup>1\*</sup>, Pedro H. Bernardo<sup>1,2</sup> and Hussam Zaher<sup>1</sup>

<sup>1</sup> Universidade de São Paulo, Museu de Zoologia, Avenida Nazaré, 481. CEP 04263-000. São Paulo, SP, Brazil.

<sup>2</sup> Royal Ontario Museum, Centre for Biodiversity and Conservation Biology, 100 Queen's Park, Toronto, ON, M5S 2C6, Canada.

\* Corresponding author. E-mail: [mcforlani@gmail.com](mailto:mcforlani@gmail.com)

**ABSTRACT:** We provide a review of the geographic distribution of *Phyllomedusa tarsius*. A new record from Guyana corresponds to the eastern limit of the range for the species, which now includes the Guiana shield.

*Phyllomedusa tarsius* Cope, 1868 is a large Phyllomedusine of the *P. tarsius* group, widespread in the Western portion of the Amazon Basin (Duellman 1974; Frost 2011). It was originally described based on a single specimen, now presumably lost (Duellman 1974), collected on the margin of the Rio Amazonas (Amazon River), below the mouth of the Napo River, Loreto, Peru. Duellman (1974) reviewed previous works concerning the genus *Phyllomedusa* and synonymized *P. nicefori*, *P. edentula* and *P. orcesi* with *P. tarsius*, providing a complete description and distribution range for the species. According to Duellman (1974), *P. tarsius* only occurs in the upper Amazon Basin, in southern Colombia, Ecuador and Peru.

The *Phyllomedusa tarsius* group was first proposed by De la Riva (1999) when describing *P. camba*. In this paper the author groups *P. boliviana*, *P. camba*, *P. coelestis*, *P. tarsius*, *P. trinitatis* and *P. venusta* in the *tarsius* group. Studying the phylogenetic relationships within hylids, Faivovich *et al.* (2005) included *P. sauvagii* in the *P. tarsius* group, but the only species analyzed by the authors was *P. tarsius*. Barrio-Amorós (2006) suggested excluding *P. boliviana* and *P. camba* (included by De la Riva 1999) and *P. sauvagii* (included by Faivovich *et al.* 2005) because those species do not share what the author suggests as synapomorphy of the group: the golden iris with black reticulations. In a phylogenetic context Faivovich *et al.* (2010) tested the hypotheses of Barrio-Amorós (2006) for *P. tarsius* group and recovered the species *P. neildi*, *P. tarsius*, and *P. trinitatis* as a monophyletic group, excluding *P. boliviana* and *P. sauvagii*, and defining *P. camba* as sister-group.

The first record of *P. tarsius* in Venezuela was given for the province of Barinas (La Marca 1996). Duellman (1997) and Markezich (1998) further recorded the species in the provinces of Bolívar and Portuguesa, respectively. In Brazil, the species was recorded in the region of Manaus (Schmidt and Amézquita 2001), and in the Reserva Adolpho Ducke


(Lima *et al.* 2006), both in Amazonas, and in the state of Rondônia (Bernarde 2007). Doan and Arriaga (2002) recorded *P. tarsius* in the Tambopata region, Peru. The record for Guyana by Reynolds *et al.* (2001) (see also Señaris and MacCulloch 2005) is problematic because it is based on a specimen without exact locality data, and whose identity has not been confirmed by the authors (R. MacCulloch pers. comm.). Here we present two new records of *P. tarsius* in Guyana, which are the first confirmed records for this country (Figure 1).

On March 24<sup>th</sup> 2011, in the vicinity of the Kuribrong River, Potaro-Siparuni District, Guyana, on the access road to the Powis Mining Camp, we collected two male specimens of *P. tarsius* on vegetation 1.5 m above the ground and near a pond on the edge of the forest (05°30'56" N; 59°03'25" W, elevation 130 m). The second collection (one male and one female) occurred on April 4<sup>th</sup> 2011, at a temporary pond beside a road (05°33'37" N; 59°17'36" W, elevation 73 m). The male was calling on vegetation approximately 1.4 m above ground, while the female was on a nearby branch 1.7 m above ground. The vegetation type in the area corresponds to a "terra-firme" forest dominated by large trees. The understory is not very dense, except in regeneration areas. These records extend the known geographic distribution of the species ca. 247 km southeast. The region is under strong anthropogenic pressure due to mining and deforestation activities. All specimens were collected under the permit reference # 030510 BR 130, issued by the Guyana Environmental Protection Agency.

In the same location we found two other species of *Phyllomedusa* (*P. bicolor* and *P. tomopterna*) and at a nearby site we found *P. vaillantii*. *Phyllomedusa hypochondrialis* also occurs in the area. *Phyllomedusa tarsius* can be easily distinguished from these four species by its size (*P. tomopterna* and *P. hypochondrialis* are much smaller while *P. bicolor* is larger and *P. vaillantii* smaller), the flanks and internal surfaces of hind limbs that are marked with

individual or confluent cream and white spots (orange or reddish with vertical black stripes on *P. tomopterna* and *P. hypochondrialis*; cream, orange or reddish ocelli in *P. vaillantii*), and the presence of a golden iris with black reticulations (iris silvery or grayish on *P. bicolor*, *P. tomopterna*, *P. hypochondrialis* and *P. vaillantii*).

All specimens (Figure 2) agree with the description



**FIGURE 1.** Geographic distribution of *Phyllomedusa tarsius*. Red circles: records from literature. Salmon shade: Geographic distribution according to IUCN (La Marca et al. 2004). Yellow triangles: new records in Guyana.

given by Duellman (1974), and exhibit the following relevant characteristics: iris bright-orange with bold black reticulations; dorsum slightly granular; first finger shorter than and opposable to second finger; first toe longer than and opposable to second toe; discs on fingers large; forearm much more robust than upper arm; calcar absent; anal opening directed posteroventrally at mid-level of thighs; dorsal surfaces and side of head green; flanks marked by discrete or confluent cream spots; white spot present on chest; pair of white para-anal spots present; throat and chest grayish brown; margin of lower lip dull cream (Figure 2). The SVL for the males agrees with Duellman's (1974) description (mean=82.03mm; n=3) although the female is smaller (89.92 mm). Specimens were deposited in the herpetological collection of the Centre for the Study of Biological Diversity at the University of Guyana under the numbers UGCSBD HA1000, 1001, 1100, 1101.

According to the International Union for Conservation of Nature (IUCN), *P. tarsius* is listed as "Least Concern" due to its wide distribution, tolerance of habitat modification, and a presumed large population size (La Marca et al. 2004). Considering the morphological variation of *P. tarsius* and its known widespread geographic distribution, a detailed taxonomic review is recommended.



**FIGURE 2.** Specimens of *Phyllomedusa tarsius* collected in Guyana. A: UGCSBD HA1000 B: UGCSBD HA1001.

**ACKNOWLEDGMENTS:** We thank JGP Consulting for logistic support during the fieldwork; E. Hingst-Zaher (Museu Biológico, Instituto Butantan, Brazil) and C. Blair (Royal Ontario Museum and University of Toronto, Canada) for the review of the manuscript, Ross MacCulloch (Royal Ontario Museum, Canada) for the information about this species in Guyana, and Pedro Peloso (American Museum, USA) and an anonymous reviewer for their comments and suggestions in a previous draft of this manuscript.

#### LITERATURE CITED

Barrio-Amorós, C. 2006. A new species of *Phyllomedusa* (Anura: Hylidae: Phyllomedusinae) from Northwestern Venezuela. *Zootaxa* 1309: 55-

68.

Bernarde, P. 2007. Ambientes e temporada de vocalização da anurofauna no Município de Espigão do Oeste, Rondônia, Sudoeste da Amazônia-Brasil (Amphibia: Anura). *Biota Neotropica* 7(2): 87-92.

De la Riva, I. 1999. A new *Phyllomedusa* from Southwestern Amazonia (Amphibia: Anura: Hylidae). *Revista Espanhola de Herpetología* 13: 123-131.

Doan, T. and W. Arriaga. 2002. Microgeographic variation in species composition of the herpetofaunal communities of Tambopata region, Peru. *Biotropica* 34(1): 101-117.

Duellman, W.E. 1974. Taxonomic notes on *Phyllomedusa* (Anura: Hylidae) from the upper Amazon Basin. *Herpetologica* 30 (2):105-112.

- Duellman, W.E. 1997. Amphibians of La Escalera Region, Southeastern Venezuela: Taxonomy, Ecology and Biogeography. *Scientific Papers, National History Museum, University Kansas* 2: 1-52.
- Faivovich, J., C.F.B. Haddad, P.C.A. Garcia, D.R. Frost, J.A. Campbell, and W.C. Wheeler. 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: a phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294: 1-240.
- Faivovich, J., C.F.B. Haddad, D. Baêta, K.H. Jungfer, G.F.R. Álvares, R.A. Brandão, C.A. Sheil, L.S. Barrientos, C.L. Barrio-Amorós, C.A.G. Cruz and W.C. Wheeler. 2010. The phylogenetic relationships of the charismatic poster frogs, Phyllomedusinae (Anura, Hylidae). *Cladistics* 26: 227-261.
- Frost, D.R. 2011. *Amphibian Species of the World: an Online Reference*. Version 5.5 (31 January, 2011). Electronic Database accessible at <http://research.amnh.org/vz/herpetology/amphibia/>. American Museum of Natural History, New York, USA. Captured on 6 July 2011.
- La Marca, E. 1996. Geographic distribution: *Phyllomedusa tarsius*. *Herpetological Review* 27: 149.
- La Marca, E. C. Azevedo-Ramos, L.A. Coloma, S. Ron and C.L. Barrio-Amorós. 2004. *Phyllomedusa tarsius*. In IUCN. *IUCN Red List of Threatened Species*. Electronic Database accessible at <http://www.iucnredlist.org>. Captured on 06 July 2011.
- Lima, A.P., W.E. Magnusson, M. Menin, L.K. Erdtmann, D.J. Rodrigues, C. Keller and W. Hödl. 2006. *Guia dos Sapos da Reserva Adolpho Ducke - Amazônia Central /Guide to the frogs of Reserva Adolpho Ducke - Central Amazonia*. Manaus: Attama 168 p.
- Markezich, A. 1998. Geographic Distribution: Anura: *Phyllomedusa tarsius*. *Herpetological Review* 29 (1): 49.
- Reynolds, R., R. MacCulloch, M. Tamessar, C. Watson, C. J. Cole, and C. Townsend. 2001. *Preliminary checklist of the herpetofauna of Guyana. Biological Diversity of the Guiana Shield Program*, Electronic database accessible at <http://botany.si.edu/bdg/guyherps.html>. Smithsonian Institution. Captured on 05 July 2011.
- Schmidt, B. and A. Amézquita. 2001. Predator-induced behavioural responses: tadpoles of the neotropical frog *Phyllomedusa tarsius* do not respond to all predators. *Herpetological Journal* 11(1): 9-16.
- Señaris, J.C. and R. MacCulloch. 2005. Amphibians. p. 9-23 In T. Hollowell and R. Reynolds (ed.). Checklist of the terrestrial vertebrates of the Guiana shield. *Bulletin of the Biological Society of Washington* 13: 1-98.

RECEIVED: October 2011

ACCEPTED: January 2012

PUBLISHED ONLINE: February 2012

EDITORIAL RESPONSIBILITY: Ross McCulloch