

Ameerega flavopicta (Lutz, 1925): First dart-poison frog (Anura: Dendrobatidae) recorded for the state of São Paulo, Brazil, with comments on its advertisement calls and taxonomy

Lucas Borges Martins^{1,2,*} and Ariovaldo Antonio Giaretta¹

1 Universidade Federal de Uberlândia, Faculdade de Ciências Integradas do Pontal, Laboratório de Taxonomia, Sistemática e Ecologia de Anuros Neotropicais. Rua 20, 1.600 - Bairro Tupã, CEP 38304-402. Ituiutaba, MG, Brazil.

2 Universidade de São Paulo, FFCLRP/Departamento de Biologia, Programa de Pós-Graduação em Biologia Comparada. Avenida dos Bandeirantes, 3900. CEP 14040-901. Ribeirão Preto, SP, Brazil.

* Corresponding author. E-mail: lucasborgesmartins@hotmail.com

ABSTRACT: *Ameerega flavopicta* is a dart-poison frog (Dendrobatidae) widely distributed throughout rocky habitats in the Brazilian states of Minas Gerais, Goiás, Tocantins, Pará, and Maranhão. Here we report for the first record of a dendrobatid frog in the State of São Paulo, Brazil, based on individuals of *A. flavopicta* found at the Municipality of Pedregulho, and also describe the advertisement calls of a male from this population. A taxonomic discussion is provided.

Ameerega flavopicta (Lutz, 1925) (Dendrobatidae: Colostethinae) is a dart-poison frog described from Belo Horizonte, State of Minas Gerais, southeastern Brazil. Currently it is known to be widely distributed throughout the Brazilian states of Minas Gerais, Goiás, Tocantins, Pará, and Maranhão (see geographic distribution map and related references in Figure 1). This species is regarded as a diurnal breeder, usually found in rocky fields or riparian forests, along streams (Haddad and Martins 1994; Toledo *et al.* 2004; Costa *et al.* 2006; Magrini *et al.* 2010).

Herein we report for the first observation of *A. flavopicta* in the State of São Paulo, southeastern Brazil, and also describe the advertisement call of a male from this locality. We also discuss the taxonomic implications of our findings.

During recent fieldworks (December 2008 and October 2011) at the region of the Jaguara hydroelectric dam (Grande river), Municipality of Pedregulho, State of São Paulo, southeastern Brazil (southern margin of the dam), we noticed *Ameerega flavopicta* males calling in open rocky habitats, along temporary streams that drain into the dam. One calling male was collected on 23 October 2011, at 13:05 h, and is housed at the Amphibian Collection of the Universidade Federal de Uberlândia (Figure 2; voucher # AAG-UFU 0659; collecting permit ICMBio/SISBIO # 30059-2; collection GPS point: 20°08'20" S, 47°18'31" W). Specimens of *A. flavopicta* were also commonly found in the northern margin of the dam (Municipality of Sacramento, State of Minas Gerais), where one specimen was collected (27 December 2008, ≈16:30 h; collection GPS point: 20°07'03" S, 47°18'57" W; voucher # AAG-UFU 4398).

The anuran fauna of the State of São Paulo is considered to be the most thoroughly surveyed in Brazil, in relation to the other states (Araújo *et al.* 2009; Rossa-Feres *et al.* 2011), but two regions still deserves additional studies




FIGURE 1. Geographic distribution of *Ameerega flavopicta*, as presently recognized. Red dot depicts the type-locality, Municipality of Belo Horizonte, State of Minas Gerais. Yellow dots represent records from the present study: municipalities of Sacramento, State of Minas Gerais (above) and Pedregulho, State of São Paulo (below). Black dots depict localities previously reported in the literature (Haddad and Martins 1994; Biavati *et al.* 2004; Toledo *et al.* 2004; Silveira 2006; Leite *et al.* 2008; Oda *et al.* 2009; Magrini *et al.* 2010; Torres and Eterovick 2010): State of Pará: 1- Parauapebas; State of Maranhão: 2- Amaro Leite; State of Tocantins: 3- Paranã; State of Goiás: 4- Minaçu, 5- Alto Paraíso de Goiás, 6- Niquelândia, 7- Formosa, 8- Pirenópolis, 9- Serra Dourada, 10- Caldas Novas; State of Minas Gerais: 11- Araguari, 12- Perdizes, 13- São Roque de Minas, 14- Alpinópolis, 15- João Pinheiro, 16- Três Marias, 17- Santana do Riacho, 18- Jaboticatubas.

(Rossa-Feres *et al.* 2011): the southwest of the state, around the Paranapanema river basin, and the northeast of the state, around the border with the State of Minas Gerais. The site where we found *A. flavopicta* is within this latter region, reinforcing the need of additional field studies there. Besides *A. flavopicta*, five other anuran species were recently registered for the first time in the State of São Paulo from this region: *Rhinella rubescens* (Lutz, 1925), *Phyllomedusa ayeaye* (Lutz, 1966), *Scinax canastrensis* (Cardoso and Haddad, 1982) and *Barycholos ternetzi* (Miranda-Ribeiro, 1937), reported by Araujo *et al.* (2007a,b; 2009), and *Leptodactylus siphax* Bokermann, 1969 reported by Martins and Silva (2009).



FIGURE 2. An adult *Ameerega flavopicta* collected at the Municipality of Pedregulho, State of São Paulo, Brazil, on 23 October 2011 (photo of live individual). Voucher housed at the Amphibian Collection of the Universidade Federal de Uberlândia, # AAG-UFU 0659.

At Pedregulho, an unvouchered male had advertisement calls recorded on 23 October 2011, around 11:35 h. Recordings were made using a M-audio Microtrack II digital recorder coupled to a Sennheiser ME66/K6 directional microphone, set to 48 KHz and 16 bit resolution. Sounds were analyzed using SoundRuler (Gridi-Papp 2007), and sonograms conected using the package Seewave (Sueur *et al.* 2008) of the R software (R Development Core Team 2010).

Analyzed calls ($N = 15$; Figure 3) consisted of a pulsed note (7.4 ± 0.7 ; 6–9 pulses), that lasts 107.0 ± 6.7 (91.3–124.2) ms, with a slight increase in frequency: dominant frequency of 3589 ± 30 (3561–3608) Hz at the start, and 3802 ± 33 (3749–3842) Hz in the end of the note. This note is repeated at a rate of about 145 notes per minute, with inter-note intervals of 300.1 ± 29.8 (272.5–403.1) ms.

Advertisement calls of *A. flavopicta* were already described for others populations (see a compilation of these data in Table 1). Our data, in general, agree with these descriptions, but a larger sample size is needed for a formal, statistically based comparison between them, and it is beyond the aim of the present work.

Costa *et al.* (2006) reported geographical variation in the advertisement calls and tadpoles of specimens of *A. flavopicta* from Caldas Novas, State of Goiás, in relation to those previously described from Santana do Riacho, State of Minas Gerais (Haddad and Martins 1994). Later, Magrini *et al.* (2010) based on broader samples, reported significant variation in the advertisement calls of specimens from Caldas Novas, when compared to populations from Minas Gerais.

Haddad and Martins (1994) reported distinctive morphological variation in specimens of *A. flavopicta* from the State of Pará, and commented that future bioacoustic analyses should help in the clarification of the taxonomic status of this population, but their advertisement calls are still unknown.

Lötters *et al.* (2009) described *Ameerega boehmei* Lötters, Schmitz, Reichle, Rödder, and Quennet, 2009 based on specimens previously considered to be from a Bolivian population of *A. flavopicta*. This species is genetically and morphologically very similar to *A. flavopicta*, but clearly differs in advertisement call features (Lötters *et al.* 2009).

All these studies are suggestive of the need of a broader taxonomic evaluation of the populations of *Ameerega flavopicta*, as variation in advertisement calls and larval morphology were already reported, and the populations from the states of Pará, Tocantins and Maranhão remain poorly known. Integrative approaches including molecular, morphological and acoustic data, as done by Lötters *et al.* (2009), should help in clarifying the taxonomic status of these populations.




FIGURE 3. Spectrogram of an advertisement call of *Ameerega flavopicta*. Recorded at the region of the Jaguara hydroelectric dam, Municipality of Pedregulho, State of São Paulo, Brazil. 23 October 2011, 13:05 h; temperature data not available. Record file: Ameerega_flavopicSP1LBM_AAGmt.

TABLE 1. Advertisement call features of *Ameerega flavopicta*. Data from the present study and literature.

LOCALITIES AND LITERATURE SOURCES	NOTE	PULSES	INTER-NOTE	NOTES	DOMINANT FREQUENCIES OF THE NOTE	
	DURATION (ms)	PER NOTE	INTERVAL (ms)	PER MINUTE	BEGINNING	END
State of São Paulo:						
Pedregulho (present study)	107	7.4	300	145	3589	3802
	91–124	6–9	273–403		3561–3608	3749–3842
State of Minas Gerais:						
Alpinópolis (Haddad <i>et al.</i> 1988)	-----	6	550	-----	Frequency band from 3100 to 3700	Frequency band from 3400 to 4000
Santana do Riacho (Haddad and Martins 1994)	110	6	480–630	-----	Frequency band from 3200 to 4200 (modulation confirmed, but not analyzed)	
Santana do Riacho (Magrini <i>et al.</i> 2010)	150	6–10	460	78–129	3203	3503
	110–180		330–690		2878–3457	3292–3705
Araguari (Magrini <i>et al.</i> 2010)	110	7–9	310	123–146	3708	4045
	100–130		250–420		3457–3981	3788–4285
Perdizes (Magrini <i>et al.</i> 2010)	170	6–9	430	94–109	3561	3889
	160–180		360–500		3181–3954	3374–4340
State of Goiás:						
Alto Paraíso de Goiás (Toledo <i>et al.</i> 2004)	92	4–6	308	169	Dominant frequency \approx 3700–3800 (modulation confirmed, but not analyzed)	
	71–98		237–486			
Caldas Novas (Costa <i>et al.</i> 2006)	144	7–8	292	139	3200	4050
Caldas Novas (Magrini <i>et al.</i> 2010)	110	6–8	260	140–175	3723	4016
	90–150		230–290		3319–4119	3650–4367

ACKNOWLEDGMENTS: We thank FAPEMIG and CNPq for financial support. We also thank CNPq for a doctoral fellowship (LBM) and a grant (AAG). J. L. Brown and P. J. R. Kok provided valuable technical and linguistic comments on the manuscript.

LITERATURE CITED

- Araujo, C.O., T.H. Condez and C.F.B. Haddad. 2007a. Amphibia, Anura, *Barycholos ternetzi*, *Chaunus rubescens*, and *Scinax canastrensis*: Distribution extension, new state record. *Check List* 3(2): 153-155.
- Araujo, C.O., T.H. Condez and C.F.B. Haddad. 2007b. Amphibia, Anura, *Phyllomedusa ayeaye* (B. Lutz, 1966): Distribution extension, new state record, and geographic distribution map. *Check List* 3(2): 156-158.
- Araujo, C.O., T.H. Condez and R.J. Sawaya. 2009. Anfíbios anuros do Parque Estadual das Furnas do Bom Jesus, sudeste do Brasil, e suas relações com outras taxocenoses no Brasil. *Biota Neotropica* 9(2): 77-98.
- Araújo, O.G.S., L.F. Toledo, P.C.A. Garcia and C.F.B. Haddad. 2009. The amphibians of São Paulo State, Brazil. *Biota Neotropica* 9(4): 197-209.
- Biavati, G.M., H.C. Wiederhecker and G.R. Colli. 2004. Diet of *Epipedobates flavopictus* (Anura: Dendrobatidae) in a Neotropical Savanna. *Journal of Herpetology* 38(4): 510-518.
- Costa, R.C., K.G. Facure and A.A. Giaretta. 2006. Courtship, vocalization, and tadpole description of *Epipedobates flavopictus* (Anura: Dendrobatidae) in southern Goiás, Brazil. *Biota Neotropica* 6(1): 1-9.
- Gridi-Papp, M. 2007. *Sound Ruler acoustic analysis. Version. 0.9.6.0.* Available at <http://soundruler.sourceforge.net>. Captured on 17 August 2007.
- Haddad, C.F.B., G.V. Andrade and A.J. Cardoso. 1988. Anfíbios anuros do Parque Nacional da Serra da Canastra, estado de Minas Gerais. *Brasil Florestal* 64: 9-20.
- Haddad, C.F.B. and M. Martins. 1994. Four species of Brazilian poison frogs related to *Epipedobates pictus* (Dendrobatidae): Taxonomy and natural history observations. *Herpetologica* 50(3): 282-295.
- Leite, F.S.F., F.A. Juncá and P.C. Eterovick. 2008. Status do conhecimento, endemismo e conservação de anfíbios anuros da Serra do Espinhaço, Brasil. *Megadiversidade* 4(1-2): 158-176.
- Lötters, S., A. Schmitz, S. Reichle, D. Rödder and V. Quenett. 2009. Another case of cryptic diversity in poison frogs (Dendrobatidae: *Ameerega*) — description of a new species from Bolivia. *Zootaxa* 2028: 20-30.
- Magrini, L., K.G. Facure, A.A. Giaretta, W.R. Silva and R.C. Costa. 2010. Geographic call variation and further notes on habitat of *Ameerega flavopicta* (Anura, Dendrobatidae). *Studies on Neotropical Fauna and Environment* 45(2): 89-94.
- Martins, L.B. and W.R. Silva. 2009. Amphibia, Anura, Leptodactylidae, *Leptodactylus syphax*: New state record. *Check List* 5(3): 433-435.
- Oda, F.H., R.P. Bastos, and M.A.C.S. Lima. 2009. Taxocenose de anfíbios anuros no Cerrado do Alto Tocantins, Niquelândia, Estado de Goiás: diversidade, distribuição local e sazonalidade. *Biota Neotropica* 9(4): 219-232.
- R development core team. 2010. *R: A Language and Environment for Statistical Computing, Version 2.10.1.* Available at <http://www.R-project.org>. Captured on 05 March 2010.
- Rossa-Feres, D.C., R.J. Sawaya, J. Faivovich, J.G.R. Giovanelli, C.A. Brasileiro, L. Schiesari, J. Alexandrino and C.F.B. Haddad. 2011. Anfíbios do Estado de São Paulo, Brasil: conhecimento atual e perspectivas. *Biota Neotropica* 11(1a): 1-19.
- Silveira, A.L. 2006. Anfíbios do município de João Pinheiro, uma área de cerrado no noroeste de Minas Gerais, Brasil. *Arquivos do Museu Nacional* 64(2): 131-139.
- Sueur, J., T. Aubin and C. Simonis. 2008. Seewave, a free modular tool for sound analysis and synthesis. *Bioacoustics* 18: 213-226.
- Toledo, L.F., L.D. Guimarães, L.P. Lima, R.P. Bastos and C.F.B. Haddad. 2004. Notes on courtship, egg-laying site, and defensive behavior of *Epipedobates flavopictus* (Anura, Dendrobatidae) from two mountain ranges of central and southeastern Brazil. *Phyllomedusa* 3(2): 145-147.
- Torres, P.F. and P.C. Eterovick. 2010. Anuran assemblage composition and distribution at a modified environment in Três Marias reservoir, southeastern Brazil. *Journal of Natural History* 44: 2649-2667.

RECEIVED: January 2012

ACCEPTED: March 2012

PUBLISHED ONLINE: June 2012

EDITORIAL RESPONSIBILITY: Philippe J. R. Kok