

Liophis dorsocorallinus Esqueda, Natera, La Marca and Ilija-Fistar, 2007 (Squamata: Dipsadidae): Distribution extension in southwestern Amazonia, state of Acre, Brazil

Jailini da Silva Araújo^{1*}, Moisés Barbosa de Souza¹, Thays de Andrade Farias¹, Danyella Paiva da Silva¹, Nathocley Mendes Venâncio¹, Júnior Marcos Lima Maciel² and Paulo Roberto Melo-Sampaio²

¹ Universidade Federal do Acre, Laboratório de Herpetologia – HerPet, Campus Universitário Áulio Gélío Alves de Souza. BR 364 km 04 Distrito Industrial. CEP 69915-900. Rio Branco, AC, Brasil.

² União Educacional do Norte – Uninorte – SICP - BR 364 km 02 - Alameda Hungria, 200 Jardim Europa II - CEP. 69.911-900. Rio Branco, AC – Brasil.

* Corresponding author. E-mail: jailini@gmail.com

ABSTRACT: We present a new distribution record of *Liophis dorsocorallinus* in the state of Acre, Brazil. The species was previously known only from three localities in Venezuela and Brazil.

The Neotropical genus *Liophis* is diverse, and the relationships between the more than 40 species it includes remain largely unknown (Dixon 1980; Fernandes *et al.* 2002). Although Zaher *et al.* (2009) proposed an amendment to the genera *Liophis* and *Erythrolamprus*, this was later refuted by Curcio *et al.* (2009).

Curcio *et al.* (2009) agree with other authors that *Liophis* is probably a polyphyletic genus, and thus the inclusion of the type species of the genera involved is highly desirable for analyses aiming to provide nomenclatural resolutions.

Until recently, *Liophis dorsocorallinus* (Figure 1) was only known from one locality in southwestern Venezuela (Esqueda *et al.* 2007). A subsequent publication reported one new location in western Brazil (Bernarde *et al.* 2011). Bernarde *et al.* (2011) also suggested that the specimen from southern Amazonas, Brazil, considered as *Liophis* sp. by França and Venancio (2010) is *Liophis dorsocorallinus*, but no additional information was provided.

Recently five additional specimens of *L. dorsocorallinus* (all females) were collected in Acre, Brazil. Specimens were collected under permits 27290-1 issued by SISBIO. We identified the specimens using the original description by Esqueda *et al.* (2007). Translated from the original description, Esqueda *et al.* (2007) stated:

"Cephalic scales black, with reddish edges (most evident in the frontal, internasals and prefrontal) on the anterior lateral portion of the parietal there is a red spot that extends to the supralabials; supralabials yellow, slightly sprinkled with black, head ventrally yellow, including also the first third of the body, then it becomes red with more or less square or rectangular black spots, interspersed with each other, which fuse the most to the cloaca (reddish the last). Dorsum reddish with black scale tips; conspicuous yellow line at the level of the first two rows of dorsolateral scales, subcaudals reddish; iris orange with black pupil".

Following these characters from Esqueda *et al.* (2007) we assigned this species to *Liophis dorsocorallinus* (for additional squamation comparisons see Table 1).



FIGURE 1. *Liophis dorsocorallinus* (UFAC-0378), female, SVL 460 mm, Parque Zoobotânico. Photo by: Paulo Roberto Melo-Sampaio


On 28 May 2008, 1000 h, during herpetofaunal sampling at Fazenda São Jorge I, municipality of Sena Madureira (09°26'11" S; 68°37'19" W) NMV found two specimens resting in the axil of a palm tree in a pasture.

On 28 November 2011 during sampling of road-killed herpetofauna on highway BR-364 between Rio Branco and Sena Madureira, PRMS and JMLM found an individual of *Liophis dorsocorallinus* near the city of Bujari (09°48'37.10" S, 67°59'6.90" W elevation 185m). Habitat adjacent to the highway was pastures and secondary forest.

On 6 January 2012 during herpetofaunal sampling at Parque Zoobotânico da Universidade Federal do Acre (09°57'26" S, 67°52'25" W) in the municipality of Rio Branco, we found a specimen in a pitfall trap with drift fence in old-growth secondary forest. On 1 February 2012, 0800 h, at the same locality we encountered another individual of *Liophis dorsocorallinus* foraging on the ground. All specimens were collected and deposited into the Coleção Herpetológica da Universidade Federal do Acre (UFAC-0322-0323, UFA-0378, UFA-0396 and UFA-0398).

TABLE 1. Comparison characters of *Liophis dorsocorallinus*. * Specimen with incomplete tail. ** Specimen road-killed. Dashes = data not available..

<i>Liophis dorsocorallinus</i>						
Character	Esqueda et al. 2007	UFAC-0322	UFAC-0323	UFAC-0378	UFAC-0396	UFAC-0398**
Supralabials	8 (4,5)	8 (4,5)	8 (4,5)	8 (4,5)	8 (4,5)	-
Infralabials	8–9 (6)	9	9	9	9	-
Preoculars	1_2	1	1	2	1	-
Postoculars	2	2	2	2	2	-
Temporals	1+2	1+2	1+2	1+2	1+2	-
Ventrals	147–153 (MALES) 151 (FEMALE)	149	145	152	153	-
Subcaudals	69–80 (MALES) 69 (FEMALE)	62	66	45*	61	68
SVL (cm)	35.5–47.4 (MALES) 49 (FEMALE)	52	39	46	49	-
TTL (cm)	48.5–62.6 (MALES) 66.4 (FEMALE)	67	62	56*	63	-

**FIGURE 3.** Geographic distribution of *Liophis dorsocorallinus*. Star = type-locality (Esqueda et al. 2007). Black dots: left, Riozinho da Liberdade (Bernarde et al. 2011); right, Boca do Acre (França and Venâncio 2010). New records (this study): Green dot – vicinity of Bujari city. Yellow dot – Parque Zoológico da Universidade Federal do Acre, Rio Branco. Blue dot – Fazenda São Jorge I, Sena Madureira.

These are the southernmost records of *Liophis dorsocorallinus* and extend its distribution 1960 km from the type locality and about 140 km southwest of the nearest locality, extending its distribution to the municipalities of Bujari, Rio Branco and Sena Madureira, eastern Acre (Figure 2).

ACKNOWLEDGMENTS: JSA and TAF thank CNPq / UFAC for financial support. PRMS and JMLM thanks to RAN / ICMBio by the permits. We thank Ross MacCulloch, Luis Felipe Esqueda and two anonymous reviewers for useful comments.

LITERATURE CITED

- Bernarde, P.S., R.A. Machado, and L.C.B. Turci. 2011. Herpetofauna da área do Igarapé Esperança na Reserva Extrativista Riozinho da Liberdade, Acre – Brasil. *Biota Neotropica* 11(3): 117-144.
- Curcio, F.F., V. Piacentini, D.S. Fernandes. 2009. On the status of the snake genera *Erythrolamprus* Boie, *Liophis* Wagler and *Lygophis* Fitzinger (Serpentes, Xenodontinae). *Zootaxa* 2173: 66-68.
- Dixon, J.R. 1980. The Neotropical colubrid snake genus *Liophis*. The generic concept. *Contributions in Biology and Geology Milwaukee Public Museum* 31: 1–40.
- Esqueda, L.F., M. Natera, E. La Marca and M. Ilija-Fistar. “2005” [2007]. Nueva especie de serpente (Reptilia: Colubridae: *Liophis*) de un bosque tropical relictual en el estado Barinas, Venezuela. *Herpetotropicos* 2: 95-103.

- Fernandes, D.S., V.J. Germano, R. Fernandes and F.L. Franco. 2002. Taxonomic status and geographic distribution of the lowland species of the *Liophis cobella* group with comments on the species from the Venezuelan Tepuis (Serpentes, Colubridae). *Boletim do Museu Nacional Nova Série Zoologia* 481: 1-14.
- França, F.G.R. and N.M. Venâncio. 2010. Reptiles and amphibians of a poorly known region in southeastern Amazonia. *Biotemas* 23 (3): 71-84.
- Pyron, R.A., F.T. Burbrink, G.R. Colli, A.N. Montes de Oca, L.J. Vitt, C.A. Kuczynski, and J.J. Wiens. 2011. The phylogeny of advanced snakes (Colubroidea), with discovery of a new subfamily and comparison of support methods for likelihood trees. *Molecular Phylogenetics and Evolution* 58: 329-342.
- Vidal, N., M. Dewynter and D.J. Gower. 2010. Dissecting the major American snake radiation: a molecular phylogeny of the Dipsadidae Bonaparte (Squamata, Caenophidia). *Comptes Rendus Biologies* 333: 48-55.
- Zaher, H., F.F. Grazziotin, J.E. Cadle, R.W. Murphy, J.C. Moura-Leite and S.L. Bonatto. 2009. Molecular phylogeny of advanced snakes (Serpentes, Caenophidia) with an emphasis on South American xenodontines: a revised classification and descriptions of new taxa. *Papéis Avulsos de Zoologia* 49: 115-153.

RECEIVED: February 2012

ACCEPTED: April 2012

PUBLISHED ONLINE: June 2012

EDITORIAL RESPONSIBILITY: Ross MacCulloch