

Amphibia, Anura, Hylidae, *Phyllomedusa camba* De la Riva, 1999: Distribution extension and geographic distribution map

Danusy Lopes Santos^{1,3*}, Sheila Pereira de Andrade¹, Elias Gonzaga de Castro¹ and Wilian Vaz-Silva^{1,2,3}

1 Centro Tecnológico de Engenharia Ltda., Rua 254, 146, Setor Coimbra. CEP 74535-440. Goiânia, GO, Brazil.

2 Centro Universitário de Goiás – Uni-Anhanguera, Departamento de Ciências Biológicas, Rua Professor Lázaro Costa, 456. CEP 74415-450. Goiânia, GO, Brazil.

3 Universidade Federal de Goiás, Instituto de Ciências Biológicas, Departamento de Ecologia. Rodovia Goiânia-Nerópolis Km 5, Campus II, Setor Itatiaia. CEP 74001-970. Goiânia, GO, Brazil.

* Corresponding author. E-mail: danusylopes@yahoo.com.br

ABSTRACT: This note reports a new record of *Phyllomedusa camba* De la Riva, 1999 in the municipality of Tangará da Serra, state of Mato Grosso, which represents the species easternmost known locality.

The genus *Phyllomedusa* currently comprises 30 species distributed in the Panamá, Pacific slopes of Colombia, and South America east of the Andes, including Trinidad, southward to northern Argentina and Uruguay (Frost 2011). According to Faivovich et al. (2005) the *Phyllomedusa tarsius* group comprises five species: *Phyllomedusa boliviana* Boulenger, 1902; *Phyllomedusa camba* De la Riva, 1999; *Phyllomedusa neildi* Barrio-Amorós, 2006; *Phyllomedusa sauvagii* Boulenger, 1882 and *Phyllomedusa tarsius* (Cope, 1868).

Phyllomedusa camba (Figure 1) has been reported to occur in the southwestern Amazon basin from southeastern Peru, western Brazil (states of Amazonas, Acre, and Rondônia) to eastern Bolivia (Frost 2011), and in state of Mato Grosso it was recorded in the municipalities of Aripuanã, Cotriguaçu, and Vila Bela da Santíssima Trindade (Rodrigues et al. 2011). *Phyllomedusa camba* inhabits primary lowland tropical moist forest, secondary forest, and also disturbed open areas (De la Riva et al. 2004).

Here we report a new record for *Phyllomedusa camba* in the municipality of Tangará da Serra (Figure 2), state of Mato Grosso. The new record in the municipality of Tangará da Serra, state of Mato Grosso (15°08'53" S, 57°27'36" W) extends the known distribution in 511 km from the type locality (Puerto Almacén, Departamento de Santa Cruz, Bolívia; 15°46' S, 62° 5' W), 568 km from Aripuanã, Mato Grosso, Brazil (10°09' S, 59°25' W), 573 km from Cotriguaçu, Mato Grosso (09°51'16.9" S, 58°14'57.7") and 200 km from the nearest locality, Vila Bela da Santíssima Trindade (14°59' S, 59°57' W).

Specimens were recorded in a temporary pond surrounded by pastures during a field expedition on 17 December 2011 for the biodiversity inventory of the hydroelectric power plant on the Juba River. Males were

observed calling in high trees. The collected specimens was euthanized with 5% lidocaine gel, fixed in 10% formalin and preserved in alcohol 70%, tissue samples were taken. Voucher specimens are deposited at the Coleção Zoológica da Universidade Federal de Goiás (ZUFG), Goiânia, Goiás, Brasil, under the labels ZUFG 6443- 6446 (Collection permit 57/2011- SUPES/MT).

Reports on the geographical distribution and its potential occurrences are extremely necessary for formulation of conservation politics. The emergence of new distribution records shows how data on the Brazilian amphibians are deficient. Information on *Phyllomedusa camba* are scarce, so this new record contributes to the knowledge of their distribution in the state of Mato Grosso. Additionally, more studies should be conducted for a better understanding of the biology of the species.



FIGURE 1. Voucher specimen of *Phyllomedusa camba* (ZUFG-6443) from Tangará da Serra municipality, state of Mato Grosso.




FIGURE 2. Geographic distribution of *Phyllomedusa camba* (shaded delimited area, IUCN, 2010) and localities where the species has already been registered: polygon: Puerto Almacén, Bolívia (15°46' S, 62°05' W), type-locality; circle: Aripuanã (10°09' S, 59°25' W); triangle: municipality of Cotriguaçu (09°51'16.9" S, 58°14'57.7" W); square: Vila Bela da Santíssima Trindade (14°59' S, 59°57' W); star: new record in Tangará da Serra (15°08'53.1" S, 57°27'36.4" W).

ACKNOWLEDGMENTS: We are grateful to Wanessa Batista, for field assistance. Roberto Leonan Novaes for assistance in the preparation of the map. CTE Engenharia is the enterprise responsible for carrying out the Faunal Inventory on the Juba River.

LITERATURE CITED

De la Riva, I. 1999. A new *Phyllomedusa* from southwestern Amazonia (Amphibia: Anura: Hylidae). *Revista Española de Herpetology* 13: 123-131.

De la Riva, I., K-H. Jungfer, A. Angulo and S. Reichle. 2004. *Phyllomedusa camba*. In IUCN 2011. *IUCN Red List of Threatened Species. Version 2009.2*. Electronic Database accessible at www.iucnredlist.org. Captured on 30 December 2011.

Faivovich, J., C.F.B. Haddad, P.C.A. Garcia, D.R. Frost, J.A. Campbell and W.C. Wheeler. 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin American Museum of Natural History* 294: 1-113.

Frost, D.R. 2011. *Amphibian Species of the World: an online reference. Version 5.5*. Electronic Database accessible at <http://research.amnh.org/vz/herpetology/amphibia/>. Captured on 30 December 2011.

Rodrigues, D.D.J., Noronha, J.D.C., Lima, M.D.M.L., Rosa, A.C. 2011. Amphibia, Anura, Hylidae, *Phyllomedusa boliviana* Boulenger, 1902 and *Phyllomedusa camba* De la Riva, 2000: Distribution extension in central Brazil. *Check List* 7(4): 397-399.

RECEIVED: February 2012

ACCEPTED: May 2012

PUBLISHED ONLINE: August 2012

EDITORIAL RESPONSIBILITY: Fernanda Werneck