

Polychaeta (Annelida) from the continental shelf off Aveiro (NW Portugal): Species composition and community structure

Ascensão Ravara* and Maria Helena Moreira

CESAM, Departamento de Biologia, Universidade de Aveiro, Campus de Santiago, 3810-193 Aveiro, Portugal.

* Corresponding author. E-mail: aravara@ua.pt

ABSTRACT: The present study provides a checklist of the polychaete fauna collected on the continental shelf off Aveiro (NW Portugal), within an area located between latitudes 40°30'N and 40°50'N and longitudes 8°40'W and 9°20'W, and a depth range of 8 to 185 m. The list includes 136 species, belonging to 37 families, from which 19 are new records for the Portuguese coast (*Isolda pulchella*, *Mesochaetopterus sagittarius*, *Aphelochaeta multibranchis*, *Chaetozone* cf. *vivipara*, *Diplocirrus hirsutus*, *Goniadella gracilis*, *Gyptis mediterranea*, *Lumbrinerides crassicephala*, *Eumida bahusiensis*, *Eumida ockelmanni*, *Nereiphylla paretti*, *Phyllodoce rosea*, *Glyphohesione klatti*, *Malmgreniella arenicolae*, *Prionospio aluta*, *Pseudopolydora paucibranchisata*, *Pseudopolydora pulchra*, *Scolecopsis mesnili*, *Polycirrus* cf. *medusa*). The shelf off Aveiro presents a well-defined pattern of sediment distribution, with finer sand on the inner (8–22 m depth) and the outer (94–184 m depth) parts of the shelf and coarser sediments on the mid-shelf. The distribution of polychaete assemblages follows closely the aforementioned sedimentary pattern, showing different specific composition and structure in each of the three areas. Depth was also found to play an important role in the distribution of some polychaete species.

INTRODUCTION

Marine soft-bottom communities support a high diversity of species with different ecological characteristics. Many studies have been published on benthic communities of the Northeastern Atlantic continental shelves (Heip and Craeymeers 1995; Hoey *et al.* 2004; Mackie *et al.* 1997; Martínez and Adarraga 2001; Parapar and Moreira 2009; among others) and the Mediterranean (Ballesteros 2006; Cebrián and Ballesteros 2004; Çinar 2005; Martin *et al.* 2000; Mohamed 1993; Sardá *et al.* 2012; Somaschini *et al.* 1998; among others). Nevertheless, little is known about the benthic communities of the Portuguese continental coast, especially at higher depths. There are only two studies on polychaete fauna from the Portuguese continental slope (including the slope off Aveiro) at depths below 200 m (Amoureux 1974; 1987). A few other works are published for Portuguese coastal waters, the majority referring to near shore zones (Almaça 1960; Amoureux and Calvário 1981; Dexter 1988; 1990; França *et al.* 2009; Guerra-García *et al.* 2011; Moreira *et al.* 1993; Sousa-Reis *et al.* 1982), and others to South and Southwest of Portugal (Bellan 1960; Gil and Sardá 1999; Hartmann-Schröder 1977; 1979; 1981; Monteiro-Marques 1979; 1987; Pérès, 1959). The western Portuguese continental shelf is of particular interest due to the combined influence of the North Atlantic and Mediterranean currents, providing conditions to the occurrence of fauna characteristic from both regions (Sousa-Reis *et al.* 1982). Polychaetes (Annelida, Polychaeta) represent a very important part of the marine macrobenthic fauna, being often the most abundant and specious group. It is, therefore, crucial to improve the knowledge on polychaete fauna composition and distribution in this area. During the summers of 1994 and 1995, two oceanographic surveys were carried out

with the aim of studying these communities and their relation to some environmental parameters on the shelf off Aveiro. These surveys resulted from a collaboration between the universities of Aveiro and Bordeaux and were supported by a Portuguese-French oceanological program (JNICT, French embassy) and by the French CIRMAT (Comite Inter-Regional Manche-Atlantique) and CNRS (Centre National de la Recherche Scientifique). The present work provides information on the species composition, distribution and ecology of benthic polychaetes of shelf depths, increasing our knowledge on the biodiversity of the Portuguese coastal waters. The data was analyzed on the basis of species composition and density and interpreted according to habitat features, namely sediment type and depth.

MATERIALS AND METHODS

Study site

The continental shelf off Aveiro is located at the NW Portuguese continental margin. The shelf-width ranges between 50.2 km (in the northern part of the study area) and 38 km (near Aveiro Canyon) and the shelf-break lies at a mean depth of about 160 m. The bathymetrical contours are generally subparallel to the coastline, having a NNE-SSW orientation. The sediment cover of the shelf is dominated by sand arranged in deposits of different origins and characteristics. The near-shore deposits are formed by siliciclastic fine to very fine sand mostly supplied by southward alongshore drift from northern river born material. The mid-shelf deposits are formed by siliciclastic coarse sand and gravel originated by past littoral reworking of terrestrial deposits. These two areas (inner- and mid-shelf) have high energetic conditions at the bottom, promoting sediment remobilization and transport. At the

100 m isobath the occurrence of oceanic thermal fronts benefit the deposition of finer sediments enriched in organic matter, which enhance the productivity of benthic fauna in this area (Martins *et al.* 2012). The subsequent outer-shelf and shelf-break deposits are formed by very fine and median sand rich in particles of biogenic origin (molluscan fragments, foraminifera tests) (Abrantes *et al.* 1994). Mud (silt and clay) content is generally low over the whole study area and increases seawards.

Data collection

The sampling took place during two oceanographic surveys carried out in July/August of 1994/95 onboard R/V “Côte d’Aquitaine” (CNRS-CIRMAT). A total of 31 sites were sampled along four transects perpendicular to the coast and covering an area between latitudes 40°30'N and 40°50'N and longitudes 8°40'W and 9°20'W, within a depth range of 8 to 185 m (Figure 1). The macrofauna was sampled with a Smith-McIntyre grab (0.1 m²) and three replicates were taken per site. Samples were fixed in 10% neutral formalin stained with rose Bengal, and later washed and the macrofauna transferred to 70% ethanol. After taxonomical identification and counting, the specimens were deposited in the Biological Research Collection of Marine Invertebrates of Universidade de Aveiro (DBUA), Portugal. Total organic matter content of surface sediments was determined by loss of weight on ignition at 450°C for 5 hours, using 1 g of dried sediment taken from each sample before fixation. For sedimentological studies an extra sample was collected at each site using a Reineck box corer, from which the upper 2 cm layer was analyzed. Sediment type was determined according to the Wentworth scale (Buchanan and Kain 1971).

Data analysis

The biological data was submitted to Non-parametric multivariate analyses using the PRIMER package (Carr *et al.* 1993). Following the prearrangement of the abundance

data into samples / species matrix, the similarity between all pairwise combinations of samples was calculated using the Bray-Curtis similarity coefficient after fourth-root transformation of the data. Hierarchical agglomerative clustering, using group-average linking, and non-metric MDS (Multi-Dimensional Scaling) ordination based on the Bray-Curtis similarity matrix were used to obtain graphical presentation of the faunal similarity changes in space. Diversity was evaluated by species richness and Shannon-Wiener index (H') (log base 2), while equitability was accessed by Pielou evenness index (J').

RESULTS AND DISCUSSION

A total of 136 species and 10,353 individuals were collected in this study. Table 1 presents a list of the species recorded, from which 19 are new records for the Portuguese coast (see species marked with a * in Table 1). For each species, the total number of individuals, habitat conditions in the study area (depth range, sediment type and organic matter content) and deposit number in the DBUA collection are given.

The classification and ordination analyses of the 31 sampled sites clustered three main groups at a similarity level higher than 30% (Figure 2). The three groups of sites occupy well-defined positions on the shelf off Aveiro corresponding to inner- (depth range from 8 to 22 m), mid- (depth range from 31 to 79 m) and outer-shelf (depth range from 94 to 184 m) (Figure 1). The mid-shelf group stands out from the others for its higher values of species richness and abundance, and for presenting coarser sediments (Table 2). The other two groups are much similar in faunal composition and are both characterized by finer sediment, although outer-shelf sites have higher organic matter content. The outer-shelf group also presents higher values of diversity and equitability. The distribution of the polychaete fauna along the shelf off Aveiro seems to be mainly related to sediment type and depth (Figure 3). Many species present a strong association with fine sediments, being very abundant on the inner-




FIGURE 1. Map of the study area on the NW Portuguese continental shelf off Aveiro with location of sampling sites. Site symbols correspond to the three clusters resulting from the classification analysis.




FIGURE 2. Classification and ordination analysis of the 31 sampled sites based on polychaete abundance data.

shelf, scarce or absent on the mid-shelf and occurring again on the outer-shelf, while other species have their distribution mainly restricted to the mid-shelf, where sediment is coarser. On the other hand, the distribution of some other species seems to be mostly determined by depth, independently of sediment type. Depth and sediment type have been pointed out by most authors as important factors affecting the distribution of marine macrobenthic communities (Hoey *et al.* 2004; Mackie *et al.* 1997; Martínez and Adarraga 2001; Parapar and Moreira 2009; Snelgrove 1998).

The comparison of the fauna collected in this area with that of the North Sea (Eleftheriou and Basford 1989; Heip and Craeymeers 1995; Hoey *et al.* 2004; Künitzer *et al.* 1992), denoted a large proportion of species in common with our study area. Furthermore, there was a similar relationship between the species distribution and the sediment type and depth in both areas. The same was observed in the coasts of France and Galicia (NW Spain) (Amoureux 1985; Martínez and Adarraga 2001; Parapar and Moreira 2009). Species like *Psammathe fusca*, *Laonice bahusiensis*, *Magelona filiformis*, *Paradiopatra hispanica*, *Diplocirrus stopbowitzi* and *Aphelocheata multibranchis* (among others), which are common in the shelf off Aveiro, are also frequent and abundant in those northern regions. Likewise, species such as *Parapionosyllis cabezli* and *Opisthodonta morena*, for example, occur mostly in the Mediterranean (Bellan 1962; 1964; 1965; Çinar 2005; Guille 1970) and are also common in the shelf off Aveiro. These findings suggest that the studied area is located in a transition zone where species characteristic of both septentrional and meridional areas can be found.




FIGURE 3. Depth distribution of the most abundant species. Dots size reflects the abundance values.

TABLE 1. List of species collected on the continental shelf off Aveiro. N, number of individuals; O.M., organic matter content; DBUA, deposit number on the DBUA collection; *, first record for Portugal. Sediment types: VFS, very fine sand; FS, fine sand; MS, median sand; CS, coarse sand; VCS, very coarse sand; G, gravel. The species *Oxydromus agilis* and *O. pallidus* (placed together in the table) require live organisms to be accurately distinguished.

LIST OF SPECIES	N	DEPTH RANGE (M)	SEDIMENT TYPE	O.M. (%)	DBUA
AMPHARETIDAE Malmgren, 1866					
<i>Ampharete finmarchica</i> (M. Sars, 1865)	9	68.9-184.2	FS, MS	0.85-4.21	00142
<i>Amphicteis gunneri</i> (M. Sars, 1835)	1	162.8	MS	3.06	00183
<i>Anobothrus gracilis</i> (Malmgren, 1866)	5	129.5-157.7	FS, MS	2.53-2.57	00143
<i>Isolda pulchella</i> F. Müller, 1858*	16	157.7-162.8	MS	2.57-3.06	00145
<i>Sosane sulcata</i> Malmgren, 1866	4	134.9-184.2	FS, MS	2.14-2.83	00144
AMPHINOMIDAE Savigny in Lamarck, 1818					
<i>Chloeia venusta</i> Quatrefages, 1866	15	35.2-184.2	MS, CS	0.58-2.83	00027
CAPITELLIDAE Grube, 1862					
<i>Mediomastus fragilis</i> E. Rasmussen, 1973	1052	8.9-162.8	FS, MS, CS, G	0.33-4.21	00126
<i>Notomastus latericeus</i> M. Sars, 1851	146	12.7-134.9	MS, CS, VCS, G	0.33-2.14	00127
<i>Notomastus profundus</i> Eisig, 1887	8	131.5-184.2	FS, MS	2.57-4.21	00128
CHAETOPTERIDAE Malmgren 1867					
<i>Mesochaetopterus sagittarius</i> (Claparède, 1870)*	25	15.6-184.2	MS, VCS, G	0.36-4.21	-
CIRRATULIDAE Ryckholt, 1851					
<i>Aphelocheata marioni</i> (Saint-Joseph, 1894)	34	39.3-184.2	FS, MS, VCS, G	0.36-4.21	-
<i>Aphelocheata multibranchis</i> (Grube, 1863)*	16	96.3-184.2	FS, MS	0.98-2.83	-
<i>Caulleriella alata</i> (Southern, 1914)	5	39.3-76.8	CS, G	0.57-1.46	-
<i>Caulleriella bioculata</i> (Keferstein, 1862)	34	30.8-76.8	CS, G	0.33-1.46	-
<i>Chaetozone setosa</i> Malmgren, 1867	37	48.3-162.8	FS, MS	0.76-4.21	-
<i>Chaetozone</i> cf. <i>vivipara</i> Christie, 1984*	4	129.5-131.5	FS	2.53-4.21	-
<i>Cirriiformia tentaculata</i> (Montagu, 1808)	5	31.8-157.7	CS, VCS, G	0.33-0.90	-
DORVILLEIDAE Chamberlin, 1919					
<i>Protodorvillea kefersteini</i> (McIntosh, 1869)	1075	30.8-124.2	CS, VCS, G	0.33-1.46	00052
<i>Schistomeringos neglecta</i> (Fauvel, 1923)	4	59.1-76.8	VCS	0.70-1.29	00053
EUNICIDAE Berthold, 1827					
<i>Eunice vittata</i> (Delle Chiaje, 1829)	10	35.2-157.7	MS, CS, VCS	0.58-2.57	00051
<i>Marphysa bellii</i> (Audouin and Milne Edwards, 1833)	12	31.8-57.3	CS, VCS, G	0.33-1.46	00050
FLABELLIGERIDAE Saint-Joseph, 1894					
<i>Diplocirrus hirsutus</i> (Hansen, 1878)*	2	21.9	VFS	0.90	-
<i>Diplocirrus stopbowitzi</i> Darbyshire and Mackie, 2009	17	48.2-79.0	VCS, G	0.70-1.29	01148
<i>Pherusa monilifera</i> (Delle Chiaje, 1841)	9	13.5-40.9	FS, VCS	0.36-0.74	-
GLYCERIDAE Grube, 1850					
<i>Glycera fallax</i> Quatrefages, 1850	27	30.8-184.2	MS, CS, VCS, G	0.36-3.06	00068
<i>Glycera lapidum</i> Quatrefages, 1866	330	30.8-184.2	CS, VCS, G	0.33-2.83	00070
<i>Glycera tridactyla</i> Schmarda, 1861	119	12.7-162.8	VFS, FS, MS	0.36-4.21	00071
GONIADIDAE Kinberg, 1866					
<i>Glycinde nordmanni</i> (Malmgren, 1866)	33	34.9-184.2	MS, VCS	0.36-2.83	00185
<i>Goniada emerita</i> Audouin and Milne Edwards, 1833	2	21.9-34.9	VFS, CS	0.42-0.90	00186
<i>Goniada maculata</i> Ørsted, 1843	16	57.3-162.8	FS, MS	0.76-4.21	00187
<i>Goniadella galaica</i> (Rioja, 1923)	63	15.6-79.0	CS, VCS, G	0.36-1.46	00188
<i>Goniadella gracilis</i> (Verrill, 1874)*	84	30.8-124.2	VCS, G	0.43-2.14	-
HESIONIDAE Grube, 1850					
<i>Gyptis mediterranea</i> Pleijel, 1993*	1	124.8	-	-	00100
<i>Oxidromus agilis</i> (Ehlers, 1864) / <i>Oxidromus pallidus</i> Claparède, 1864	25	35.2-76.8	CS, VCS	0.58-1.29	00101
<i>Psamathe fusca</i> Johnston 1836	96	12.7-97.5	CS, VCS, G	0.33-1.46	00102
LUMBRINERIDAE Schmarda, 1861					
<i>Lumbrinerides amoureuksi</i> Miura, 1980	19	30.8-76.8	MS, CS, VCS, G	0.33-2.14	00039
<i>Lumbrinerides crassicephala</i> Hartman, 1965*	9	124.6-162.8	FS, MS, G	2.14-4.21	01323
<i>Lumbrineris futilis</i> Kinberg, 1865	5	46.1-184.2	FS, CS, G	0.33-2.83	01322
<i>Lumbrineris lusitanica</i> Martins, Carrera-Parra, Quintino and Rodrigues, 2012	31	57.3-184.1	FS, MS, CS, VCS, G	0.78-4.21	00040
<i>Scoletoma</i> cf. <i>magnidentata</i> (Winsnes, 1981)	1	39.3	G	0.57	-
MAGELONIDAE Cunningham and Ramage, 1888					
<i>Magelona allenii</i> Wilson, 1958	5	13.5-124.2	FS	0.74-1.35	00083
<i>Magelona filiformis</i> Wilson, 1959	40	12.7-162.8	VFS, FS	0.36-3.06	00082
<i>Magelona johnstoni</i> Fiege, Licher and Mackie 2000	230	8.9-93.9	VFS, FS	0.36-1.03	00084
<i>Magelona minuta</i> Eliason, 1962	76	93.9-162.8	FS, MS	0.76-4.21	00081
MALDANIDAE Malmgren, 1867					
<i>Leiochone leiopygos</i> (Grube, 1860)	32	46.1-184.2	MS, CS, VCS, G	0.33-4.21	00129
<i>Maldane glebifex</i> Grube, 1860	2	129.5-157.7	MS, VCS	0.36-2.57	00130
NEPHTYIDAE Grube, 1850					

TABLE 1. CONTINUED.

LIST OF SPECIES	N	DEPTH RANGE (M)	SEDIMENT TYPE	O.M. (%)	DBUA
<i>Aglaophamus agilis</i> (Langerhans, 1880)	25	12.7-131.5	FS, CS, VCS	0.42-4.21	00062
<i>Nephtys assimilis</i> Ørsted, 1843	24	8.9-100.5	VFS, FS, CS, VCS	0.36-1.35	00060
<i>Nephtys cirrosa</i> Ehlers, 1868	18	8.9-34.9	FS, CS	0.42-0.83	00061
<i>Nephtys hombergii</i> Savigny in Lamarck, 1818	29	8.9-184.9	VFS, FS	0.76-3.06	00059
NEREIDIDAE Blainville, 1818					
<i>Eunereis longissima</i> (Johnston, 1840)	14	35.2-124.2	CS, G	0.36-0.98	00022
OENONIDAE Kinberg, 1865					
<i>Notocirrus scoticus</i> McIntosh, 1869	1	59.1	VCS	0.70	-
ONUPHIDAE Kinberg, 1865					
<i>Aponuphis bilineata</i> (Baird, 1869)	245	30.8-184.2	MS, CS, VCS, G	0.33-4.21	00043
<i>Aponuphis fauveli</i> (Rioja, 1918)	24	129.5-184.2	FS, MS	2.14-4.21	00077
<i>Diopatra neapolitana</i> Delle Chiaje, 1841	9	12.7-93.9	VFS, FS	0.74-0.99	-
<i>Hyalinoecia tubicola</i> (O.F. Müller, 1776)	1	162.8	MS	3.06	-
<i>Paradiopatra hispanica</i> (Amoureux, 1972)	19	134.9-184.2	FS, MS, G	2.14-3.06	00044
<i>Rhamphobranchium</i> (<i>Spinigerium</i>) <i>brevibrachiatum</i> (Ehlers, 1874)	1	184.2	FS	2.83	00045
OPHELIIDAE Malmgren, 1867					
<i>Ophelia neglecta</i> Schneider, 1892	5	15.6-68.9	CS	0.85	-
<i>Ophelina cylindrica</i> (Hansen, 1878)	3	129.5-157.7	FS, MS	2.53-4.21	00085
ORBINIIDAE Hartman, 1942					
<i>Orbinia latreillii</i> (Audouin and Milne Edwards, 1833)	2	12.7-13.8	FS	0.99-1.03	00091
<i>Orbinia sertulata</i> (Savigny, 1822)	3	12.7-131.5	FS	0.99-4.21	00090
OWENIIDAE Rioja, 1917					
<i>Galathowenia oculata</i> (Zachs, 1923)	49	100.5-184.2	FS, MS	1.35-4.21	00078
<i>Owenia fusiformis</i> Delle Chiaje, 1844	39	12.7-162.8	FS, MS	0.36-2.14	00004
PARAONIDAE Cerruti, 1909					
<i>Aricidea</i> (<i>Acmira</i>) <i>catherinae</i> Laubier, 1967	21	97.5-130.1	FS, MS	0.76-2.53	00582
<i>Aricidea</i> (<i>Acmira</i>) <i>cerrutii</i> Laubier, 1966	2	76.8-157.7	MS	1.29-2.57	00197
<i>Aricidea</i> (<i>Acmira</i>) <i>laubieri</i> Hartley, 1981	10	93.9-134.9	FS, MS	0.76-2.53	00581
<i>Aricidea</i> (<i>Acmira</i>) <i>simonae</i> Laubier and Ramos, 1974	2	93.9-100.5	FS	0.76-1.35	00196
<i>Aricidea</i> (<i>Allia</i>) <i>roberti</i> Hartley, 1984	5	76.5-96.3	FS	0.76-1.29	00580
<i>Aricidea</i> (<i>Aricidea</i>) <i>pseudoarticulata</i> Hobson, 1972	23	76.8- 97.5	FS, MS	0.76-3.06	00199
<i>Aricidea</i> (<i>Aricidea</i>) <i>wassi</i> Pettibone, 1965	6	93.9-184.2	FS	0.76-2.83	00195
<i>Levensenia gracilis</i> (Tauber, 1879)	1	96.3	FS	-	00583
<i>Paradoneis lyra</i> (Southern, 1914)	118	8.9-162.8	MS, CS, VCS, G	0.36-2.57	00198
PECTINARIIDAE Quatrefages, 1866					
<i>Amphictene auricoma</i> (O.F. Müller, 1776)	1	13.8	FS	1.03	-
<i>Lagis koreni</i> Malmgren, 1866	6	13.5-96.3	FS, VFS	0.74-0.90	-
PHYLLODOCIDAE Ørsted, 1843					
<i>Eteone picta</i> Quatrefages, 1866	2	13.5-15.6	FS	0.74	-
<i>Eulalia mustela</i> Pleijel, 1987	109	30.8-157.7	CS, VCS, G	0.36-2.57	00113
<i>Eumida bahusiensis</i> Bergström, 1914*	73	12.7-97.5	FS, CS	0.36-1.29	00110
<i>Eumida ockelmanni</i> Eibye-Jacobsen, 1987*	6	35.2-71.7	VCS, G	0.58-0.90	00109
<i>Hesionura elongata</i> (Southern, 1914)	189	30.8-71.7	CS, VCS, G	0.33-1.46	00111
<i>Nereiphylla paretii</i> Blainville, 1828*	4	35.2-157.7	MS, VCS	0.58-2.57	00119
<i>Paranaitis kosteriensis</i> (Malmgren, 1867)	15	35.2-157.7	MS, VCS	0.36-2.57	00118
<i>Phyllodoce lineata</i> (Claparède, 1870)	9	21.9-184.2	FS, MS	0.76-3.06	00120
<i>Phyllodoce longipes</i> Kinberg, 1866	1	12.7	VFS	0.99	-
<i>Phyllodoce rosea</i> (McIntosh, 1877)*	3	12.7-21.9	VFS	0.90-0.99	-
<i>Pseudomystides limbata</i> (Saint-Joseph, 1888)	19	30.8-162.8	MS, CS, VCS, G	0.33-3.06	00112
PILARGIDAE Saint-Joseph, 1899					
<i>Ancistrosyllis groenlandica</i> McIntosh, 1878	1	124.2	FS	-	00098
<i>Glyphohesione klatti</i> Friedrich, 1950*	1	124.4	-	-	00099
PISIONIDAE Ehlers, 1901					
<i>Pisione remota</i> (Southern, 1914)	1153	8.9-79.0 m	CS, VCS, G	0.33-1.46	00001
POECILOCHAETIDAE Hannerz, 1956					
<i>Poecilochaetus serpens</i> Allen, 1905	6	131.5	G	4.21	00387
POLYGORDIIDAE Czerniavsky, 1881					
<i>Polygordius appendiculatus</i> Fraipont, 1887	1251	30.8-79.0	CS, VCS, G	0.36-1.46	00079
POLYNOIDAE Malmgren, 1867					
<i>Harmothoe antilopes</i> McIntosh, 1876	2	96.3-124.2	FS, MS	-	00035
<i>Harmothoe glabra</i> (Malmgren, 1865)	9	30.8-129.5	FS, CS, VCS, G	0.43-2.53	00036
<i>Malmgreniella arenicolae</i> (Saint-Joseph, 1888)*	16	40.9-162.8	MS, CS, VCS	0.36-3.06	00037
<i>Malmgreniella ljunghmani</i> (Malmgreni, 1867)	167	30.8-79.0	CS, VCS, G	0.33-1.46	00038

TABLE 1. CONTINUED.

LIST OF SPECIES	N	DEPTH RANGE (M)	SEDIMENT TYPE	O.M. (%)	DBUA
SABELLIDAE Latreille, 1825					
<i>Paradialychone filicaudata</i> (Southern, 1914)	99	39.3-184.2	MS, CS, VCS, G	0.33-4.21	00663/4
SCALIBREGMATIDAE Malmgren, 1867					
<i>Scalibregma inflatum</i> Rathke, 1843	15	76.8-184.2	FS, MS	1.29-3.06	00002
SERPULIDAE Rafinesque, 1815					
<i>Ditrupa arietina</i> (O.F. Müller, 1776)	1	157.7	MS	2.57	00072
<i>Hydroides norvegicus</i> Gunnerus, 1768	22	40.9-93.9	CS, VCS, G	0.36-0.85	00073
<i>Serpula concharum</i> Langerhans, 1880	14	34.9-157.7	MS, CS, VCS	0.36-2.57	00075
<i>Spirobranchus triqueter</i> (Linnaeus, 1758)	42	35.2	VCS	0.58	00074
SIGALIONIDAE Malmgren, 1867					
<i>Sigalion squamosus</i> Delle Chiaje, 1830	1	100.5	FS	1.35	-
<i>Sthenelais limicola</i> (Ehlers, 1864)	12	76.8-131.5	FS	0.76-4.21	00030
SPIONIDAE Grube, 1850					
<i>Aonides oxycephala</i> (M. Sars, 1862)	278	30.8-184.2	MS, CS, VCS, G	0.33-3.06	00156
<i>Laonice bahusiensis</i> Söderström, 1920	44	57.3-157.7	MS, CS, VCS, G	0.70-2.57	00175
<i>Prionospio aluta</i> Maciolek, 1985*	67	59.1-184.2	FS, CS	0.70-4.21	00157
<i>Prionospio caspersi</i> Laubier, 1962	2	157.7-184.2	FS, MS	2.57-2.83	-
<i>Prionospio fallax</i> Söderström, 1920	30	68.9-134.9	FS, MS	0.76-2.53	00158
<i>Pseudopolydora paucibranchiata</i> (Okuda, 1937)*	238	97.5-184.2	FS, MS	0.98-3.06	00163
<i>Pseudopolydora pulchra</i> (Carazzi, 1895)*	2	34.9-46.1	CS	0.33-0.42	00164
<i>Scolecopsis mesnili</i> (Bellan and Lagardère, 1971)*	3	8.9-15.6	FS	0.83	-
<i>Scolecopsis cf. tridentata</i> (Southern, 1914)	5	12.7-131.5	FS, CS	0.85-4.21	-
<i>Spio decoratus</i> Bobretzky, 1870	548	8.9-131.5	FS, CS	0.33-4.21	00176
<i>Spiophanes afer</i> Meißner, 2005	12	96.3-184.2	FS	1.35-4.21	00173
<i>Spiophanes bombyx</i> (Claparède, 1870)	212	8.9-184.2	VFS, FS, MS	0.98-4.21	00174
SYLLIDAE Grube, 1850					
<i>Dioplosyllis cirrosa</i> Gidholm, 1962	2	48.2	CS	0.7	00009
<i>Eurysyllis tuberculata</i> Ehlers, 1864	6	31.8-71.7	CS, VCS	0.58-0.85	00012
<i>Exogone (Exogone) naidina</i> Ørsted, 1845	2	48.2	CS	0.7	00011
<i>Myrianida brachycephala</i> (Marenzeller, 1874)	2	31.8-71.7	VCS	0.58-0.80	00006/8
<i>Odontosyllis ctenostoma</i> Claparède, 1868	1	79.0	G	0.80	-
<i>Opisthodonta morena</i> Langerhans, 1879	26	46.4-76.8	CS, VCS	0.33-1.33	00014
<i>Palposyllis prosostoma</i> Hartmann-Schröder, 1977	35	30.8-71.7	CS, G	0.33-1.46	00029
<i>Paraehlersia ferrugina</i> (Langerhans, 1881)	48	30.8-76.8	CS, G	0.33-1.46	00013
<i>Parapionosyllis cabezali</i> Parapar, San Martín and Moreira, 2000	52	28.8-71.7	CS, G	0.44-1.46	00180
<i>Sphaerosyllis bulbosa</i> Southern, 1914	146	31.8-79.0	CS, G	0.33-1.46	00017
<i>Sphaerosyllis hystrix</i> Claparède, 1863	15	40.9-76.8	CS, VCS	0.36-1.29	00016
<i>Sphaerosyllis taylora</i> Perkins, 1981	11	31.8-71.7	VCS	0.36-1.29	00093
<i>Streptodonta pterochaeta</i> (Southern, 1914)	35	40.9-76.8	CS, G	0.33-1.46	00015
<i>Syllis garciai</i> (Campoy, 1982)	231	30.8-93.9	CS, VCS, G	0.33-1.46	00028
<i>Syllis licheri</i> Ravara, San Martín and Moreira in San Martín, 2003	31	46.1-100.5	MS, G	0.33-1.35	00589
<i>Syllis parapari</i> San Martín and López, 2000	20	31.8-100.5	CS, VCS	0.36-1.35	00429
<i>Syllis pontxioi</i> San Martín and López, 2000	213	30.8-79.0	CS, G	0.33-1.46	00019
<i>Trypanosyllis coeliaca</i> Claparède, 1868	39	30.8-76.8	CS, G	0.39-1.46	00020
TEREBELLIDAE Grube, 1850					
<i>Janice conchilega</i> (Pallas, 1766)	16	12.7-162.8	FS, MS, CS, VCS	0.33-3.06	-
<i>Loimia medusa</i> (Savigny, 1822)	1	59.1	VCS	0.70	00153
<i>Pista malmgreni</i> Saphronova and Jirkov in Jirkov, Leontovich and Saphronova, 2001	36	35.2-79.0	CS, VCS, G	0.33-1.46	00154
<i>Polycirrus cf. medusa</i> Grube, 1850*	211	39.4-184.2	MS, CS, VCS, G	0.33-3.06	-
TRICHOBRANCHIDAE Malmgren, 1866					
<i>Terebellides stroemii</i> Sars, 1835	2	124.2-184.2	FS	2.83	00155

TABLE 2. Mean values of species richness (S), number of individuals (N), diversity (H'), equitability (J'), organic matter content (O.M.) and sediment type (VFS, very fine sand; FS, fine sand; MS, median sand; CS, coarse sand; VCS, very coarse sand; G, gravel) for each group of sites resultant from classification analysis.

	S	N	H'	J'	O.M. (%)	Sed. type
Inner-shelf	14.7	57.8	2.69	0.703	0.898	VFS, FS
Mid-shelf	39.4	193.8	3.89	0.736	0.730	CS, VCS, G
Outer-shelf	28.9	41.2	4.03	0.828	2.270	FS, MS

ACKNOWLEDGMENTS: Thanks are due to the chief scientist of the research cruises, Dr. Jean Claude Sorbe, for the opportunity of carrying out this work, and to the crew of RV Côte d'Aquitaine and all the participants in the cruises, for the fieldwork. The authors also wish to thank Dr.

Guillermo San Martín, Dr. Eduardo López, Dr. José Manuel Viéitez, Dr. Julio Parapar, Dr. Florencio Aguirrezabalaga, Dr. Alex Muir, Dr. Gordon Patersen and Dr. Fredrik Pleijel for the assistance on taxonomic identification of the species. This research was partially supported by the Junta Nacional

de Investigação Científica e Tecnológica (JNICT) with two grants (PRAXIS XXI/BM/6595/95 and PRAXIS XXI/BTI/16724/98) and by the CZCM research unit from Universidade de Aveiro.

LITERATURE CITED

- Abrantes, I., F. Magalhães and J.M.A. Dias. 1994. Characterization of the surface sediments of the continental shelf and upper slope between Espinho and Aveiro. *Gaia* 8: 97-104.
- Almaça, C. 1960. Contribuição para o estudo da zonação marinha do litoral português. *Boletim da Sociedade Portuguesa de Ciências Naturais* 23: 201-208.
- Amanieu, M., O. Guelorget and P. Michel. 1977. Richesse et diversité de la macrofaune benthique d'une lagune littorale méditerranéenne. *Vie et Milieu* 27(1B): 85-109.
- Amanieu, M., O. Guelorget and J. Nouguié-Soule. 1981. Analyse de la diversité de la macrofaune benthique d'une lagune littorale méditerranéenne. *Vie et Milieu* 31(3-4): 303-312.
- Amoureux, L. 1974. Annélides polychètes recueillies sur les pentes du talus continental au nord-ouest de l'Espagne et du Portugal (Campagne 1972 de la "Thalassa"). *Cuadernos de Ciencias Biológicas* 3: 121-154.
- Amoureux, L. 1985. Annélides polychètes d'un transect du plateau continental Sud-Gascogne. *Cahiers de Biologie Marine* 26: 341-354.
- Amoureux, L. 1987. Annélides polychètes du talus continental européen. Données complémentaires concernant les campagnes 1970-1971-1972 de la "Thalassa". Données globales sur l'ensemble des six campagnes (1967 à 1973). *Cahiers de Biologie Marine* 28: 569-577.
- Amoureux, L. and J. Calvário. 1981. Annélides Polychètes du Portugal. Données nouvelles. *Arquivos do Museu Bocage, série B* 1: 145-156.
- Ballesteros, E. 2006. Mediterranean coralligenous assemblages: a synthesis of present knowledge. *Oceanography and Marine Biology Annual Review* 44: 123-195.
- Bellan, G. 1960. Annélides polychètes. *Résultats Scientifiques de la Campagne du N.R.P. "Faial" dans les eaux côtières du Portugal (1957)*. Lisboa 2: 1-31.
- Bellan, G. 1962. Ille Campagne Méditerranéenne du N/O "Président Théodore Tissier" de l'Institut Scientifique et Technique des Pêches. Annélides Polychètes. *Recueil des Travaux de la Station Marine d'Endoume* 25(39): 23-37.
- Bellan, G. 1964. Contribution à l'étude systématique, bionomique et écologique des Annélides Polychètes de la Méditerranée. *Recueil des Travaux de la Station Marine d'Endoume* 49(33): 1-371.
- Bellan, G. 1965. Contributions à l'étude bionomique de la Méditerranée Occidentale (Côte du Var et des Alpes maritimes - côte occidentale de Corse). *Bulletin de l'Institut Océanographique de Monaco* 65(1345): 1-24.
- Buchanan, B.A. 1963. The bottom fauna communities and their sediment relationships off the coast of Northumberland. *Oikos* 14(2): 154-175.
- Buchanan, J.C. and J.M. Kain. 1971. Measurement of the physical and chemical environment; p. 30-58 In N.A. Holme and A.D. McIntyre (ed.). *Methods for the study of marine benthos*. Oxford: Blackwell.
- Field, J.G., K.R. Clarke and R.M. Warwick. 1982. A practical strategy for analysing multispecies distribution patterns. *Marine Ecology Progress Series* 8: 37-52.
- Cebrián, E. and E. Ballesteros. 2004. Zonation patterns of benthic communities in La Herradura (Alboran Sea, western Mediterranean). *Scientia Marina* 68: 69-84.
- Çinar, M.E. 2005. Polychaetes from the coast of northern Cyprus (eastern Mediterranean Sea), with two new records for the Mediterranean Sea. *Cahiers de Biologie Marine* 46: 143-159.
- Dexter, D.M. 1988. The sandy beach fauna of Portugal. *Arquivos do Museu Bocage, nova série* 1: 101-110.
- Dexter, D.M. 1990. The effect of exposure and seasonality on sandy beach community structure in Portugal. *Ciência Biológica - Ecology Systematics* 10: 31-50.
- Dyer, M.F., W.G. Fry, P.D. Fry and G.J. Cranmer. 1983. Benthic regions within the North Sea. *Journal of the Marine Biology Association of the United Kingdom* 63: 683-693.
- Eleftheriou, A. and D.J. Basford. 1989. The macrobenthic infauna of the offshore northern North Sea. *Journal of the Marine Biology Association of the United Kingdom* 69: 123-143.
- França, S., C. Vinagre, M.A. Pardal and H.N. Cabral. 2009. Spatial and temporal patterns of benthic invertebrates in the Tagus estuary, Portugal: comparison between subtidal and intertidal mudflat. *Scientia Marina* 73(2): 307-318.
- Gil, J. and R. Sardá. 1999. New records of Annelida Polychaeta for the Portuguese fauna (with comments on some already known species). *Arquivos do Museu Bocage, nova série* 3(10): 287-336.
- Guerra-García, J.M., M.P. Cabezas, E. Baeza-Rojano, D. Izquierdo, J.R. Corzo, M. Ros, J.A. Sánchez, A. Dugo-Cota, A.M. Flores-Léon and M.M. Soler-Hurtado. 2011. Abundance patterns of macrofauna associated to marine macroalgae along the Iberian Peninsula. *Zoologica baetica* 22: 3-17.
- Guille, A. 1970. Bionomie benthique du plateau continental de la côte catalane française. II. Les communautés de la macrofaune. *Vie et Milieu* 21: 149-280.
- Hartmann-Schröder, G. 1977. Polychaeten aus dem Sublitoral und Bathyal vor der portugiesischen und marokkanischen Küste. Auswertung der Fahrt 8 (1967) von F. S. "Meteor". *"Meteor" Forschungsergebnisse D* 26: 65-99.
- Hartmann-Schröder, G. 1979. Die Polychaeten der "Atlantischen Kuppenfahrt" von F. S. "Meteor" (Fahrt 9 c, 1967). 1. Proben aus Schleppgeräten. *"Meteor" Forschungsergebnisse D* 31: 63-64.
- Hartmann-Schröder, G. 1981. Die Polychaeten der Fahrten 11, 19, 21 und 26 (1967-1972) von F. S. "Meteor" in das Gebiet zwischen dem Golf von Biscaya und dem Auftriebsgebiet vor Westafrika. *"Meteor" Forschungsergebnisse* 33: 23-36.
- Heip, C. and J.A. Craeymeersch. 1995. Benthic community structures in the North Sea. *Helgoländer Meeresuntersuchungen* 49: 313-328.
- Hoey, G.V., S. Degraer and M. Vincx. 2004. Macrobenthic community structure of soft-bottom sediments at the Belgian continental shelf. *Estuarine Coastal and Shelf Science* 59: 599-613.
- Künitzer, A., D. Basford, J.A. Craeymeersch, J.M. Dewarumez, J. Dörjes, G.C.A. Duineveld, A. Eleftheriou, C. Heip, P. Herman, O.P. Kingston, U. Niermann, E. Rachor, H. Rumohr and P.A.J. de Wilde. 1992. The benthic infauna of the North Sea: species distribution and assemblages. *ICES Journal of Marine Science* 49: 127-143.
- Mackie, A.S.Y., C. Parmiter and L.K.Y. Tong. 1997. Distribution and diversity of Polychaeta in the Southern Irish Sea. *Bulletin of Marine Science* 60: 467-481.
- Martin, D., S. Pinedo and R. Sardá. 2000. Distribution patterns and trophic structure of soft-bottom polychaete assemblages in a north-western Mediterranean shallow-water bay. *Ophelia* 53(1): 1-17.
- Martins, V., I. Abrantes, C. Granjeira, P. Martins, R. Nagai, S.H.M. Sousa, L.L.M. Laut, J.M.A. Dias, J.M. Dias, E.F. Silva and F. Rocha. 2012. Records of sedimentary dynamics in the continental shelf and upper slope between Aveiro-Espinho (N Portugal). *Journal of Marine Systems* 96-97: 48-60.
- Martínez J. and I. Adarraga. 2001. Distribución batimétrica de comunidades macrobentónicas de sustrato blando en la plataforma continental de Guipúzcoa (Golfo de Vizcaya). *Boletín Instituto Español de Oceanografía* 17: 33-48.
- Mohamed, M. 1993. Étude faunistique et écologique des peuplements infralittoraux superficiels des côtes rocheuses du Maroc. I - Peuplement à *Padina pavonica*. *Arquivos do Museu Bocage, nova série* 2(14): 287-300.
- Monteiro-Marques, V. 1979. Contribution à l'étude du plateau continental des côtes portugaises (Algarve, Sud Portugal): étude des peuplements de substrats meubles. *Téthys* 9(2): 191-195.
- Monteiro-Marques, V. 1987. A plataforma continental do Algarve. Definição qualitativa das biocenoses de substrato móvel. *Documentos Técnicos do Instituto Hidrográfico, Lisboa* 31: 1-204.
- Moreira, M.H., H. Queiroga, M.M. Machado and M.R. Cunha. 1993. Environmental gradients in a southern Europe estuarine system: Ria de Aveiro, Portugal. Implications for soft bottom macrofauna colonization. *Netherlands Journal of Aquatic Ecology* 27(2-4): 465-482.
- Parapar, J. and J. Moreira. 2009. Polychaeta of the "DIVA-Artabria I" project (cruise 2002) in the continental shelf and upper slope off Galicia (NW Spain). *Cahiers de Biologie Marine* 50: 57-78.
- Pérès, J.M. 1959. Aperçu bionomique sur les communautés benthiques des côtes sud du Portugal. *Résultats Scientifiques de la campagne du N.R.P. "Faial" dans les eaux côtières du Portugal (1957)*. Lisboa 1: 1-35.
- Picard, J. 1965. Recherches qualitatives sur les biocenoses marines des substrats meubles dragables de la région marseillaise. *Recueil des Travaux de la Station Marine d'Endoume* 52(36): 1-60.
- Sardá, R., S. Rossi, X. Martí and J.M. Gili. 2012. Marine benthic cartography of the Cap de Creus (NE Catalan Coast, Mediterranean Sea). *Scientia Marina* 76(1): 159-171.
- Snelgrove, P.V.R. 1998. The biodiversity of macrofaunal organisms in marine sediments. *Biodiversity and Conservation* 7: 1123-1132.
- Somaschini, A., N. Martini, M.F. Gravina, A. Belluscio, F. Corsi and G.D. Aridizzone. 1998. Characterization and cartography of some Mediterranean soft-bottom benthic communities (Ligurian Sea, Italy). *Scientia Marina* 62:27-36.
- Sousa-Reis, C., V.M. Monteiro-Marques, J. Calvário, J.C. Marques, R. Melo and R. Santos. 1982. Contribuição para o estudo dos povoamentos bentónicos (substrato móvel) da costa ocidental portuguesa. *Oecologia Aquatica* 6: 91-105.

RECEIVED: November 2012

ACCEPTED: May 2013

PUBLISHED ONLINE: June 2013

EDITORIAL RESPONSIBILITY: Luis E. Arruda Bezerra