

Strobilurus torquatus Wiegmann, 1834 (Squamata: Tropiduridae): New records from the Brazilian state of Paraíba and a geographic distribution map

Keoma Coutinho Rodrigues^{1*}, Fagner Ribeiro Delfim², Carla Soraia Soares de Castro¹, Frederico Gustavo Rodrigues França¹, Edinaldo Leite Filho², Daniel Oliveira Mesquita², Felipe Araújo de Oliveira², Alinny Costa Araújo dos Santos³, Stephen Francis Ferrari³, Mônica Mafra Valençá-Montenegro⁴

¹ Universidade Federal da Paraíba, Centro de Ciências Aplicadas e Educação, Pós-Graduação em Ecologia e Monitoramento Ambiental, Departamento de Engenharia e Meio Ambiente. Rua da Mangueira. CEP 58297-000. Rio Tinto, PB, Brazil.

² Universidade Federal da Paraíba, Centro de Ciências Exatas e da Natureza, Departamento de Sistemática e Ecologia. Cidade Universitária. CEP 58051-900. João Pessoa, PB, Brazil.

³ Universidade Federal de Sergipe. Departamento de Biologia. Avenida Marechal Rondon, Rosa Elze. CEP 49000-100. São Cristóvão, SE, Brazil.

⁴ Centro Nacional de Pesquisa e Conservação de Primatas Brasileiros, Instituto Chico Mendes de Conservação da Biodiversidade. Praça Antenor Navarro, 5, Varadouro. CEP 58010-480. João Pessoa, PB, Brazil.

* Corresponding author. E-mail: keoma.rodrigues09@gmail.com

ABSTRACT: We provide the first records of *Strobilurus torquatus* from the Brazilian state of Paraíba and review its distribution based on specimens available in Brazilian collections and published data. This result emphasizes the need for more surveys on reptiles at this region.

The Tropiduridae is one of the most species-rich lizard families at the Neotropical region (Torres-Carvajal 2004). The family comprises 117 species distributed in eight genera – *Eurolophosaurus*, *Microlophus*, *Plica*, *Stenocercus*, *Strobilurus*, *Tropidurus*, *Uracentron*, and *Uranoscodon* (Uetz 2012). Tropidurids are widespread in Brazil, where a total of 35 species is found (Bérnails and Costa 2011), with representatives of the family having been recorded in all terrestrial biomes (Rodrigues 1987, Martinez *et al.* 2011).

The systematics of Tropiduridae went through a number of reviews in recent years (Jackson 1978, Rodrigues 1987, Frost 1992, Frost *et al.* 2001). Based on a cladistic analysis, Frost (1992) considered *Plica*, *Uracentron*, and *Strobilurus* to be junior synonyms of *Tropidurus*. Following this classification, *Strobilurus torquatus* Wiegmann, 1834 was renamed *Tropidurus strobilurus*, given that the binomial *Tropidurus torquatus* Wied, 1820 had already been assigned. More recently, Frost *et al.* (2001) presented molecular and morphological evidence that *Strobilurus* should be considered a distinct and valid genus, revalidating it.

Strobilurus torquatus (Figure 1A) differs from the other tropidurid species found in Northeastern Brazil due to the presence of a relatively shorter spiny tail with enlarged, heavily keeled and mucronated scales. It is considered a forest-dwelling heliophilic lizard, which feeds mainly on ants found on tree trunks at different levels in the forest, including the ground (Jackson 1978, Rodrigues *et al.* 1989). Its currently known distribution lies within the Brazilian Atlantic Forest, ranging from Rio de Janeiro state (Salles and Silveira 2010) at its southern end, and Pernambuco in the north (Rodrigues *et al.* 1989), and

isolated rainforest enclaves within the semi-arid caatinga scrublands of Ceará, further west (Jackson 1978, Borges-Nojosa and Caramaschi 2003).

This study presents the first records of *Strobilurus torquatus* (Figure 2) from the Brazilian state of Paraíba, and reviews its geographic distribution based on the available published data, and including specimens deposited at herpetological collections of the following Brazilian institutions (see Table 1): Museu de Zoologia da Universidade de São Paulo (MZUSP) in São Paulo; Museu Nacional, Universidade Federal do Rio de Janeiro (MNRJ) in Rio de Janeiro; Universidade de Brasília (CHUNB); Universidade de Sergipe (CHUFS) in Aracaju; and Museu de História Natural da Universidade de Alagoas (MUFAL) in Maceió. A non-examined specimen from Espírito Santo state, whose record is available on Species Link (2010) was also included. This specimen is deposited at Museu de Biologia Professor Mello Leitão (MBML) in Santa Teresa, Espírito Santo.

The collections were authorized by the federal environment agency, through SISBIO/ICMBio permit number 33909-1. The specimens were deposited in the herpetological collection of the Universidade Federal da Paraíba, under catalog numbers CHUFPB 9163 and 10761.

One individual (Figure 1B) was collected at 15:02 h on February 23rd, 2012, in the Engenho Gargaú Private Natural Heritage Reserve (RPPN Engenho Gargaú), a 1058.62 ha remnant of Atlantic Forest inserted within a matrix of sugarcane plantations in the municipality of Santa Rita in Paraíba state, Brazil (6°59'52" S, 34°57'30" W). The area is covered with a well-preserved forest, canopy reaching up to 12-15 m in height (Figure 3A). The lizard had been partially ingested by a female blonde

capuchin monkey, *Sapajus flavius* Schreber 1774, which discarded its prey when disturbed. It was not possible to determine the sex of this individual.

A second specimen, a male, (Figure 1C) was collected on April 23, at Mata do Buraquinho, part of the Benjamin Maranhão Botanical Garden (JBBM), located in the municipality of João Pessoa, Paraíba ($7^{\circ}08'42''S$, $34^{\circ}51'54''W$). The lizard was captured in glue traps installed in the tabular roots of a fig tree (*Ficus doliaria* Mart.) at the edge of a trail. Another individual was observed in the same tree at the same day, but not captured. The study area has

irregularly continuous canopy of 15-25 m in height, with a well-developed shrubby understory (Figure 3B). This site has 515 ha of forest, and is located in the southwestern portion of João Pessoa municipality, and is one of the few Atlantic Forest remnants in northeastern Brazil inserted entirely within an urban matrix.

The present records of *Strobilurus torquatus* from Paraíba help fill the lacuna in its distribution at the northeastern of Brazil (Figure 3). The state of Paraíba have suffered intense deforestation for the cultivation of crops such as sugarcane (Campanili and Schaffer 2010) that may




FIGURE 1. A: *Strobilurus torquatus* observed and not collected at RPPN Engenho Gargaú, Santa Rita, Paraíba (Brazil). Photograph by Caio Brito. B: *Strobilurus torquatus* (CHUFPB 9163) collected at RPPN Engenho Gargaú, Santa Rita, Paraíba (Brazil). The specimen was discarded by a female blonde capuchin monkey, *Sapajus flavius*, when disturbed. Photograph by Keoma Rodrigues. C: *Strobilurus torquatus* specimen (CHUFPB 10761) from Mata do Buraquinho, Benjamin Maranhão Botanical Garden, João Pessoa, Paraíba (Brazil). Photograph by Fagner Delfim.




FIGURE 2. Known geographic distribution of *Strobilurus torquatus*, including the new records reported here. Black Circles = collection specimens; White Circles = published records; Red Asterisks = new records from Paraíba. Legend: Alagoas state: 1 – Campo Alegre; 2 – Coruripe; 3 – Flexeiras; 4 – Maceió; Ipioca; 5 – Murici; 6 – Passo de Camaragibe; 7 – Quebrangulo; 8 – Rio Largo; 9 – São Miguel dos Campos; Bahia state: 10 – Ilhéus; 11 – Itabuna; 12 – Mata de São João; 13 – Porto Seguro; 14 – São José da Vitoria; 15 – Una; Ceará state: 16 – Baturité; 17 – Ibiapaba; 18 – Maranguape; 19 – Pacatuba; Espírito Santo state: 20 – Aracruz, Santa Cruz; 21 – Linhares; 22 – Presidente Kennedy; 23 – Santa Teresa; Minas Gerais state: 24 – UHE Barra do Braúná; Pernambuco state: 25 – Itamaracá; 26 – São Caetano; 27 – Timbaúba; 28 – Vicência, Água Azul; Rio de Janeiro state: 29 – Pedra do Garrafão; Sergipe state: 30 – Serra de Itabaiana; Paraíba state: 31 – João Pessoa; 32 – Santa Rita.

have accentuated the difficulty in recording forest dwellers reptilian species, as *S. torquatus*.

One of the locations, where the species was recorded has been sampled intensively for at least three years (Santana *et al.* 2007 and later by Pereira-Filho, unpublished data). Even with such efforts the species has not been

found, what may indicate a low abundance or deficiency in sampling methods. We believe that our findings demonstrate important records for the distribution of *S. torquatus*, and enhance the importance of more and better sampling efforts at the Atlantic Forest areas in the Northeast of Brazil.



FIGURE 3. Habitat on which the specimens of *Strobilurus torquatus* were collected at the Brazilian state of Paraíba : A - RPPN Engenho Gargaú, Santa Rita ($6^{\circ}59'52''$ S, $34^{\circ}57'30''$ W). Photograph by Keoma Rodrigues; B – Mata do Buraquinho, Benjamin Maranhão Botanical Garden, João Pessoa ($7^{\circ}08'48.31''$ S, $34^{\circ}51'23.20''$ W). Photograph by Suênia Oliveira.

TABLE 1. Records of *Strobilurus torquatus* based on museum specimens and literature data. Acronyms: CHUNB –Herpetological Collection of the Universidade de Brasília; MZUSP – Museu de Zoologia da Universidade de São Paulo; MUFAL – Museu de História Natural da Universidade Federal de Alagoas ; CHUFS – Herpetological Collection of the Universidade Federal de Sergipe; MNRJ – Museu Nacional do Rio de Janeiro, Rio de Janeiro; MBML – Museu de Biologia Professor Mello Leitão (Species Link data). * New records.

LOCALITY	STATE	LATITUDE	LONGITUDE	REFERENCE
Campo Alegre	Alagoas	$9^{\circ}46'46.27''$ S	$36^{\circ}21'01.20''$ W	MUFAL 6720
Coruripe	Alagoas	$10^{\circ}08'00.62''$ S	$36^{\circ}10'33.89''$ W	MUFAL 6227
Flexeiras	Alagoas	$9^{\circ}16'21.54''$ S	$35^{\circ}43'25.92''$ W	MUFAL 1918
Maceió, Ipioca	Alagoas	$9^{\circ}31'27.52''$ S	$35^{\circ}35'45.15''$ W	MUFAL 5410
Murici	Alagoas	$9^{\circ}18'17.64''$ S	$35^{\circ}56'30.24''$ W	MUFAL 1781
Passo de Camaragibe	Alagoas	$9^{\circ}14'55.18''$ S	$35^{\circ}29'24.41''$ W	MNRJ 3404
Quebrangulo	Alagoas	$9^{\circ}19'24.14''$ S	$36^{\circ}28'27.85''$ W	MNRJ 3906
Rio Largo	Alagoas	$9^{\circ}29'48.71''$ S	$35^{\circ}51'28.76''$ W	MUFAL 441
São Miguel dos Campos	Alagoas	$9^{\circ}46'48.68''$ S	$36^{\circ}05'54.05''$ W	Jackson 1978
Ilhéus	Bahia	$14^{\circ}47'49.84''$ S	$39^{\circ}02'03.84''$ W	MNRJ 3441
Itabuna	Bahia	$14^{\circ}47'21.40''$ S	$39^{\circ}16'40.09''$ W	MZUSP 29626
Mata de São João	Bahia	$12^{\circ}27'43.00''$ S	$38^{\circ}14'01.00''$ W	Fonseca <i>et al.</i> 2012
Porto Seguro	Bahia	$16^{\circ}27'04.04''$ S	$39^{\circ}03'52.79''$ W	MZUSP 95220
São José da Vitória	Bahia	$15^{\circ}05'00.00''$ S	$39^{\circ}21'00.00''$ W	Rodrigues <i>et al.</i> 1989
Una	Bahia	$15^{\circ}16'10.43''$ S	$39^{\circ}04'09.92''$ W	MZUSP 91625
Baturité	Ceará	$4^{\circ}19'40.56''$ S	$38^{\circ}53'04.55''$ W	Borges-Nojosa and Caramaschi 2003
Ibiapaba	Ceará	$5^{\circ}03'00.00''$ S	$40^{\circ}55'00.00''$ W	Borges-Nojosa and Caramaschi 2003
Maranguape	Ceará	$3^{\circ}53'26.39''$ S	$38^{\circ}40'54.74''$ W	Borges-Nojosa and Caramaschi 2003
Pacatuba	Ceará	$3^{\circ}58'57.63''$ S	$38^{\circ}38'13.28''$ W	Borges-Nojosa and Caramaschi 2003
Aracruz, Santa Cruz	Espírito Santo	$19^{\circ}56'02.61''$ S	$40^{\circ}08'18.89''$ W	MZUSP 36715
Linhares	Espírito Santo	$19^{\circ}23'21.42''$ S	$40^{\circ}03'52.25''$ W	MZUSP 30747
Presidente Kennedy	Espírito Santo	$21^{\circ}05'16.95''$ S	$41^{\circ}01'55.45''$ W	CHUNB 24129
Santa Teresa	Espírito Santo	$19^{\circ}56'08.16''$ S	$40^{\circ}36'01.08''$ W	MBML 281
UHE Barra do Braúná	Minas Gerais	$21^{\circ}26'42.36''$ S	$42^{\circ}25'43.68''$ W	Costa 2011
João Pessoa	Paraíba	$7^{\circ}08'48.31''$ S	$34^{\circ}51'23.20''$ W	CHUFPB*
Santa Rita	Paraíba	$6^{\circ}59'52.30''$ S	$34^{\circ}57'30.65''$ W	CHUFPB*
Itamaracá	Pernambuco	$7^{\circ}44'58.57''$ S	$34^{\circ}49'39.72''$ W	MZUSP 21481
São Caetano	Pernambuco	$8^{\circ}22'09.00''$ S	$36^{\circ}04'60.00''$ W	MZUSP 66131
Timbaúba	Pernambuco	$7^{\circ}30'11.17''$ S	$35^{\circ}18'50.46''$ W	MNRJ 12940
Vicência, Água Azul	Pernambuco	$7^{\circ}40'06.32''$ S	$35^{\circ}19'38.84''$ W	MZUSP 23072
Pedra do Garrafão	Rio de Janeiro	$21^{\circ}12'18.00''$ W	$41^{\circ}20'27.60''$ W	Salles and Silveira 2010
Serra de Itabaiana	Sergipe	$10^{\circ}44'18.91''$ S	$37^{\circ}23'03.22''$ W	CHUFS 363

ACKNOWLEDGMENTS: We thank the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) and the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for providing scholarships, and personnel at Japungu Agroindustrial S.A. and the Jardim Botânico Benjamin Maranhão for logistic support and permission to collect data within their properties. Caio Brito and Carine Fiúza provided the images.

LITERATURE CITED

- Bérnilds, R.S. and H.C. Costa. 2011. *Brazilian reptiles – List of species. Electronic*. Database accessible at <http://www.sbherpetologia.org.br/>. Captured on 29 February 2012.
- Borges-Nojosa, D.M. and U. Caramaschi. 2003. Composição e Análise Comparativa da Diversidade e das Afinidades Biogeográficas dos Lagartos e Anfíbiosbenídeos (Squamata) dos Brejos Nordestinos; p.489-540. In I. Leal, J.M.C. Silva and M. Tabarelli (ed.). Ecologia e Conservação da Caatinga. Recife: UFPE.
- Campanili, M. and Schaffer, W.B. 2010. *Mata Atlântica: patrimônio nacional dos brasileiros*. Brasília: Ministério do Meio Ambiente. 408 p.
- Costa, H.C. 2011. *Strobilurus torquatus*: Geographic Distribution. *Herpetological Review* 42(4): 570.
- Fonseca, E., M. S. Tinôco, R. Marques, H. C. Browne-Ribeiro and C. P. Fazolato. 2012. *Strobilurus torquatus* (Amazon Lava Lizard): Geographic Distribution. *Herpetological Review* 43(3): 446.
- Frost, D.R. 1992. Phylogenetic analysis and taxonomy of the *Tropidurus* group of lizards (Iguania: Tropiduridae). *American Museum Novitates* (3033): 1-68.
- Frost, D.R., M.T. Rodrigues, T, and T.A. Titus. 2001. Phylogenetics of the Lizard Genus *Tropidurus* (Squamata: Tropiduridae: Tropidurinae): Direct Optimization, Descriptive Efficiency, and Sensitivity Analysis of Congruence Between Molecular Data and Morphology. *Molecular Phylogenetics and Evolution* 21 (3): 352-371.
- Jackson, J. F. 1978. Differentiation in the Genera *Enyalius* and *Strobilurus* (Iguanidae): Implications for Pleistocene Climatic Changes in Eastern Brazil. *Arquivos de Zoologia* 30(1): 1-80.
- Martinez, L.E., L.J. Avila, C.H.F. Perez, D.R. Perez, J.W. Sites, Jr. and M. Morando. 2011. A new species of *Liolaemus* (Squamata, Iguanidae, Liolaemini) endemic to the Auca Mahuida volcano, northwestern Patagonia, Argentina. *Zootaxa* 3010: 31 – 46.
- Rodrigues, M.T., Y. Yonenaga-Yassuda and S. Kasahara. 1989. Notes on the ecology and karyotypic description of *Strobilurus torquatus* (Sauria, Iguanidae). *Revista Brasileira de Genética* 12 (4): 747-759.
- Salles, R. O. Land A. L. Silveira. 2010. Primeiro Registro de *Strobilurus torquatus* Wiegmann, 1834 (Reptilia, Squamata, Tropiduridae) no estado do Rio de Janeiro, Brasil. *Evolução e Conservação da Biodiversidade* 1 (1): 13.
- Santana, G.G., W.L.S. Vieira, G.A. Pereira-Filho, F.R. Delfim, Y.C.C. Lima and K.S. Vieira. 2008. Herpetofauna em um fragmento de Floresta Atlântica no Estado da Paraíba, Região Nordeste do Brasil. *Biotaem* 21(1): 75-84.
- Species Link 2010. *Species link*. Electronic Database accessible at <http://www.splink.cria.org.br/>. Captured on 29 February 2012.
- Torres-Carvajal, O. 2004. The abdominal skeleton of tropidurid lizards (Squamata: Tropiduridae). *Herpetologica* 60 (1): 75-83.
- Uetz, P.(editor) 2012. *The Reptile Database*. Electronic database accessible at <http://reptile-database.org>, Captured on 29 February 2012.

RECEIVED: November 2012

ACCEPTED: Feberuary 2013

PUBLISHED ONLINE: June 2013

EDITORIAL RESPONSIBILITY: Pedro M. S. Nunes