

Occurrence of *Phyllostomus elongatus* (Geoffroy St.-Hilaire, 1810) (Chiroptera, Phyllostomidae) in the Cerrado of Tocantins and a compilation of its Brazilian distribution

Roberto Leonan Morim Novaes^{1*}, Renan de França Souza², Saulo Felix¹, Gabriella Jacob¹, Cristal Sauwen¹ and Leonardo dos Santos Avilla¹

¹ Universidade Federal do Estado do Rio de Janeiro, Departamento de Zoologia. Av. Pasteur, 458, Sala 501, Urca, CEP 22290-240, Rio de Janeiro, RJ, Brazil.

² Universidade do Estado do Rio de Janeiro, Programa de Pós-Graduação em Ecologia e Evolução. Av. São Francisco Xavier 524, Maracanã, CEP 20550-013, Rio de Janeiro, RJ, Brazil.

* Corresponding author. E-mail: robertoleonan@gmail.com

ABSTRACT: The lesser spear-nosed bat, *Phyllostomus elongatus*, is endemic of South America and in Brazil this species is recorded in Amazon, Atlantic Forest, Cerrado, and Pantanal. Here, we present a new record for *P. elongatus* in the savanna of central Brazil, known as the Cerrado. In January 2012, five individuals of *P. elongatus* were captured and recorded in a limestone cave in the Aurora do Tocantins county, Tocantins State. *Phyllostomus elongatus* has already been registered in 32 locations and 14 Brazilian states, and their records appear to be associated with humid habitats and forested areas.

The genus *Phyllostomus* Lacépède, 1799, is represented by four species, all of which occur in Brazil (Williams and Genoways 2008): *Phyllostomus discolor* Wagner, 1843, *Phyllostomus elongatus* (Geoffroy St.-Hilaire, 1810), *Phyllostomus hastatus* (Pallas, 1767), and *Phyllostomus latifolius* (Thomas, 1901). *Phyllostomus elongatus* is endemic of South America and distributed from Venezuela, Bolivia, Peru, Ecuador, Colombia, and the Guyanas to the southeast of Brazil (Baker *et al.* 1988; Williams and Genoways 2008). *P. elongatus* occurs east of the Andes, in lowlands, and its distribution is associated with tropical rainforests (Williams and Genoways 2008). Furthermore, an isolated population of this species is known to the west of the Andes, occurring in Colombia and Ecuador (Williams and Genoways 2008). This species occupies primary and secondary forests, disturbed areas, and karst environments (Fischer *et al.* 1997, Simmons and Voss 1998, Fregonezi *et al.* 2013). *P. elongatus* can form harems of more than 10 individuals or colonies composed of up to 50 non-reproductive males (Fregonezi *et al.* 2013). *P. elongatus* can be considered omnivorous like the other species of the genus, since their diet includes insects, nectar, fruits and small vertebrates (Tuttle 1970, Reis and Peracchi 1987, Fischer *et al.* 1997, Bernard 2002).

In Brazil, *P. elongatus* has been registered in four biomes: the Amazon (Handley 1967; Piccinini 1974; Marques 1985; Bernard *et al.* 2011a), the Atlantic Forest (Vieira 1942; Faria *et al.* 2006; Faria and Baumgarten 2007), the Pantanal (Bordignon and França 2009; Alho *et al.* 2011), and the Cerrado (Sousa *et al.* 2013). The first record assigned to the Brazilian Cerrado, the type-locality for the species (Rio Branco, Mato Grosso), is, in fact, a region that combines a set of different landscapes, characterized by presenting transitional vegetation between the Pantanal and the Cerrado (Eva *et al.* 2002). Records of this species in the Caatinga biome were mentioned by

Marinho-Filho and Sazima (1998), however, this data was actually from an Atlantic Forest region in the northeastern Brazil (Vieira 1953, 1955; Souza-Lopes 1978; Souza *et al.* 2004; Nogueira *et al.* 2007). Thus, here we present the first record of *P. elongatus* for the central Cerrado and also for the Tocantins State, Brazil.

We recorded *P. elongatus* during a bat survey in a karstic area of the Aurora do Tocantins municipality, Southeast Tocantins State, Northern Brazil. The site has several karstic limestone complexes, each with dozens of caves. This karstic complex is inserted into a mosaic landscape composed of large remnants of the Neotropical savanna and farms with cattle ranching and maize, cassava, rice, and soybeans plantations. Aurora do Tocantins presents typical vegetation of the Cerrado *sensu strictu*, with enclaves of gallery forest (known as Cerradão, in Portuguese) in areas near the limestone massifs.

We conducted eight nights of bat surveys, with five nights consisting of twelve hour surveys (18:00–06:00 h) and three nights of six hour surveys (18:00–00:00h) using ten mist-nets (Zootech® 9 x 3 m, mesh: 20 mm), totaling 21,260 m².h of sampling effort (Straube and Bianconi 2002). The mist-nets were placed (1) at cave entrances (6,390 m².h); (2) from a distance of 15 meters to cave entrances (6,380 m².h); (3) in edges, trails, and glades of Cerrado fragments (6,380 m².h); and, (4) around artificial ponds (2,110 m².h). Captured individuals were marked with a perforation-code in the dactylopatagium (Bonaccorso and Smythe 1972) and they were released afterwards at the same location of their capture sites. Some individuals were collected and deposited in the mammalian collection of the Museu Nacional do Rio de Janeiro (MN78389, MN78390, MN78385), which was authorized by the SISBIO/IBAMA (4028-1/28717). We followed the guidelines of Sikes *et al.* (2011) for handling the animals. Individuals were identified using the taxonomic characters proposed

by Simmons and Voss (1998), Lim and Engstrom (2001), Reis *et al.* (2007, 2013). The compilation of the distribution of *P. elongatus* in Brazil was made through records obtained from the scientific literature (see Reis *et al.* 2007, Williams and Genoways 2008).

Five individuals of *Phyllostomus elongatus* (Figure 1) were captured in areas around the caves Gruta dos Moura and Gruta do Urso ($12^{\circ}34' 53''$ S and $46^{\circ}30' 59''$ W, ca. 400 m.a.s.l.) during a bat survey in Aurora do Tocantins, Tocantins State, northern Brazil. *P. elongatus* represented 0.96% of a total of 518 captures of 30 species. Four individuals were captured as they were leaving the Gruta dos Moura cave during the first hours after sundown. Another individual of *P. elongatus* was captured in a gallery forest area about one hour before dawn (Table 1).

According to Nogueira *et al.* (2007), *Phyllostomus elongatus* is a medium-sized bat, with a total body length between 99 and 115 millimeters (mm), and forearm between 61 and 71 mm. Additionally, *P. elongatus* has round-edged ears that are longer than its head length, which differentiates the species from its congeners. Furthermore, *P. elongatus*'s calcaneus is longer than its foot length, which is an important characteristic that distinguishes it from *P. hastatus* and *P. discolor*, with *P. elongatus* having the longest calcaneus. In addition, *P. elongatus* can be distinguished from *P. latifolius* by its forearm length and for having a white spot on the tip of the wings (Koopman 1994, Fregonezi *et al.* 2013).



FIGURE 1. Adult male *Phyllostomus elongatus* (MN78390) captured in karstic area of Aurora do Tocantins, TO, northern Brazil. Photo: Roberto Leonan Morim Novaes.

Captures were conducted in a gallery forest that has similar physiognomic and abiotic characteristics to rainforests, such as high humidity and a canopy that can reach up to 12m in height (Oliveira-Filho and Ratter 2002).

Phyllostomus elongatus has been recorded in 32 Brazilian localities, distributed in 14 states, including the new record for the State of Tocantins (Table 2, Figure 2). Most of the records occur in the Amazon rainforest and the At-




FIGURE 2. Geographic distribution of *Phyllostomus elongatus* in Brazil, including a new record from the Brazilian Cerrado (gray) of Tocantins State (hollow circle).

lantic Forest. Additionally, there is also a record for the Pantanal on an area of semideciduous forest (Bordignon and França 2009). The previous record of *P. elongatus* for the Cerrado was captured in a gallery forest in the transition between the Cerrado and Amazonia (Sousa et al. 2013). The compilation and revision of the geographic distribution of *P. elongatus* suggest that this species is more frequent on forested habitats with higher humidity. Williams and Genoways (2008) made a distribution review of *P. elongatus* noting the possibility of occurrence of this species in the Cerrado of Central Brazil, however the authors did not confirm its occurrence in this region.

TABLE 1. Individuals of *Phyllostomus elongatus* captured in Cerrado of Aurora do Tocantins, Tocantins State, Northern Brazil.

DATE	HOUR	INDIVIDUAL	SEX	REPRODUCTIVE STAGE	SKULL LENGTH	FOREARM LENGTH	BODY MASS
14/jan/12	19:30	MN78390	Male	Abdominal testes	27.8	63.9	44.0
17/jan/12	19:00	MN78388	Male	Abdominal testes	28.9	64.5	30.0
13/jan/12	18:00	MN78389	Female	Inactive	29.1	65.4	33.0
15/jan/12	04:10	Released	Female	Post-lactating	-	62.0	40.0
17/jan/12	19:00	Released	Female	Post-lactating	-	68.2	32.0

TABLE 2. Collecting localities of *Phyllostomus elongatus* in Brazil. Amazonia (Ama), Atlantic Forest (Atl), Cerrado (Cer), and Pantanal (Pan).

LOCALITY	STATE	BIOME	COORDINATES	REFERENCE
PN Montanhas do Tumucumaque	Amapá	Ama	03°12' N, 52°01' W	Martins et al. (2011)
EB Ilha de Maracá	Roraima	Ama	02°27' N, 60°59' W	Robinson (1998)
FLONA do Amapá	Amapá	Ama	01°18' N, 51°52W	Martins et al. (2011)
Macapá	Amapá	Ama	00°02' N, 51°04' W	Mok et al. (1982)
Mazagão	Amapá	Ama	00°07' S, 51°16' W	Piccinini (1974)
Maruanum	Amapá	Ama	00°10' S, 51°16' W	Piccinini (1974)
RDS Rio Iratapuru	Amapá	Ama	00°18' S, 52°26' W	Martins et al. (2011)
Utinga	Pará	Ama	01°20' S, 48°32' W	Handley (1967)
Belém	Pará	Ama	01°27' S, 48°25' W	Mok and Lacey (1980)
Santarém	Pará	Ama	02°24' S, 54°42' W	Bernard (2001)
RDS Amanã	Amazonas	Ama	02°35' S, 64°40' W	Pereira et al. (2010)
PARNA da Amazônia	Pará	Ama	03°48' S, 56°40' W	Marques (1985)
Not provided	Rondônia	Ama	Not provided	Bernard et al. (2011a)
80km N of Manaus	Amazonas	Ama	02°24' S, 59°43' W	Sampaio et al. (2003)
Refúgio do Maruaga	Amazonas	Ama	02°54' S, 59°41' W	Fischer et al. (1997)
Manaus	Amazonas	Ama	03°01' S, 60°07' W	Reis and Guillaumet (1983)
Rio Madeira	Amazonas	Ama	05°27' S, 60°45' W	Bobrowiec (2012)
FLONA de Carajás	Pará	Ama	06°08' S, 50°18' W	Tavares et al. (2012)
Rio Aripuanã	Amazonas	Ama	06°08' S, 60°11' W	Bobrowiec (2012)
PARNA Serra do Divisor	Acre	Ama	07°40' S, 73°40' W	Nogueira et al. (1999)
Recife	Pernambuco	Atl	08°03' S, 34°52' W	Souza-Lopes (1978)
Caruaru	Pernambuco	Atl	08°21' S, 36°01' W	Souza et al. (2004)
Rio Largo	Alagoas	Atl	09°28' S, 35°51' W	Vieira (1953)
Aurora do Tocantins	Tocantins	Cer	12°34' S, 46°30' W	Present Study
Nova Xavantina	Mato Grosso	Cer	14°38' S, 52°21' W	Sousa et al. (2013)
Salvador	Bahia	Atl	12°47' S, 38°27' W	Vieira (1955)
Ilheus	Bahia	Atl	14°47' S, 39°01' W	Faria et al. (2006)
Rio Branco	Mato Grosso	Pan	15°14' S, 58°07' W	Type-locality
Una	Bahia	Atl	15°16' S, 39°04' W	Faria and Baumgarten (2007)
Maciço do Urucum	Mato Grosso do Sul	Pan	19°24' S, 57°22' W	Bordignon and França (2009)
Povoação Village	Espírito Santo	Atl	19°33' S, 39°48' W	Lage et al. (2013)
Teresópolis	Rio de Janeiro	Atl	22°25' S, 42°57' W	Vieira (1942)

ACKNOWLEDGMENTS: We are grateful for the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for supporting the project; Prefeitura Municipal de Aurora do Tocantins, Anselmo Rodrigues, and Wagner José Moura for structural and logistical support on the performed activities; for Ruby Malzoni for extensive revision and correction; for anonymous reviewers and Paúl Velasco for their valuable suggestions.

Studies on the bat fauna of the Cerrado are very recent; the first ones were made in the early 1990's (Aguiar and Zortéa 2008). Although knowledge about the diversity of bats in this biome is still limited and the information about the species distribution in the Cerrado is incipient (Bernard et al. 2011b), this biome presents a high bat diversity (Marinho-Filho 1996, Zortéa and Alho 2008). In addition, bat surveys are lacking or poorly conducted for many areas of the Brazilian Cerrado, including the state of Tocantins (Bernard et al. 2011b, Gregorin et al. 2011), which indicates that new species of bats can still be recorded for this biome.

LITERATURE CITED

- Aguiar, L.M.S. and M. Zortéa. 2008. A diversidade de morcegos conhecida para o Cerrado. *Anais do IX Simpósio Nacional do Cerrado*. Brasília: Empresa Brasileira de Pesquisa Agropecuária. 6 pp.
 Alho, C.J.R., G. Camargo and E. Fischer. 2011. Terrestrial and aquatic mammals of the Pantanal. *Brazilian Journal of Biology* 71(1): 297-310.

- Baker, R.J., C.G. Dunn and K. Nelson. 1988. Allozymic study of the relationships of *Phyllosderma* and four species of *Phyllostomus*. *Occasional Papers, Museum Texas Tech University* 125: 1–14.
- Bernard, E. 2001. Species list of bats (Mammalia, Chiroptera) of Santarém area, Pará State, Brazil. *Revista Brasileira de Zoologia* 18(2): 455–463.
- Bernard, E. 2002. Diet, activity and reproduction of bat species (Mammalia, Chiroptera) in Central Amazonia, Brazil. *Revista Brasileira de Zoologia* 19(1): 173–188.
- Bernard, E., V.C. Tavares and E. Sampaio. 2011a. Compilação atualizada das espécies de morcegos (Chiroptera) para a Amazônia brasileira. *Biota Neotropica* 11(1): 1–12.
- Bernard, E., L.M.S. Aguiar and R.B. Machado. 2011b. Discovering the Brazilian bat fauna: a task for two centuries? *Mammal Review* 41(1): 23–39.
- Bobrowiec, P.E.D. 2012. A Chiroptera preliminary survey in the middle Madeira River region of Central Amazonia, Brazil. *Mammalia* 76: 277–283.
- Bonaccorso, F.J. and N. Smythe. 1972. Punch-marking bats: an alternative to banding. *Journal of Mammalogy* 53(2): 389–390.
- Bordignon, M.O. and A.O. França. 2009. Riqueza, diversidade e variação altitudinal em uma comunidade de morcegos filostomídeos (Mammalia, Chiroptera) no centro-oeste do Brasil. *Chiroptera Neotropical* 15(1): 425–433.
- Eva, H.D., E.E. Miranda, C.M. Di Bella, V. Gond, O. Huber, M. Sgrenzaroli, S. Jones, A. Coutinho, A. Dorado, M. Guimarães, C. Elvidge, F. Achard, A.S. Belward, E. Bartholomé, A. Baraldi, G. De Grandi, O. Vogt, S. Fritz and A. Hartley. 2002. *A vegetation map of South America*. Luxembourg: Official Publications of the European Communities, Joint Research Centre. 48 pp.
- Faria, D. and J. Baumgarten. 2007. Shade cacao plantations (*Theobroma cacao*) and bat conservation in southern Bahia, Brazil. *Biodiversity and Conservation* 16: 291–312.
- Faria, D., B. Soares-Santos and E. Sampaio. 2006. Bats from the Atlantic rainforest of southern Bahia, Brazil. *Biota Neotropica* 6(2): 1–13.
- Fischer, E., W.A. Pedro, S. Borges, M.R. Pinheiro and A. Vicentini. 1997. Predation of *Carollia perspicillata* by *Phyllostomus cf. elongatus* in Central Amazonia. *Chiroptera Neotropical* 3(1): 67–68.
- Fregonezi, M.N., N.R. Reis and A.L. Peracchi. 2013. Subfamília Phyllostominae; pp. 71–106, in: Reis, N.R., M.N. Fregonezi, A.L. Peracchi and O.A. Shibatta (Eds.). *Morcegos do Brasil: guia de campo*. Rio de Janeiro: Technical Books Editora.
- Gregorin, R., E. Gonçalves, C.C. Aires and A.P. Camignotto. 2011. Morcegos (Mammalia, Chiroptera) da Estação Ecológica Serra Geral do Tocantins: composição específica e considerações taxonômicas. *Biota Neotropica* 11(1): 299–311.
- Handley Jr., C.O. 1967. Bats of the canopy of an Amazonian forest. *Atas do Simpósio sobre a Biota Amazônica* 5: 211–215.
- Koopman, K.F. 1994. *Chiroptera: systematics. Handbook of Zoology, VIII (Mammalia)*. Berlin and New York: Walter de Gruyter. 217 pp.
- Lage, S.B., R.S. Cipriano, A.C. Ferreguetti and R.L. Martins. 2013. First record of *Phyllostomus elongatus* (É. Geoffroy, 1810) (Mammalia: Chiroptera) for the state of Espírito Santo, southeastern Brazil. *Check List* 9(4): 880–882.
- Lim, B.K., M.D. Engstrom. 2001. Species diversity of bats (Mammalia: Chiroptera) in Iwokrama Forest, Guyana, and the Guianan subregion: implications for conservation. *Biodiversity and Conservation* 10: 613–657.
- Marinho-Filho, J. 1996. The Brazilian Cerrado bat fauna and its conservation. *Chiroptera Neotropical* 2(1): 37–39.
- Marinho-Filho, J.M. and I. Sazima. 1998. Brazilian bats and conservation biology: a first survey; pp. 282–294, in: T.H. Kunz and P.A. Racey (ed.). *Bat biology and conservation*. Washington: Smithsonian Institution Press.
- Marques, S.A. 1985. Novos registros de morcegos do Parque Nacional da Amazônia (Tapajós), com observações do período de atividade noturna e reprodução. *Boletim do Museu Paraense Emílio Goeldi, série Zoologia* 2(1): 71–83.
- Martins, A.C., E. Bernard, R. Gregorin and W.A.S. Silva. 2011. Filling data on the diversity and distribution of Amazonian bats (Chiroptera): the case of Amapá, easternmost Brazil. *Zoologia* 28(2): 177–185.
- Mok, W.Y. and L.A. Lacey. 1980. Algumas considerações ecológicas sobre morcegos vampiros na epidemiologia da raiva humana na Bacia Amazônica. *Acta Amazônica* 10(2): 335–342.
- Mok, W.I., D.E. Wilson, L.A. Lacey and R.C.C. Luizão. 1982. Lista atualizada de quirópteros da Amazônia Brasileira. *Acta Amazônica* 12(4): 817–823.
- Nogueira, M.R., A.L. Peracchi and R. Moratelli. 2007. Subfamília Phyllostominae; pp. 61–98, in: N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (ed.) *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina.
- Nogueira, M.R., A. Pol and A.L. Peracchi. 1999. New records of bats from Brazil with a list of additional species for the chiropteran fauna of the state of Acre, western Amazon basin. *Mammalia* 63: 363–368.
- Oliveira-Filho, A.T. and J.A. Ratter. 2002. Vegetation physiognomies and woody flora of the Cerrado biome, pp. 91–120, in: P.S. Oliveira and T.J. Marquis (ed.). *Cerrados of Brazil: ecology and natural history of a Neotropical savanna*. New York: Columbia University Press.
- Pereira, M.J.R., J.T. Marques and J.M. Palmerin. 2010. Vertical stratification of bats assemblages in flooded and unflooded Amazonian forests. *Current Zoology* 56(4): 469–478.
- Piccinini, R.S. 1974. Lista provisória dos quirópteros da coleção do Museu Paraense Emílio Goeldi (Chiroptera). *Boletim do Museu Paraense Emílio Goeldi, Série Zoologia* 77: 1–32.
- Reis, N.R. and J.L. Guillaumet. 1983. Les chauves-souris frugivores de la région de Manaus et leur rôle dans la dissémination des espèces végétales. *Revue d'Ecologie la Terre et la Vie* 38: 147–169.
- Reis, N.R. and A.L. Peracchi. 1987. Quirópteros da região de Manaus, Amazonas, Brasil (Mammalia, Chiroptera). *Boletim do Museu Paraense Emílio Goeldi, série Zoologia* 3(2): 161–182.
- Reis, N.R., A.L. Peracchi, W.A. Pedro and I.P. Lima. 2007. *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina. 253 pp.
- Reis, N.R., M.N. Fregonezi, A.L. Peracchi and O.A. Shibatta. 2013. *Morcegos do Brasil: guia de campo*. Rio de Janeiro: Technical Books Editora. 252 pp.
- Robinson, F. 1998. The bats of the Ilha de Maracá; pp. 165–188, in: J.A. Ratter and W. Milliken (ed.). *Maracá: the biodiversity and environment of an Amazonian Rainforest*. London: John Wiley & Sons.
- Sampaio, E.M., E.K.V. Kalko, E. Bernard, B. Rodríguez-Herrera and C.O. Handley Jr. 2003. A biodiversity assessment of bats (Chiroptera) in a tropical lowland rainforest of Central Amazonia, including methodological and conservation considerations. *Studies on Neotropical Fauna and Environment* 38(1): 17–31.
- Sikes, R.S., W.L. Gannon and Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92(1): 235–253.
- Simmons, N.B. and R.S. Voss. 1998. The mammals of Paracou, French Guiana: a Neotropical lowland rainforest fauna, Part 1. Bats. *Bulletin of the American Museum of Natural History* 237: 1–219.
- Sousa, R.F., P.C. Venere and K.C. Faria. 2013. Bats in forest remnants of the Cerrado savanna of eastern Mato Grosso, Brazil. *Biota Neotropica* 13(2): 326–341.
- Souza M.A.N., A. Langguth and E.A. Gimenez. 2004. Mamíferos dos brejos de altitude da Paraíba e Pernambuco; pp. 229–254, in: K.C. Porto, J.J.P. Cabral and M. Tabarelli (ed.). *Brejos de altitude em Pernambuco e Paraíba: história natural, ecologia e conservação*. Brasília: Ministério do Meio Ambiente.
- Souza-Lopes, M.J. 1978. Cariótipo de duas espécies de morcegos de Pernambuco (Chiroptera, Phyllostomatidae). *Revista Nordestina de Biologia* 1: 113–17.
- Straube, F.C. and G.V. Bianconi. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com utilização de redes-de-neblina. *Chiroptera Neotropical* 8: 150–152.
- Tavares, V.C., C.F.S. Palmuti, R. Gregorin and T.T. Dornas. 2012. Morcegos; pp. 162–179, in: F.D. Martins, A.F. Castilho, J. Campos, F.M. Hatano and S.G. Rolim (org.). *Fauna da Floresta Nacional de Carajás: estudos sobre vertebrados terrestres*. São Paulo: Nitro Editorial, Belo Horizonte.
- Tuttle, M.D. 1970. Distribution and zoogeography of Peruvian bats, with comments on natural history. *The University of Kansas, Science Bulletin* 49(2): 45–86.
- Vieira, C.O.C. 1942. Ensaio monográfico sobre os quirópteros do Brasil. *Arquivos de Zoologia do Estado de São Paulo* 3: 219–471.
- Vieira, C.O.C. 1953. Sobre uma coleção de mamíferos do estado de Alagoas. *Arquivos de Zoologia do Estado de São Paulo* 8: 209–224.
- Vieira, C.O.C. 1955. Lista remissiva dos mamíferos do Brasil. *Arquivos de Zoologia do Estado de São Paulo* 8: 341–474.
- Williams, S.L. & H.H. Genoways. 2008. Subfamily Phyllostominae; pp. 255–299, in: A.L. Gardner, (ed.). *Mammals of South America Volume 1: marsupials, xenarthrans, shrews, and bats*. Chicago: The University of Chicago Press.
- Zortéa, M. and C.J.R. Alho. 2008. Bat diversity of a Cerrado habitat in central Brazil. *Biodiversity and Conservation* 17: 791–805.

RECEIVED: August 2013

ACCEPTED: November 2013

PUBLISHED ONLINE: February 2014

EDITORIAL RESPONSIBILITY: Paúl M. Velazco