

## *Ceratophrys joazeirensis* Mercadal, 1986 (Anura: Ceratophryidae): New state record

Daniel Oliveira Santana<sup>1\*</sup>, Renato Gomes Faria<sup>2</sup>, Francis Luiz Santos Caldas<sup>1</sup> and Crizanto Brito De-Carvalho<sup>2</sup>

<sup>1</sup> Universidade Federal da Paraíba, Programa de Pós Graduação em Ciências Biológicas (Zoologia), Cidade Universitária, CEP 58059-900, João Pessoa, PB Brasil.

<sup>2</sup> Universidade Federal de Sergipe, Programa de Pós-graduação em Ecologia e Conservação. Cidade Universitária Prof. José Aloísio de Campos, Avenida Marechal Rondon, Jardim Rosa Elze. CEP 49100-000. São Cristóvão, SE, Brazil.

\* Corresponding author. E-mail: [danielbioufs@yahoo.com.br](mailto:danielbioufs@yahoo.com.br)

**ABSTRACT:** This article provides new records to *Ceratophrys joazeirensis* of two new localities from the state of Sergipe, northeast Brazil. These records extend the geographical distribution of the species for municipalities from Poço Redondo and Canindé do São Francisco. These points are the first records of *Ceratophrys joazeirensis* for the State of Sergipe.

The genus *Ceratophrys* Wied-Neuwied, 1824 currently comprises eight species occurring in Tropical South America (Frost 2013) in forest and non-forest areas (Lynch 1982). Five species occur in Brazil *Ceratophrys aurita* (Raddi, 1823), *Ceratophrys cornuta* (Linnaeus, 1758), *Ceratophrys cranwelli* Barrio, 1980 *Ceratophrys joazeirensis* Mercadal de Barrio, 1986 and *Ceratophrys ornata* (Bell, 1843). The Caatinga Horned Frog *Ceratophrys joazeirensis* Mercadal (1986) is the only species of the genus that occurs in the Caatinga biome, a typical semi-arid morphoclimatic domain in northeastern Brazil (Ab'Saber 2003).

*Ceratophrys joazeirensis* has no population information available in the literature, since it is known from only a few specimens. It is a species of dry Caatinga savannah, which presumably breeds in temporary ponds by larval development (Skuk and Juncá 2004).


This species is reported from the type locality (vicinities of Juazeiro, Bahia, Brazil; Mercadal 1986), north of Triunfo, Pernambuco (Santos *et al.* 2009), Araruna, northern Paraíba, from the vicinities of Passa-e-Fica, northern Rio Grande do Norte (Vieira *et al.* 2006), from Santa Maria, Rio Grande do Norte (Jorge *et al.* 2012) and from Pindobaçu, Bahia (Zaidan and Leite 2012). Recently, two new records were reported, from São Raimundo Nonato, Piauí (Roberto *et al.* 2013) and Buritizeiro, Minas Gerais (Maciel *et al.* 2013).

2013). We also present a map (Figure 1) and data in Table 1 of the current known distribution of this species based on literature.

The present study relates the occurrence of *C. joazeirensis* (Figure 2) in two localities in Caatinga area in Sergipe, in northeast Brazil. The first record corresponds Serra da Guia (09°58'52" S, 37°52'05" W, 727 m above sea level) in the municipality of Poço Redondo, Sergipe (Figure 1). The second record corresponds Xingó Power Plant (09°37'38" S, 37°48'08" W; 100 above sea level) in the municipality of Canindé do São Francisco, Sergipe (Figure 1). This records extends the known distribution of *C. joazeirensis* approximately 300 km east of the type locality. The individuals of *C. joazeirensis* were collected, fixed in formalin 4% and preserved in alcohol 70%. The specimen had their identification confirmed based on color patterns of head as a blot between the interocular space to the nostrils and upper eyelids fairly pronounced, based on the description of Mercadal (1986) and the nomenclature followed the classification recommended by the Brazilian Herpetology Society (Segalla *et al.* 2012). Voucher specimens were deposited at the Herpetological Collection of the Universidade Federal de Sergipe (CHUFS), (C348 and C353). Collecting permit (# 10504-1) was granted by Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis - IBAMA.

**TABLE 1.** Localities of known distribution of *Ceratophrys joazeirensis*.

STATE	MUNICIPALITY	LATITUDE (S)	LONGITUDE (W)	REFERENCE
Bahia	Juazeiro	09°24'38,00"	40°30'26,00"	Mercadal (1986)
Bahia	Pindobaçu	10°44'44,84"	40°21'28,65"	Zaidan and Leite (2012)
Minas Gerais	Buritizeiro	17°21'03,00"	44°57'43,00"	Maciel <i>et al.</i> (2013)
Pernambuco	Triunfo	07°50'17,48"	38°06'06,59"	Santos <i>et al.</i> (2009)
Paraíba	Araruna	06°26'07,00"	35°39'41,00"	Vieira <i>et al.</i> (2006)
Piauí	São Raimundo Nonato	09°01'05,46"	42°42'09,66"	Roberto <i>et al.</i> (2013)
Rio Grande do Norte	Passa-e-Fica	06°26'15,19"	35°38'37,61"	Vieira <i>et al.</i> (2006)
Rio Grande do Norte	Santa Maria	05°51'14,40"	35°42'03,60"	Jorge <i>et al.</i> (2012)
Sergipe	Poço Redondo	09°58'52,00"	37°52'05,00"	New Record
Sergipe	Canindé do São Francisco	09°37'38,80"	37°48'08,38"	New Record



**FIGURE 1.** Geographic distribution of *Ceratophrys joazeirensis* in Brazil.



**FIGURE 2.** *Ceratophrys joazeirensis* (C348). Photo by Crizanto Brito De Carvalho.

**ACKNOWLEDGMENTS:** We thank Universidade Federal de Sergipe and Secretaria de Meio Ambiente e Recursos Hídricos – SEMARH/SE for logistic. The Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for fellowships, and an anonymous reviewer for their helpful comments and suggestions in of this manuscript.

#### Literature Cited

- Ab'Saber, A.N. 2003. *Os domínios de natureza no Brasil: potencialidades paisagísticas*. São Paulo: Ateliê Editorial. 159 pp.
- Frost, D.R. 2013. *Amphibian Species of the World: an Online Reference*. Version 5.6 (9 January 2013). Accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA. Captured on 17 September 2013.
- Jorge, J.S., M.N.C. Kokubum, R.D.F. Sales and E.M.X. Freire. 2012. Geographic distribution: *Ceratophrys joazeirensis*. *Herpetological Review* 43(2): 297–298.
- Lynch, J.D. 1982. Relationships of the frogs of the genus *Ceratophrys* (Leptodactylidae) and their bearing on hypotheses of Pleistocene forest refugia in South America and punctuated equilibria. *Systematic Zoology* 31(2): 166–179.
- Maciel, N.M., M.N. C. Kokubum, P.H.P. Braga, A.T. Queiróz-Júnior and R.H. Matsushita. 2013. Distribution extension, new state record and geographic distribution map of *Ceratophrys joazeirensis* Mercadal, 1986 (Anura: Ceratophryidae). *Herpetology Notes* 6: 447–450.
- Mercadal de Barrio, I.T. 1986. *Ceratophrys joazeirensis* sp. n. (Ceratophryidae, Anura) del noreste de Brasil. *Amphibia-Reptilia* 7(4): 313–334.
- Roberto, I.J., S.C. Ribeiro, and D. Loebmann. 2013. Amphibians of the state of Piauí, Northeastern Brazil: a preliminary assessment. *Biota Neotropica* 13(1): 322–330.
- Santos, E.M., G.L. Silva, T.F. Campo and M.S. Quirino. 2009. Geographic distribution: *Ceratophrys joazeirensis*. *Herpetological Review* 40(1): 107.
- Segalla, M, U. Caramachi, C.A.G. Cruz, P.C.A. Garcia, T. Grant, C.F.B. Haddad and J. Langone. 2012. *Brazilian amphibians – list of species*. Accessible at <http://www.sbsherpetologia.org.br>. Captured on 04 December 2013.
- Skuk, G. and F. Juncá 2004. *Ceratophrys joazeirensis*, in: IUCN 2013. *IUCN Red List of Threatened Species*. Version 2013.2. Accessible at <http://www.iucnredlist.org>. Captured on 04 December 2013.
- Vieira, K.S., C. Arzabe, and W.L.S. Vieira. 2006. Amphibia, Ceratophryidae, Ceratophryinae, *Ceratophrys joazeirensis*: distribution extension. *Check List* 2(2): 28–29.
- Zaidan, B.F. and F.S.F. Leite. 2012. Advertisement call of the rare, explosive breeding caatinga horned frog *Ceratophrys joazeirensis* Mercadal de Barrio, 1986 (Anura, Ceratophryidae). *Zootaxa* 3540: 65–66.

RECEIVED: December 2013

ACCEPTED: February 2014

PUBLISHED ONLINE: May 2014

EDITORIAL RESPONSIBILITY: Camila C. Both