



# *Typha orientalis* Presl (Typhaceae): a new species record for India

Aijaz Hassan Ganie<sup>1\*</sup>, A. R. Dar<sup>2</sup>, Mehboob Ashraf<sup>2</sup> and Zafar A. Reshi<sup>1</sup>

<sup>1</sup> University of Kashmir, Department of Botany, Srinagar, 190 006, Jammu and Kashmir, India

<sup>2</sup> Government Degree College Ganderbal, Jammu and Kashmir, India

\* Corresponding author. E-mail: [aijazku@gmail.com](mailto:aijazku@gmail.com)

**Abstract:** *Typha orientalis* C. Presl (Typhaceae) is recorded for the first time from the Kashmir Himalaya, India and for the first time in the entire Indian sub-continent. A detailed taxonomic description and photographs of the diagnostic characters are provided to facilitate its identification in the field. Also provided are diagnostic characters used to distinguish *T. orientalis* C. Presl from *T. latifolia* L.

**Key words:** *Typha latifolia*, Kashmir Himalaya, Indian subcontinent

*Typha* L. (Typhaceae), or cattails, includes 10 to 15 species and comprises highly productive aquatic weeds (Boyd 1971) and rapid colonizers of disturbed or newly formed wetlands (Steinbachova-Vojtiskova *et al.* 2006). The genus is widely distributed in temperate and tropical regions of the Northern and Southern Hemispheres, existing on all continents except Antarctica (Smith 1987; Kim 2002).


In the Indian subcontinent, four species of *Typha* (*T. angustata* Bory & Chaub., *T. elephantina* Roxb., *T. javanica* Schnizl. and *T. laxmannii* Lepech.) have been reported (Hooker 1893). Saha (1968) reported 4 species from India, which include *T. angustata*, *T. elephantina*, *T. latifolia* and *T. laxmannii*, of which the first two occur throughout India while the other two are restricted to Kashmir, Punjab, Decan and probably Gujarat. Cook (1996) recorded five species (*T. latifolia*, *T. laxmannii*, *T. minima*, *T. domingensis* [= *T. javanica*] and *T. elephantina*) from India. Of these, *T. latifolia*, *T. laxmannii* and *T. minima* are restricted in distribution to Western Himalayan part in India. Kaul and Zutshi (1967) recorded only *T. angustata* from the wetlands of Srinagar, India. Stewart (1972) recorded 3 species of the genus (*T. angustata* [= *T. domingensis*], *T. elephantina*, and *T. laxmannii*) from the whole of the former Jammu and Kashmir. Kak (1990) reported 3 species (*T. angustata*, *T. elephantina*, and *T. laxmannii*) from Kashmir Himalaya, India.

While carrying out botanical surveys in the Kashmir Valley, full blooming specimens of *Typha* species were collected from the Ganderbal (1,590 m above sea level [a.s.l.]), Gadura (1,590 m a.s.l.), Nigeen Lake (1,595 m a.s.l.), Sundoo (1605 m a.s.l.), Awantipora (1,600 m a.s.l.), Ahansar wetland (1,585 m a.s.l.) Narbal (1,603 m a.s.l.), Shalbugh wetland (1,590 m a.s.l.), Wular Lake (1,645 m a.s.l.). After critical examination of morphological characters, the

specimens were identified as *Typha orientalis* C. Presl. Study of the relevant taxonomic literature (Hooker 1893; Kaul and Zutshi 1967; Stewart 1972; Kak 1990; Cook 1996) revealed that this species is previously unreported from India. Therefore, this report represents the first record of *Typha orientalis* C. Presl for the flora of India. A detailed taxonomic description and photographs of diagnostic characters of this species are given here to facilitate its identification in the field and also to validate this new record for India.

*Typha orientalis* is distributed in China (Anhui, Guangdong, Guizhou, Hebei, Heilongjiang, Henan, Hubei, Jiangsu, Jiangxi, Jilin, Liaoning, Nei Mongol, Shaanxi, Shandong, Shanxi, Taiwan, Yunnan, Zhejiang), Japan, Korea, Mongolia, Myanmar, Philippines, Russia, Australia and now India (Kashmir Himalaya).

The study site was in the Kashmir Valley, situated in northern fringe of the Indian sub-continent between 33°22' N and 34°50' N and 073°55' E and 073°33' E (Figure 1), covering an area of ca. 16,000 km<sup>2</sup>. Standard herbarium methods were followed (Ganie *et al.* 2014).



**Figure 1.** Map of Kashmir Himalaya showing study sites (1, Wular Lake; 2, Ahansar; 3, Ganderbal; 4, Gadura; 5, Shalbugh; 6, Narbal; 7, Nigeen Lake; 8, Awantipora; 9, Sundoo).

***Typha orientalis*** C. Presl, *Abh. Königl. Böhm. Ges. Wiss.*, ser. 5, 6: 599; *Epimel. Bot.* 239. 1851.

Vernacular (Kashmiri) name: Zab/Peitz

Perennial herb (Figures 2 and 3) up to 270 cm tall; rhizome robust creamish white in colour; stem with nodes and internodes, solid, stout clasped by leaf sheaths; leaves dark green, leaf lamina of lower leaves 140–190 × 2–2.5 cm, and upper leaves 80–155 × 1.3–1.8 cm, leaf sheath of lower leaves 45–110 cm and 45–80 cm of lower leaves, abaxially convex, transverse section semicircular to lanceolate; male and female parts of inflorescence contiguous, male part of spikes 12–17 cm, with 1–3 deciduous bracts at base or occasionally in middle portion and upper portion; female part of spikes 11–24 cm, with 1 deciduous bract at base; male flowers: stamens 3, rarely 2 or 4; anthers ca. 3 mm; female flowers without bracteoles; ovary fusiform to lanceolate; stalk 2–2.5 mm, slender; styles 1.5–3 mm; stigmas spatulate, 0.5–0.8 mm; hairs on stalk ca. as long as style; fruit elliptic.

**SPECIMENS EXAMINED:** India, Jammu and Kashmir, Ganderbal, 5 June 2014, Aijaz, Rashid and Mehboob 01991 [Kashmir University Herbarium (KASH)]; Gadura-Ganderbal, 5 June 2014, Aijaz, Rashid and Mehboob 01992 (KASH); Nigeen Lake, 7 June 2014, Aijaz, Rashid and Mehboob 01993 (KASH); Sundoo-Anantnag, 15 June 2014, Aijaz, Rashid and Mehboob 01994 (KASH); Awantipora-Pulwama, 20 June 2014, Aijaz, Rashid and Mehboob 01995 (KASH); Narbal-Srinagar, 1 July 2014, Aijaz, Rashid and Mehboob 01996 (KASH), Shalbugh-Ganderbal, 01 July 2014, Aijaz, Rashid and Mehboob 01997 (KASH); Wular Lake 10 July 2014, Aijaz, Rashid and Mehboob 01998 (KASH).

**ADDITIONAL SPECIMENS EXAMINED** (determination of specimens deposited in Kashmir University Herbarium [KASH]): After critical examination of the previous herbarium specimens, it was also found that many specimens of *Typha orientalis* in the Kashmir University Herbarium (KASH), which had been deposited over a period of time were misidentified are given as *T. latifolia*: India, Kashmir: Kishtiwari, 20 July 2007, A.A. Khuroo, Akhtar H. Malik and Z.S. Khan 639; Saidakadal-Nigeen Lake, 13 September 2012, Mahpara 52.

During this study, *Typha orientalis* was collected from Srinagar, Ganderbal, Anantnag, Baramulla, Bandipora and Pulwama districts of the Kashmir Himalaya (Figure 1).

**Flowering period:** The species are in full bloom during May–July in different aquatic habitats of Kashmir Himalaya.

**Fruiting period:** It was observed that the species produce fruits during September to October

The taxonomy and systematics of *Typha* is unclear and controversial. This is due to species having very similar gross morphology and a large range of variability in vegetative and reproductive traits that are used to delimit taxa (Smith 1967). Since the early systematic treatment of Kronfeld (1889),

several regional revisions have tried to establish taxonomic boundaries between taxa based on morphological traits (Saha 1968; Cook 1980; Sharma and Gopal 1980; Bokhari 1983; Finlayson et al. 1985; Kuehn and White 1999; Hamdi and Assadi 2003, Kim et al. 2003). Other revisions have focused on particular species complexes (Kuehn et al. 1999) or correct taxonomic identification of various species of the genus (Sharma and Gopal 1980).

Morphologically, *Typha orientalis* appears similar to *T. latifolia* because the male and female parts of the inflorescence in both the species are not separated; both species are ebracteolate. Based on the distinguishing characters of the two species (Table 1), both *T. orientalis* and *T. latifolia* occur in Kashmir Himalaya.

In Kashmir Himalaya, the bewildering taxonomy of the genus *Typha* may be resolved in the future by molecular studies.

## ACKNOWLEDGEMENTS

We are thankful to the Head, Department of Botany, University of Kashmir, Srinagar, for providing necessary facilities. We acknowledge the kind help provided by non-teaching staff, particularly Mr. Zahoor Ahmad Reshi Department of Botany, Government Degree College Ganderbal.


## LITERATURE CITED

- Bokhari, M.H. 1983. The aquatic plants of Iran and Pakistan. III. Typhaceae. *Biologia* 29: 85–91.
- Boyd, C.E. 1971. Further studies on productivity, nutrient and pigment relationships in *Typha latifolia* populations. *Bulletin of the Torrey Botanical Club* 98: 144–150.
- Cook, C.D.K. 1980. Typhaceae; pp. 275–276, in: Tutin, T.G., V.H. Heywood, N.A. Burges, D.A. Valentine, S.M. Walters and D.A. Webb (eds.). *Flora Europaea*, Vol. 5. Cambridge: Cambridge University Press.
- Cook, C.D.K. 1996. *Aquatic and wetland plants of India*. New Delhi: Oxford University Press. 385 pp.
- Finlayson, M., R.I. Forrester, D.S. Mitchell and A.J. Chick. 1985. Identification of native *Typha* species in Australia. *Australian Journal of Botany* 33: 101–107.
- Ganie, A.H., B.A. Tali, A.K. Khuroo, I.A. Nawchoo and A.M. Rather. 2014. *Rheum spiciforme* Royle (Polygonaceae): a new record to the flora of Kashmir Valley, India. *National Academy Science Letters* 37(6): 561–565.
- Hamdi, S.M.M. and M. Assadi. 2003. Typhaceae; pp.299–317, in: Hamdi, S.M.M. and M. Assadi. (eds.), *Flora of Iran* 42. Berlin/Hamburg: Verlag Paul Parey.
- Hooker, J.D. 1893. *The Flora of British India*. Vol. 6. L. London: Reeve & Co., Vol. VI. 488–489.
- Kak, A.M. 1990. Aquatic and wetland vegetation of the Kashmir Himalaya. *Journal of Economic and taxonomic Botany* 14(1): 1–14.
- Kaul, V. and D.P. Zutshi. 1967. A study of aquatic and marshland vegetation of Srinagar. *Proceedings of the Natural Institute of Science, India*, B 33(3–4): 111–127.

**Table 1.** Morphological comparison between *Typha orientalis* and *T. latifolia*.


Trait	<i>T. orientalis</i>	<i>T. latifolia</i>
Leaf lamina	50–118 × 1.5–2.5cm, dark green in colour, T.S. linear-lanceolate to lanceolate (Figure 2)	50–105 × 1.0–1.5cm, light green in colour, T.S. semilunar (Figure 2)
Male flower	Usually 3 staminate	Usually 2 staminate
Female flower	Hairs on stalk as long as style, gynophore 2–2.5 mm long, ovary fusiform, stigma spatulate (Figure 2)	Hairs on stalk shorter than style, gynophore 3.5–4.0 mm long, ovary lanceolate, stigma lanceolate (Figure 2)
Fruit	Elliptic	Lanceolate





**Figure 2:** Vegetative and floral parts of *T. orientalis* **A:** Natural population (scale = 0.01 mm). **B:** Habit (scale = 0.02 mm) **C:** Leaf apical portion (scale = 0.06 mm). **D:** Inflorescence (scale = 0.1 mm). **E:** Male flower (scale = 10 cm). **F:** Female flower; note hairs on stalk as long as style (scale = 7.5 cm). **G:** Female flower with spatulate stigma (scale = 9 cm). **H:** Carpel with gynophore (scale = 6.5 m). **I:** Fusiform ovary (scale = 20 cm).





**Figure 3. A–D: *Typha orientalis*.** **A:** T.S. of leaf (scale = 2 cm). **B:** T.S. of female part of inflorescence; note hairs on stalk as long as style (scale = 2.5 cm). **C:** Female flower; note fusiform ovary and medium sized gynophore (scale = 3.7cm). **D:** Spathulate stigma (scale = 30 cm). **E–G: *Typha latifolia*.** **E:** T.S. of leaf (scale = 2 cm). **F:** T.S. of female part of inflorescence; note hairs on stalk shorter than style. **G:** Female flower; note lanceolate ovary and long gynophore (scale = 3.3 cm). **H:** Lanceolate stigma (scale = 25 cm).

- Kim, C. 2002. Morphology and molecular systematic of Typhaceae. Dissertation, Ajou University, Suwon, Republic of Korea.
- Kim, C., H. Shin and H. K. Choi. 2003: A phenetic analysis of *Typha* in Korea and Far East Russia. *Aquatic Botany* 75: 33–43.
- Kronfeld, M. 1889. Monographie der Gattung *Typha* Tourn. Verhandlungen der Zoologisch-Botanischen Gesellschaft 39: 89–192.
- Kuehn, M.M., J.E. Minor and B.N. White. 1999. An examination of hybridization between the cattail species *Typha latifolia* and *Typha angustifolia* using random amplified polymorphic DNA and chloroplast DNA markers. *Molecular Ecology* 8: 1981–1990. doi: [10.1046/j.1365-294x.1999.00792.x](https://doi.org/10.1046/j.1365-294x.1999.00792.x)
- Saha, S. 1968. The genus *Typha* in India—its distribution and uses. *Journal of the Botanical Society of Bengal* 22: 11–18.
- Sharma, K.P. and B. Gopal. 1980. A note on the identity of *Typha elephanita* Roxb. *Aquatic Botany* 9: 381–387.
- Smith, S.G. 1967. Experimental and natural hybrids in North America *Typha* (Typhaceae). *American Midland Naturalist* 78: 257–287.
- Steinbachova-Vojtiskova, L., E. Tylova, A. Soukup, H. Novicka, O. Votrubova, H. Lipavska and H. Cizkova. 2006. Influence of nutrient supply on growth, carbohydrate, and nitrogen metabolic relations in *Typha angustifolia*. *Environmental and Experimental Botany* 57: 246–257.
- Stewart, R.R. 1972. An annotated catalogue of the vascular plants of West Pakistan and Kashmir. Karachi: Fakhri Printing Press. xviii + 1028 pp.
- Authors' contribution statement:** AHG, ARD and MA has collected the plant material, the analysis of data were carried out in the laboratory by AHG, ARD, MA and ZAR. AHG, ARD and ZAR wrote the manuscript.
- Received:** October 2014
- Accepted:** January 2015
- Editorial responsibility:** Nik Fadzly