



# Checklist of phytobenthos from Boipeba Island, Bahia, Brazil, emphasizing the morphological features of *Nitophyllum punctatum* (Rhodophyta, Ceramiales)

**Carlos Wallace do Nascimento Moura\***, **Aigara Miranda Alves**, **Alana Araujo dos Santos**,  
**Wellington Romualdo de Almeida**, **Kátia Lidiane Moniz-Brito**, **Ingryd Santana de Oliveira**,  
**Nelma dos Santos Freitas**, **Ivania Batista de Oliveira** and **Geraldo José Peixoto Ramos**

Universidade Estadual de Feira de Santana, Departamento de Ciências Biológicas, Laboratório de Ficologia, Avenida Transnordestina s/n, Bairro Novo Horizonte, CEP 44036-900, Feira de Santana, BA, Brazil

\* Corresponding author. E-mail: [wallace@uefs.br](mailto:wallace@uefs.br)

**Abstract:** We report here the results of a survey of the phytobenthos of Boipeba Island, Bahia state, Brazil. Samples were collected during three periods (July 2007 and March and December 2012) in the intertidal regions of offshore reefs near the beaches at Ponta dos Castelhanos, Bainema, Moreré, and Tassimirim. A total of 159 taxa were identified: Rhodophyta (83), Chlorophyta (52), Heterokontophyta/Phaeophyceae (21), Cyanobacteria (02), and Magnoliophyta (01). The most representative orders were Ceramiales (Rhodophyta), Bryopsidales (Chlorophyta), and Dictyotales (Heterokontophyta/Phaeophyceae) with 30, 23 and 15 taxa respectively; vouchers were deposited in the herbarium of the Universidade Estadual de Feira de Santana (HUEFS). This work adds 150 new taxa to those already known for Boipeba Island, including *Nitophyllum punctatum* (Stackh. in With.) Grev., whose occurrence is now confirmed for the coast of Bahia. The number of species identified in this study was elevated in comparison with previous phytobenthos studies undertaken on Brazilian islands.

**Key words:** algae, marine flora, northeastern coast, reef communities

## INTRODUCTION

Bahia has the most extensive shoreline (approximately 1,100 km) of any Brazilian state (Bahia 1999), comprising a wide variety of coastal environments and one of the most highly diverse phycological floras in the country (Pereira 2002; Nunes and Guimarães 2008; Bicudo and Menezes 2010).

Although the phycological benthic marine flora of Bahia has been relatively well studied (Nunes 1998;

Széchy and Paula 2010; Creed et al. 2010; Moura 2010; Santos and Moura 2011; Alves et al. 2011a, 2011b, 2012a, 2012b, 2012c; Jesus 2012; Almeida et al. 2012; Almeida et al. 2013; Torrano-Silva and Oliveira 2013; Almeida et al. 2014; Moura et al. 2014a), there is still much to be discovered about its coastal ecosystems (especially along the extreme northern and southern beaches) and oceanic islands (such as the Abrolhos Archipelago), as well as the Todos os Santos and Camamu bays and their associated islands (such as Tinharé and Boipeba).

Boipeba Island is part of the Tinharé-Boipeba Environmental Protection Area - APA (created by State Decree n° 1.240, of 05/06/1992) and is located approximately 500 m off the southern coast of Bahia state near Camamu Bay (Rio dos Patos). There are also fringe reefs in the area bordering the islands (that remain submerged during the syzygy low tides), shallow reefs at depths between 5 and 10 m, as well as deep reefs at depths from 10 to 20 m (Kikuchi et al. 2008).

Kikuchi et al. (2008) recorded the presence of nine coral species and one hydrocoral species on the reefs in the Tinharé-Boipeba APA and Camamu during a visual census using AGRRA (Ginsburg et al. 1998) and PETRORISCO (Dutra and Kikuchi 2003) protocols. The endemic species *Mussismilia hispida* Verrill, 1902, *Mussismilia brasiliensis* Verrill, 1867, *Mussismilia hartti* Verrill, 1868, *Favia gravida* Verrill, 1868, and *Siderastrea stellata* Verrill, 1868, and the hydrocoral *Millepora alcicornis* L., 1758 are common elements of the reef fauna in the area. According to Kikuchi et al. (2008), the corals on this reef area grow in pools and canals in association with large numbers of organisms that constitute the reef communities, such as fish, crustaceans, mollusks, segmented worms, etc., and large quantities of algae (coralline, filamentous, and fleshy algae). Kikuchi et al.

(2008) pointed out the importance of green algae along the reefs at Tinhare-Boipeba and Camamu, as they cover approximately 80% of the shallow reef areas there.

In spite of the great diversity of phytobenthos at Boipeba Island, there had been no specific studies of its phycological flora. A number of species are known to the island, including *Halymenia duchassaingii* (J.Agardh) Kylin (Nunes 2005), *Rhizoclonium africanum* Kütz., and *Valonia aegagropila* C.Agardh (Alves et al. 2009, 2010), as well as three species of *Cladophora* (*C. corallicola* Børgesen, *C. dalmatica* Kütz. and *C. vagabunda* [L.] C.Hoek) (Alves et al. 2012c), *Hypnea musciformis* (Wulfen in Jacq.) J.V.Lamour. (Jesus 2012), *Parvocaulis pusillus* (M.Howe) S.Berger et al., and *Parvocaulis parvulus* (Solms) S.Berger et al. (Moura et al. 2014a). In terms of cyanobacteria, Caires (2013) reported the presence of *Symploca* cf. *hydnoides* Gomont.

Studies of the structures of phytobenthic communities have been used to evaluate environmental impacts on marine environments. According to Oliveira et al. (1999), the biodiversity of marine organisms is correlated to a certain degree with algal community diversity — and this diversity increases ecosystem stability — as the presence of greater numbers of functionally equivalent species with different levels of tolerance to environmental factors make these systems better able to resist environmental impacts (including alterations caused by anthropogenic activities).

The aim of the present study was to undertake a taxonomic survey of the phytobenthos at Boipeba Island, Bahia state and confirm the occurrence of *Nitophyllum punctatum* (Stackh. in With.) Grev. along the Brazilian coast, based on its morphological and reproductive features.

## MATERIALS AND METHODS

The study area includes four beaches: Tassimirim ( $13^{\circ}34' S$ ,  $038^{\circ}55' W$ ), Moreré ( $13^{\circ}35' S$ ,  $038^{\circ}54' W$ ), Bainema ( $13^{\circ}38' S$ ,  $038^{\circ}53' W$ ), and Ponta dos Castelhanos ( $13^{\circ}40' S$ ,  $038^{\circ}54' W$ ), located on Boipeba Island. Boipeba Island is part of Cairu city, on the southeastern coast of Bahia state, in northeastern Brazil (between the latitudes  $13^{\circ}22'$  and  $13^{\circ}51' S$ , and longitudes  $38^{\circ}51'$  and  $39^{\circ}03' W$ ; Figure 1).

Sampling was undertaken at random during three different periods (July 2007 and March and December 2012) in the intertidal region of the coastal reefs on the four beaches and in the Bainema mangrove swamp; the specimens were removed with the aid of spatulas and held in labeled plastic bags until processed.

The specimens were fixed and preserved in 4% formaldehyde (using seawater) and subsequently identified with the aid of both a stereomicroscope and a light microscope. The taxonomic arrangement used follows Wynne (2011), with the exception of *Feldmannia*

*mitchelliae* (Harv.) H.S.Kim, which was listed in Wynne (2011) as *Hincksia mitchelliae* (Harv.) P.C.Silva, but was considered a synonym by Kim (2010). The taxon author names follow Brummitt and Powell (1992). The voucher specimens were deposited in the Herbarium of the Universidade Estadual de Feira de Santana (HUEFS), Bahia state, Brazil.

## RESULTS AND DISCUSSION

A total of 159 taxa were identified (Table 1). The most representative was the phylum Rhodophyta, with 83 taxa, followed by phylum Chlorophyta (52), Heterokontophyta/Phaeophyceae (21), Cyanobacteria (2), and Magnoliophyta (1).


This high richness of red algae (Rhodophyta) had previously been observed in surveys in tropical and warm-temperate regions by Kain and Norton 1990, Marins et al. 2008, and Torrano-Silva and Oliveira 2013. Among the 21 orders encountered, the most representative were: Ceramiales (Rhodophyta) with seven families and 30 taxa; Bryopsidales (Chlorophyta) with five families and 23 taxa; and Dictyotales (Heterokontophyta/ Phaeophyceae), with one family (Dictyotaceae) being represented by 15 taxa (Figure 2). In addition to the latter family, Rhodomelaceae and Gracilariaeae (Rhodophyta) also demonstrated high floristic richness, with 17 and nine taxa respectively. The most representative genera were: *Caulerpa* J.V.Lamour. (with 11 taxa), *Gracilaria* Grev. (9), *Cladophora* Kütz. (6), *Dictyota* J.V.Lamour. (5), followed by *Amphiroa* J.V.Lamour., *Bostrychia* Mont. and *Hypnea* J.V.Lamour. (with 4 taxa each).

The number of species encountered in the present study was similar to previous surveys undertaken on Brazilian oceanic islands. Collections at Fernando de Noronha Island yielded 172 species (Villaça et al. 2006); Atol das Rocas, 93 species (Oliveira and Ugadim 1974, 1976; Villaça et al. 2010); São Pedro and São Paulo, 39 species (Villaça et al. 2006); Trindade and Martin Vaz, 131 species (Villaça et al. 2006); and the Abrolhos Archipelago, 164 species (Torrano-Silva and Oliveira 2013).

Ceramiales and Bryopsidales were the most representative orders in surveys undertaken by Villaça et al. (2006), Villaça et al. (2010), and Torrano-Silva & Oliveira (2013). High numbers of Chlorophyta were encountered on the islands of Trindade and Martin Vaz, with Bryopsidales being the predominant order (65% of the taxa identified), followed by Ceramiales (Villaça et al. 2006).

Among the beaches surveyed at Boipeba Island, Ponta dos Castelhanos demonstrated the greatest taxonomic richness with 109 taxa, followed by Bainema (91), Tassimirim (84), and Moreré (80) (Table 1).

The order Bryopsidales (with five families) and the




**Figure 1.** Map of Boipeba Island on the southern coast of Bahia state, Brazil, with highlighting to the beaches studied 1 - Ponta dos Castelhanos, 2 - Bainema, 3 - Moreré, 4 - Tassimirim (adapted from the map of the Tinharé-Boipeba APA and the roadmap of Bahia state).

families Caulerpaceae (eight taxa) and Cladophoraceae (seven taxa) were the richest Chlorophyta taxa at Ponta dos Castelhanos Beach. Among the Heterokontophyta/Phaeophyceae, the family Dictyotaceae (Dictyotales) stood out with 10 taxa. The most expressive order of the Rhodophyta was Ceramiales, with 23 taxa (57% of which belonged to the family Rhodomelaceae).

Bainema Beach demonstrated a flora comparable to that of Ponta dos Castelhanos, although exclusive taxa were collected at the former that had only occasionally been reported for the coast of Bahia, such as *Trichogloea*

*requienii* (Mont.) Kütz., *Parvocaulis myriosporus* (A.B.Joly & Cord.-Mar.) C.W.N.Moura & J.C.De Andrade, and *Parvocaulis parvulus* (Solms) S.Berger et al. The latter species has only recently been cited to the southern Atlantic (Moura et al. 2014a).

Among the drift specimens on Ponta dos Castelhanos Beach, we were able to confirm the presence of *Nitophyllum punctatum* for the coast of Bahia state. Torrano-Silva and Oliveira (2013) had provisionally listed this taxon for the region of the Abrolhos Archipelago (off the coast of southern Bahia), but had not encountered

**Figure 2.** Distributions of the orders of phytobenthos found at Boipeba Island, Bahia state, Brazil.

plants with spermatangial or tetrasporangial sori that could confirm their identification.

The specimens of *N. punctatum* collected at Boipeba Island had thallus blades up to 10 cm tall with delicate appearances, light-red to almost transparent, and composed of repeatedly bifurcating strips that decreased in width toward the terminal portions of the thallus, becoming wedge-shaped; thallus without a central vein, with smooth and undulating edges. Lamina monostromatic, 3–4 cells thick in the basal portion, composed of

elongated and polygonal cells, 80–100 µm × 30–50 µm, diminishing in size towards the margins and united by numerous pit-connections; chloroplasts discoid, numerous. Reproductive structures spread along the lamina, except at the base; cystocarps hemispherical, developing on both sides of the lamina, 500–900 µm in diameter, with smooth pericarp and projected ostiole; tetrasporangial sori elliptical to circular, prominent, distributed along the entire thallus on both sides of the lamina, with spherical tetrahedral tetrasporangia 100–160 µm in

**Table 1.** Phytobenthos from Boipeba Island, Bahia State, Brazil. (A) Ponta dos Castelhanos Beach, (B) Moreré Beach, (C) Bainema Beach, (D) Tassimirim Beach, (1) 2007, (2) 2012, (+) Present and (-) Absent.

Phylum/Order/Family	Species	Beach								Number in the Herbarium HUEFS
		A 1	A 2	B 1	B 2	C 1	C 2	D 1	D 2	
<b>Rhizophyta</b>										
Erythrocystidales		-	-	-	+	-	+	-	-	187829, 190752
Erythrocystidiaceae	<i>Erythrocystis carnea</i> (Dillwyn) J.Agardh	-	-	-	+	-	+	-	-	187829, 190752
<b>Corallinales</b>										
Corallinaceae	<i>Amphiroa anastomosans</i> Weber Bosse	-	+	+	-	+	+	-	+	155482, 155481, 134605, 155479
	<i>Amphiroa fragilissima</i> (L.) J.V.Lamour.	-	+	+	+	-	+	+	+	155489, 155486, 155488, 155484
	<i>Amphiroa rigida</i> J.V.Lamour.	-	+	-	-	-	+	-	-	155477, 154478
	<i>Amphiroa vanbosseae</i> Me. Lemoine	-	+	-	-	-	-	-	-	155483
	<i>Corallina panizzoi</i> Schnetter & U.Richt.	-	+	-	-	-	-	-	-	155495
	<i>Jania adhaerens</i> J.V.Lamour.	+	+	+	-	+	+	-	+	155490, 134610, 134611, 155491
	<i>Jania subulata</i> (J.Ellis & Sol.) Sond.	-	-	-	-	-	+	-	-	155494
	<i>Jania cubensis</i> Mont. ex Kütz.	-	+	-	-	-	-	-	-	155493
<b>Nemaliales</b>										
Liagoraceae	<i>Liagora ceranoides</i> J.V.Lamour.	-	-	-	-	-	-	-	+	187810
	<i>Trichogloea requienii</i> (Mont.) Kütz.	-	-	-	-	-	+	-	-	190766
Galaxauraceae	<i>Dichotomaria marginata</i> (J.Ellis & Sol.) Lam.	-	+	+	-	+	-	-	-	187808, 130912, 130913
	<i>Dichotomaria obtusata</i> (J.Ellis & Sol.) Lam.	+	-	-	+	-	-	+	+	130911, 187851, 130914
	<i>Galaxaura rugosa</i> (J.Ellis & Sol.) J.V.Lamour.	-	+	+	+	-	+	-	+	187849, 134575, 187848, 187850
	<i>Tricleocarpa cylindrica</i> (J.Ellis & Sol.) Huisman & Borow	-	-	+	-	-	+	+	+	130903, 187836, 130904
	<i>Tricleocarpa fragilis</i> (L.) Huisman & R.A.Towns.	-	-	+	-	+	-	-	-	130906, 130907
<b>Bonnemaisoniales</b>										
Bonnemaisoniaceae	<i>Asparagopsis taxiformis</i> (Delile) Trevis.	-	-	+	-	-	-	+	-	134607, 190771

*Continued*

**Table 1.** Continued.

Phylum/Order/Family	Species	Beach								Number in the Herbarium HUEFS
		A 1	A 2	B 1	B 2	C 1	C 2	D 1	D 2	
<b>Ceramiales</b>										
Callithamniaceae	<i>Crouania attenuata</i> (C.Agardh) J.Agardh	-	+	-	-	-	-	-	-	187797
Ceramiaceae	<i>Centroceras gasparrinii</i> (Menegh.) Kütz.	+	-	+	-	-	-	+	-	155497, 134576, 134577,
Dasyaceae	<i>Dasya rigidula</i> (Kütz.) Ardis.	+	-	-	-	-	-	-	-	134572
	<i>Heterosiphonia crispella</i> (C.Agardh) M.J.Wynne	-	+	-	-	-	-	-	-	190737
Delesseriaceae	<i>Caloglossa leprieurii</i> (Mont.) G.Martens	+	-	-	-	-	-	+	-	187787, 189919
	<i>Nitophyllum punctatum</i> (Stackh. in With.) Grev.	+	-	-	-	-	-	-	-	155514
Rhodomelaceae	<i>Acanthophora muscooides</i> (L.) Bory	-	+	-	-	+	+	-	-	190741, 134578
	<i>Acanthophora spicifera</i> (Vahl) Børgesen	+	+	-	+	-	+	-	+	134595, 187796, 187843, 190740
	<i>Bostrychia montagnei</i> Harv.	+	-	-	-	+	+	+	-	189914, 190755, 164817
	<i>Bostrychia moritziana</i> (Sond. ex Kütz.) J.Agardh	-	-	-	-	-	+	-	-	187815
	<i>Bostrychia radicans</i> (Mont.) Mont. in Orb.	+	-	-	-	-	+	-	-	189912, 187792
	<i>Bostrychia tenella</i> (J.V.Lamour.) J.Agardh	+	-	-	-	+	+	+	-	189911, 189909, 164816
	<i>Bryocladia cuspidata</i> (J.Agardh) De Toni	-	-	-	-	-	-	+	-	146371
	<i>Bryothamnion seaforthii</i> (Turner) Kütz.	-	+	-	-	-	-	+	-	187789, 147326
	<i>Chondria collinsiana</i> M.Howe	-	+	-	-	-	-	-	-	155509
	<i>Digenea simplex</i> (Wulfen) C.Agardh	-	+	-	+	+	+	+	-	190742, 187834, 147281, 187818
	<i>Herposiphonia tenella</i> (C.Agardh) Ambronn	-	-	-	-	+	-	-	-	155511
	<i>Laurencia</i> sp.	-	-	-	-	+	-	-	-	134604
	<i>Laurencia dendroidea</i> J.Agardh	+	+	+	+	+	+	+	+	187676, 187680, 187681, 187682
	<i>Murrayella periclados</i> (C.Agardh) F.Schmitz	+	-	-	-	-	-	+	-	189910, 164808
	<i>Neosiphonia gorgoniae</i> (Harv.) S.M.Guim. & M.T.Fujii	+	-	-	-	-	-	-	-	134574
	<i>Palisada perforata</i> (Bory) K.W.Nam	-	+	-	+	-	+	-	+	187814, 187811, -187801, 187816
	<i>Vidalia obtusiloba</i> J.V.Lamour.	+	+	-	-	-	-	+	-	134557, 134558
Spyridiaceae	<i>Spyridia clavata</i> Kütz.	-	+	-	-	-	-	-	-	190747
	<i>Spyridia filamentosa</i> (Wulfen) Harv. in Hook.	-	+	-	-	-	+	-	-	190735, 190739
	<i>Spyridia hypnoides</i> (Bory in Belanger) Papenf.	-	-	-	-	+	-	-	-	136240
Wrangeliaceae	<i>Griffithsia schousboei</i> Mont. in Webb. var. <i>schousboei</i>	-	-	-	-	-	+	-	-	155510
	<i>Griffithsia schousboei</i> var. <i>anastomosans</i> E.C.Oliveira	-	+	-	-	-	-	-	-	190769
	<i>Haloplegma duperreyi</i> Mont.	-	-	-	-	+	-	+	-	134573, 134609
	<i>Tiffaniella gorgoniae</i> (Mont.) Doty & Meñez	-	+	-	-	-	+	-	+	133789, 133791, 133794
<b>Gelidiales</b>										
Gelidiaceae	<i>Gelidium</i> sp.	-	-	-	-	+	-	-	-	190792
Gelidiellaceae	<i>Gelidiella acerosa</i> (Forssk.) Feldmann & Hamel	-	+	-	+	+	+	+	-	187798, 190746, 134603, 187831
<b>Gigartinales</b>										
Cystocloniaceae	<i>Hypnea cornuta</i> (Kütz.) J.Agardh	-	-	-	+	-	-	-	-	187845
	<i>Hypnea musciformis</i> (Wulfen in Jacq.) J.V.Lamour.	+	+	+	+	-	+	+	+	136246, 136245, 187794, 136241
	<i>Hypnea nigrescens</i> Grev. ex J.Agardh	-	-	-	+	-	-	-	-	187844
	<i>Hypnea spinella</i> (C.Agardh) Kütz.	+	+	+	-	-	-	-	-	136242, 136244
Gigartinaceae	<i>Chondracanthus acicularis</i> (Roth) Frederiq in Hommersand et al.	-	-	-	-	+	-	-	-	190770
Rhizophyllidaceae	<i>Ochtodes secundiramea</i> (Mont.) M.Howe	-	+	+	+	+	-	+	+	187795, 134532, 134531, 134530
Solieriaceae	<i>Agardhiella ramossissima</i> (Harv.) Kylin	+	+	-	-	-	-	-	-	134556
	<i>Agardhiella subulata</i> (C.Agardh) Kraft & M.J.Wynne	+	-	+	-	-	-	-	-	134554, 134555
	<i>Meristotheca gelidium</i> (J.Agardh) E.J.Faye & Masuda in E.J.Faye et al.	+	-	+	-	+	+	-	-	130910, 130909, 187800
<b>Gracilariales</b>										
Gracilariaeae	<i>Gracilaria birdiae</i> Plastino & E.C.Oliveira	-	-	+	-	+	+	+	+	133777, 133776, 133775
	<i>Gracilaria cervicornis</i> (Turner) J.Agardh	+	+	+	+	+	+	+	+	190758, 133780, 133779, 133778
	<i>Gracilaria cuneata</i> Aresch.	+	+	+	-	+	+	-	-	133773, 133774, 133772
	<i>Gracilaria curtissiae</i> J.Agardh	+	+	+	-	-	+	+	+	133768, 133766, 187805, 133767
	<i>Gracilaria domingensis</i> (Kütz.) Sond. ex Dickie	+	-	+	-	+	+	+	+	133762, 133759, 133761, 133763
	<i>Gracilaria isabellana</i> Gurgel, Frederiq & J.N.Norris	+	-	+	+	+	+	+	+	133770, 133764, 133771, 133769,
	<i>Gracilaria</i> sp.	+	-	-	-	+	-	+	-	133781, 133783, 133782
	<i>Hydropuntia caudata</i> (J.Agardh) Gurgel & Frederiq	+	+	+	+	+	+	+	+	190744, 187788, 187839, 187840
	<i>Hydropuntia cornea</i> (J.Agardh) M.J.Wynne	+	+	+	+	+	+	+	+	134585, 134586, 134583, 134584
<b>Halymeniales</b>										
Halymeniaceae	<i>Cryptonemia crenulata</i> (J.Agardh) J.Agardh	+	-	-	-	-	-	+	+	136238, 136239
	<i>Cryptonemia limensis</i> (Kütz.) J.A.Lewis	-	-	-	-	-	-	+	-	136237

*Continued*


**Table 1.** Continued.

Phylum/Order/Family	Species	Beach								Number in the Herbarium HUEFS
		A 1 2	B 1 2	C 1 2	D 1 2					
	<i>Cryptonemia seminervis</i> (C.Agardh) J.Agardh	-	-	-	+	-	+	+	-	190745, 136236
	<i>Halymenia duchassaingii</i> (J.Agardh) Kylin	-	+	-	-	-	+	-	-	190757, 190753
	<i>Halymenia elongata</i> C.Agardh	-	+	-	-	-	-	-	-	187833
	<i>Halymenia floressi</i> (Clemente) C.Agardh	-	+	-	-	-	-	-	-	187803
<b>Sebdeniales</b>										
Sebdeniaceae	<i>Sebdenia flabellata</i> (J.Agardh) P.G.Parkinson	+	-	-	-	-	-	-	-	134580
<b>Peyssonneliales</b>										
Peyssonneliaceae	<i>Peyssonnelia inamoena</i> Pilg.	-	-	+	-	-	-	-	-	155496
<b>Rhodymeniales</b>										
Rhodymeniaceae	<i>Botryocladia occidentalis</i> (Børgesen) Kylin	+	+	+	+	-	+	+	-	134535, 134537, 187821, 134534
	<i>Botryocladia pyriformis</i> (Børgesen) Kylin	-	-	+	-	-	+	-	-	134536, 187785
Champiaceae	<i>Champia parvula</i> (C.Agardh) Harv.	-	+	-	+	-	+	-	+	187819, 190774, 187835, 187804
	<i>Champia feldmanii</i> Diaz-Pif.	+	-	-	+	-	-	-	-	187820, 190775
	<i>Champia vieillardii</i> Kütz.	-	-	-	-	-	-	+	-	136233
Lomentariaceae	<i>Ceratodictyon planicaule</i> (W.R.Taylor) M.J.Wynne	-	-	-	-	+	-	-	-	134553
	<i>Ceratodictyon repens</i> (Kütz.) R.E.Norris	-	-	+	-	+	-	-	-	148274, 148275
	<i>Ceratodictyon variabile</i> (Grev. ex J.Agardh) R.E.Norris	-	-	-	-	-	-	+	-	134548
<b>Heterokontophyta</b>										
<b>Dictyotales</b>										
Dictyotaceae	<i>Canistrocarpus cervicornis</i> (Kütz.) De Paula & De Clerck	+	+	+	+	+	+	+	+	134518, 134515, 134517, 133817
	<i>Canistrocarpus crispatus</i> (J.V.Lamour.) De Paula & De Clerck	-	-	-	+	-	-	-	-	164805
	<i>Dictyopteris delicatula</i> J.V.Lamour.	+	+	+	+	-	+	+	+	133785, 133786, 190754, 133784
	<i>Dictyopteris jamaicensis</i> W.R.Taylor	+	+	-	-	-	-	-	-	133788
	<i>Dictyopteris plagiogramma</i> (Mont.) Vickers	+	-	-	-	-	-	-	-	133787
	<i>Dictyota bartayresiana</i> J.V.Lamour.	-	-	-	-	-	-	-	+	189939
	<i>Dictyota ciliolata</i> Sond. ex Kütz.	+	+	+	+	+	+	+	+	134518, 134523, 134525, 134524
	<i>Dictyota crenulata</i> C.Agardh	-	-	+	-	+	+	+	+	134519, 134521, 134520
	<i>Dictyota menstrualis</i> (Hoyt) Schnetter, Hörnig & Weber-Peukert	+	-	-	-	+	-	+	-	134529, 134527, 134526
	<i>Dictyota mertensii</i> (Mart.) Kütz.	+	+	-	+	-	+	+	-	134529, 189928, 134529, 134528
	<i>Lobophora variegata</i> (J.V.Lamour.) Womersley ex E.C. Oliveira	+	+	+	+	+	+	+	+	133756, 133757, 133755, 133758
	<i>Padina antillarum</i> (Kütz.) Picc.	-	-	+	-	-	-	+	-	133717, 130916
	<i>Padina boergesenii</i> Allender & Kraft	+	+	+	+	+	-	+	+	108746, 130920, 130917, 130919,
	<i>Padina sanctae-crucis</i> Børgesen	-	-	-	-	-	+	+	-	187922, 130918
	<i>Spatoglossum schroederi</i> (C.Agardh) Kütz.	+	-	+	-	+	+	+	-	134613, 136234, 134612, 134614
<b>Sphaereliales</b>										
Sphaereliaceae	<i>Sphaerelaria rigidula</i> Kütz.	-	+	-	-	-	-	-	-	155498
<b>Fucales</b>										
Sargassaceae	<i>Sargassum polyceratum</i> Mont.	-	-	-	-	+	-	-	-	148293
	<i>Sargassum vulgare</i> C.Agardh	+	-	-	-	-	-	-	-	148292
	<i>Sargassum</i> sp.	-	-	+	-	-	-	-	+	147578, 148278
<b>Ectocarpales</b>										
Acinetosporaceae	<i>Feldmannia mitchelliae</i> (Harvey) H.-S.Kim	-	-	-	-	-	-	+	-	134579
Scytoniphonaceae	<i>Colpomenia sinuosa</i> (Roth) Derbès & Solier	+	+	+	+	-	+	+	-	134568, 134569, 187915, 134570
<b>Chlorophyta</b>										
<b>Ulvales</b>										
Gayraliaceae	<i>Gayralia brasiliensis</i> Pellizzari, M.C.Oliveira & N.S.Yokoia	-	-	-	-	-	+	+	-	190767, 134608
Ulvaceae	<i>Ulva chaetomorphoides</i> (Børgesen) H.S.Hayden et al.	-	+	-	-	-	-	-	-	190773
	<i>Ulva fasciata</i> Delile	-	-	-	-	-	-	+	-	133754
	<i>Ulva flexuosa</i> Wulfen	+	-	+	-	+	+	+	-	133748, 133747, 133746, 133749
	<i>Ulva flexuosa</i> Wulfen subsp. <i>paradoxa</i> (C.Agardh) M.J.Wynne	-	-	-	-	-	+	-	-	187860
	<i>Ulva lactuca</i> L.	+	+	+	+	+	+	+	-	133752, 133753, 133751, 133750
<b>Cladophorales</b>										
Anadyomenaceae	<i>Anadyomene stellata</i> (Wulfen in Jacq.) C.Agardh	+	+	+	+	+	-	+	-	133815, 133814, 133813, 133812
Cladophoraceae	<i>Chaetomorpha aerea</i> (Dillwyn) Kütz.	-	+	-	-	-	-	-	-	155512
	<i>Chaetomorpha antennina</i> (Bory) Kütz.	-	+	-	-	-	-	-	-	187903
	<i>Chaetomorpha brachygona</i> Harv.	+	-	-	+	-	-	-	-	164811, 190785

*Continued*

**Table 1.** Continued.

Phylum/Order/Family	Species	Beach								Number in the Herbarium HUEFS
		A 1 2	B 1 2	C 1 2	D 1 2					
	<i>Cladophora brasiliiana</i> G.Martens	-	-	-	+	-	-	-	-	190780
	<i>Cladophora corallicola</i> Børgesen	+	-	-	-	-	-	-	+	136424, 136423
	<i>Cladophora dalmatica</i> Kütz.	+	+	+	+	-	-	-	-	136476, 147505
	<i>Cladophora laetevirens</i> (Dillwyn) Kütz.	+	+	-	-	+	-	-	-	136499, 136498
	<i>Cladophora socialis</i> Kütz.	-	-	-	+	-	-	-	-	190790
	<i>Cladophora vagabunda</i> (L.) C.Hoek	+	-	+	-	+	+	-	-	136551, 136550, 136549
	<i>Rhizoclonium africanum</i> Kütz.	-	-	-	-	-	-	+	-	136299
	<i>Rhizoclonium riparium</i> (Roth) Kütz. ex Harv.	+	-	-	-	+	-	-	-	164803, 155500
<b>Siphonocladales</b>										
Bodaleaceae	<i>Cladophoropsis membranacea</i> (H. Bang ex C.Agardh) Børgesen	+	-	-	+	-	-	+	-	136612, 190791, 136613
Siphonocladaceae	<i>Phyllodictyon anastomosans</i> (Harv.) Kraft & M.J.Wynne	-	+	+	-	+	-	+	-	155499, 136327, 136326, 136325
Valoniaceae	<i>Dictyosphaeria versluysi</i> Weber Bosse	+	+	+	+	+	+	+	+	134588, 136365, 134589, 134591
	<i>Valonia aegagropila</i> C.Agardh	+	+	+	+	+	-	+	-	134606, 134596, 130852, 136638
<b>Bryopsidales</b>										
Bryopsidaceae	<i>Bryopsis hypnoides</i> J.V.Lamour.	-	+	-	+	-	-	-	+	187852, 187854, 187855
	<i>Bryopsis pennata</i> J.V.Lamour.	+	-	+	-	+	-	+	+	134546, 134541, 134544, 134542
Codiaceae	<i>Bryopsis plumosa</i> (Huds.) C.Agardh	+	-	+	-	+	-	+	-	134546, 134543, 134544, 134539
	<i>Codium intertextum</i> Collins & Herv.	+	-	+	+	+	+	+	-	133793, 133794, 133792, 133795
	<i>Codium isthmocladum</i> Vickers	+	+	+	-	+	+	+	+	133790, 133794, 133791, 133789
	<i>Codium taylorii</i> P.C.Silva	-	-	-	+	-	-	-	-	190788
Caulerpaceae	<i>Caulerpa cupressoides</i> (H.West in Vahl) C.Agardh	-	-	-	-	-	-	+	-	133718
	<i>Caulerpa cupressoides</i> (H.West in Vahl) C.Agardh var. <i>lycopodium</i>	+	-	+	-	+	-	+	-	133730, 133723, 133724, 133728
	<i>Caulerpa fastigiata</i> Mont.	-	+	-	-	-	-	+	-	187872, 133731
	<i>Caulerpa kempfi</i> A.B.Joly & S.Pereira	-	+	-	-	-	+	-	-	187875, 187876
	<i>Caulerpa macrophysa</i> Sond. ex Kütz.	-	+	+	-	+	-	+	-	187867, 133732, 133734, 133733
	<i>Caulerpa prolifera</i> (Forssk.) J.V.Lamour.	-	-	-	-	-	-	+	-	133740
	<i>Caulerpa racemosa</i> (Forssk.) J.Agardh	+	-	+	-	+	-	-	-	133720, 133721, 133722
	<i>Caulerpa racemosa</i> (Forssk.) J.Agardh var. <i>occidentalis</i> (J.Agardh) Børgesen	+	+	-	+	-	+	+	+	133735, 133736,
	<i>Caulerpa scalpelliformis</i> (R.Br. ex Turner) C.Agardh	+	+	+	+	+	+	+	+	133739, 133743, 133738, 133737
	<i>Caulerpa sertularioides</i> (S.G.Gmel.) M.Howe	+	+	+	+	+	+	+	+	133742, 133743, 133744, 133741
Halimedaceae	<i>Caulerpa verticillata</i> J.Agardh	-	-	+	-	-	-	-	-	190768
	<i>Halimeda cuneata</i> K.Hering	+	+	+	+	+	+	+	-	134564, 134563, 134567, 134565
	<i>Halimeda opuntia</i> (L.) J.V.Lamour.	-	+	+	+	+	+	+	+	187888, 134560, 134559, 134562
Udoteaceae	<i>Boodleopsis pusilla</i> (Collins) W.R.Taylor, A.B.Joly & Bernat.	+	-	-	-	-	-	-	-	164814
	<i>Penicillus capitatus</i> Lam.	+	+	+	+	+	+	+	+	134551, 134552, 134547, 134550
	<i>Udotea cyathiformis</i> Decne var. <i>cyathiformis</i> f. <i>sublittoralis</i> (W.R.Taylor) D.S.Littler & Littler	-	+	-	-	+	-	-	-	187899, 134593
	<i>Udotea flabellum</i> (J.Ellis & Sol.) J.V.Lamour.	+	+	-	-	-	-	+	-	134594, 134592
<b>Dasycladales</b>										
Dasycladaceae	<i>Neomeris annulata</i> Dickie	+	+	+	+	+	+	+	+	134599, 134601, 134600, 134602
Polyphysaceae	<i>Acetabularia calyculus</i> J.V.Lamour. in Quoy & Gaimard	-	+	-	-	-	-	-	-	155504
	<i>Acetabularia crenulata</i> J.V.Lamour.	+	+	-	+	-	-	-	-	134582, 187889
	<i>Acetabularia schenckii</i> K.Möbius	+	+	-	-	-	-	-	-	134581
	<i>Parvocaulis myriosporus</i> (A.B.Joly & Cord.-Mar.) C.W.N.Moura & J.C. De Andrade	-	-	-	-	-	+	-	-	155503
	<i>Parvocaulis parvulus</i> (Solms) S.Berger et al.	-	-	-	-	-	+	-	-	155501
	<i>Parvocaulis pusillus</i> (M.Howe) S.Berger et al.	+	+	-	-	+	-	-	-	130897, 130896
<b>Cyanobacteria</b>										
<b>Nostocales</b>										
Sympytonemataceae	<i>Bachytrichia quoy</i> (C.Agardh) Bornetii & Flahault	+	-	-	-	-	-	-	-	189944
Oscillatoriales										
Oscillatoriaceae	<i>Oscillatoria</i> sp.	-	-	-	-	+	-	-	-	155508
<b>Magnoliophyta</b>										
<b>Alismatales</b>										
Cymodoceaceae	<i>Halodule wrightii</i> Asch.	-	-	-	-	+	-	-	-	190772



**Figure 3.** *Nithophyllum pannosum* (Stackh.) Grev. **A.** Tetrasporic plant. **B.** Detail of the sori of the tetrasporangia distributed on the bifurcated lamina; note the undulating margin. **C.** Female plant. **D.** Cystocarps. **E.** Detail of a cystocarp, with ostiole prominent in lateral view. **F.** Surface view of thallus cells with pit-connections. **G.** Margin of the smooth lamina. **H.** Transversal section of the monostromatic lamina. **I.** Tetrasporangial sorus in surface view. **J.** Transversal section of a tetrasporangial sorus. **K.** Detail of the tetrahedral tetrasporangia.

**Table 2.** Comparisons of the vegetative and reproductive structures of *Nitophyllum punctatum*, *N. adhaerens*, *N. wilsoniae*, and *Haraldiophyllum bonnemaisonii*. ND = no data available

Characteristics	<i>Nitophyllum punctatum</i> (Stack. in With.) Grev.	<i>Nitophyllum adhaerens</i> M.J.Wynne	<i>Nitophyllum wilsoniae</i> Collins & Herv.	<i>Haraldiophyllum bonnemaisonii</i> (Kylin) Zinova
Thalli length	3-50 cm	up to 3-4 cm	up to 10 cm	2-15 cm
Branching	irregular to dichotomous lobed	irregular; deeply lobed	di- to polychotomous	more-or-less regularly dichotomously divided into overlapping lobes
Midrib-like	absent	absent	absent	present; one or more development basally and extending upwards on blade
Blade margins	smooth, slightly undulate	smooth, slightly undulate; presence of peripheral rhizoids uni- or multicellular	small teeth along the margin, undulate	smooth, slightly undulate
Cystocarps (diam.)	(500-) 650-900 µm	650-860 µm	ND	430-600 µm
Ostiole	projecting	projecting	ND	non-projecting
Tetrasporangial sorus distribution	scattered on blade	in terminal lobes	scattered on blade	scattered, over upper two-thirds of blade
Tetrasporangia (diam.)	100-175 µm	38-75 µm	45-60 (-75) µm	50-70 µm
Type locality	Weymouth, Dorset, England	Bocana Reef, Estacion Puerto Morelos, Quintana Roo, Mexico	Bermuda	Orkney Island, Scotland
References	Maggs and Hommersand 1993, This work	Wynne 1997, Surati and Guimarães 2007	Dawes and Mathieson 2008, Joly et al. 1965, Littler and Littler 2000	Maggs and Hommersand 1993

diameter (Figure 3). The material examined demonstrated characteristics similar to those described by Maggs and Hommersand (1993) for the British Isles and by Torrano-Silva and Oliveira (2013) for the Abrolhos Archipelago.

Two other species of *Nitophyllum* Grev. have been cited in literature (Moura et al. 2014b) for the Brazilian coast: *N. wilsoniae* Collins & Herv. (reported for the coasts of Bahia, Espírito Santo, and São Paulo) and *N. adhaerens* M.J.Wynne (reported for Espírito Santo, Rio de Janeiro, São Paulo, and Santa Catarina). The characteristics that separate the species are listed in Table 2.

According to Maggs and Hommersand (1993), *N. punctatum* is morphologically similar to *Haraldiophyllum bonnemaisonii* (Kylin) Zinova, although the latter differs by presence of a midrib, and has a smaller cystocarp, without a projected ostiole, and smaller tetrasporangia, approximately 50–70 µm in diameter (Table 2).

The present work added 150 taxa to marine flora the Boipeba Island, including *N. punctatum*, and confirmed its occurrence along the Bahian coast.

This survey of the phytobenthic biodiversity of Boipeba Island was undertaken to increase our knowledge of the insular marine flora and reef communities that compose the rich natural ecosystems along the coast of Bahia state, Brazil.

## ACKNOWLEDGEMENTS

The authors are grateful to the Universidade Estadual de Feira de Santana for logistic support, as well as the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)/ Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) through the Projeto Flora da Bahia (483909/2012).

## LITERATURE CITED

- Almeida, W.R., A.M. Alves, S.M.P.B. Guimarães and C.W.N. Moura. 2012. Cladophorales and Siphonocladales (Chlorophyta) from Bimbarras Island, Todos os Santos Bay, Bahia state, Brazil. Iheringia. Série Botânica 67(2): 149–164. [http://www.fzb.rs.gov.br/upload/2014032809594ih67\\_2\\_p149\\_164.pdf](http://www.fzb.rs.gov.br/upload/2014032809594ih67_2_p149_164.pdf)
- Almeida, W.R., S.M.P.B. Guimarães and C.W.N. Moura. 2013. *Bangiopsis subsimplex* (Mont.) F. Schmitz (Stylonematiales, Rhodophyta) on the northeastern coast of Brazil. Acta Botanica Brasilica 27(1): 231–236. doi: [10.1590/S0102-33062013000100022](https://doi.org/10.1590/S0102-33062013000100022)
- Almeida, W.R., S.M.P.B. Guimarães and C.W.N. Moura. 2014. Novas adições à flora marinha bentônica da costa nordeste do Brasil. Iheringia. Série Botânica 69(1): 97–105. [http://www.fzb.rs.gov.br/upload/20140805153146ih69\\_1\\_p97\\_105.pdf](http://www.fzb.rs.gov.br/upload/20140805153146ih69_1_p97_105.pdf)
- Alves, A.M., C.W.N. Moura, G.L. Alves and L.M.S. Gustinari. 2009. Os gêneros *Chaetomorpha* Kütz. nom. cons. e *Rhizoclonium* Kütz. (Chlorophyta) do litoral do Estado da Bahia, Brasil. Revista Brasileira de Botânica 32(3): 545–570. doi: [10.1590/S0100-84042009000300014](https://doi.org/10.1590/S0100-84042009000300014)
- Alves, A.M., L.M.S. Gustinari and C.W.N. Moura. 2010. La familia Valoniaceae (Chlorophyta) en el estado de Bahía, Brasil: aspectos morfológicos y de distribución. Hidrobiología 20(2): 171–184. <http://www.scielo.org.mx/pdf/hbio/v20n2/v20n2a8.pdf>
- Alves, A.M., L.M.S. Gustinari and C.W.N. Moura. 2011a. Morphology and taxonomy of *Anadyomene* species (Cladophorales, Chlorophyta) from Bahia, Brazil. Botanica Marina 54: 135–145. doi: [10.1515/bot.2011.015](https://doi.org/10.1515/bot.2011.015)
- Alves, A.M., L.M.S. Gustinari and C.W.N. Moura. 2011b. *Microdictyon* (Chlorophyta, Anadyomenaceae) do Estado da Bahia, Brasil. Sitientibus 11(1): 57–61. <http://pkp.ufes.br/ojs/index.php/sitientibusBiologia/article/download/88/86>
- Alves, A.M., L.M.S. Gustinari and C.W.N. Moura. 2012a. Flora of Bahia: Siphonocladaeae. Sitientibus. Série Ciências Biológicas 12(2): 167–177. doi: [10.13102/scb126](https://doi.org/10.13102/scb126)
- Alves, A.M., L.M.S. Gustinari and C.W.N. Moura. 2012b. Flora of Bahia: Boodleaceae. Sitientibus. Série Ciências Biológicas 12(2): 179–188. doi: [10.13102/scb128](https://doi.org/10.13102/scb128)
- Alves, A.M., L.M.S. Gustinari, I.S. Oliveira, K.L. Moniz-Brito and

- C.W.N. Moura. 2012c. The genus *Cladophora* (Chlorophyta) in the littoral of Bahia, Brazil. *Nova Hedwigia* 95(3-4): 337–372. doi: [10.1127/0029-5035/2012/0025](https://doi.org/10.1127/0029-5035/2012/0025)
- Bahia. 1999. Coordenação de Desenvolvimento do Turismo. Programa de desenvolvimento turístico da Bahia — oportunidades de investimento. Governo do Estado da Bahia, Salvador, Brasil, 67 pp.
- Bicudo, C.E.M. and M. Menezes. 2010. Introdução: as algas do Brasil; pp. 49–60, in: R.C. Forzza et al. (org.). Catálogo de plantas e fungos do Brasil. Rio de Janeiro: Instituto de Pesquisas Jardim Botânico do Rio. <http://books.scielo.org/id/z3529/pdf/forzza-9788560035083-06.pdf>
- Brummitt, R.K. and C.E. Powell. 1992. Authors of plant names. A list of authors of scientific names of plants, with recommended standard forms of these names including abbreviations. Kew: Royal Botanic Gardens. 732 pp.
- Caires, T.A. 2013. Cianobactérias marinhas bentônicas filamentosas do litoral do Estado da Bahia (MSC. thesis). Universidade Estadual de Feira de Santana. Feira de Santana. 140 pp.
- Creed, M., M.T. Fujii, M.B.B. Barreto, S.M.P.B. Guimarães, V. Cassano, S.M.B. Pereira, M.F.O. Carvalho and S. Khader. 2010. Rhodophyceae. pp. 416–436, in: R.C. Forzza et al. (org.). Catálogo de plantas e fungos do Brasil. Vol. 1. Rio de Janeiro: Instituto de Pesquisas Jardim Botânico do Rio de Janeiro. <http://books.scielo.org/id/z3529/12>
- Dawes, C.J. and A.C. Mathieson. 2008. The seaweeds of Florida. Gainesville, Florida: University Press of Florida. 591 pp.
- Dutra, L.X.C. and R.K.P. Kikuchi. 2003. Método do vídeo transecto para diagnóstico de recifes de corais na Baía de Todos os Santos, Bahia. Proceedings VI Congresso de Ecologia do Brasil 1: 341–343.
- Ginsburg, R.N., P.A. Kramer, J.C. Lang, P. Sale and R.S. Steneck. 1998. Atlantic and Gulf Reef Assessment (AGRRA) - REVISED Rapid Assessment Protocol (RAP). Accessed at <http://www.agrra.org/>, 6 June 2014.
- Jesus, P.B. 2012. O gênero *Hypnea* J.V. Lamour. (Gigartinales, Rhodophyta) no litoral do estado da Bahia, Brasil: aspectos morfológicos e anatômicos (MSC. thesis). Universidade Estadual de Feira de Santana. Feira de Santana. 185 pp.
- Joly, A. B., M.C. Marino, N. Yamaquishi-Tomita, Y. Ugadim, E.C. Oliveira Filho and M.M. Ferreira. 1965. Additions to the marine flora of Brazil V. Arquivos da Estação de Biologia Marinha da Universidade Federal do Ceará 5(1): 65–78.
- Kain, J.M and T.A. Norton. 1990. Marine Ecology; pp. 377–422, in: K.M. Cole and R.G. Sheath (eds.). Biology of the red algae. New York: Cambridge University Press.
- Kikuchi, R.K.P., M.D.M. Oliveira, Z.M.A.N. Leão, R.M. Silva and P.M.R. Martins. 2008. Os recifes de Tinhare-Boipeba-Camamu, Bahia. Proceedings of Rio Oil and Gas Expo and Conference 1: 1–8.
- Kim, H.-S. 2010. Heterokontophyta: Phaeophyceae: Ectocarpales. Marine brown algae I; pp. 3–137. in: H.-S. Kim and S.-M. Boo (eds). Algal flora of Korea. Volume 2, Number 1. Ectocarpaceae, Acinetopsoraceae, Chordariaceae. Incheon: National Institute of Biological Resources.
- Littler, D.S. and M.M. littler, 2000. Caribbean reef plants. An identification guide to the reef plants of the Caribbean, Bahamas, Florida and Gulf of Mexico. Washington, D.C.: OffShore Graphics. 542 pp.
- Maggs, C.A. and M.D. Hommersand. 1993. Seaweeds of the British Isles. Vol. 1. Rhodophyta, Part 3A Ceramiales. London: Natural History Museum Publication. 444 pp.
- Marins, B.V., P.A. Brasileiro, M.B.B. Barreto, J.M.C. Nunes, Y. Yoneshigue-Valentin and G.M. Amado Filho. 2008. Subtidal benthic marine algae of the Todos os Santos Bay, Bahia state, Brazil. *Oecologia Brasiliensis* 12: 229–242. <http://dialnet.unirioja.es/descarga/articulo/2881794.pdf>
- Moura, C.W.N. 2010. Ulvophyceae. pp. 438–448, in: R.C. Forzza et al. (org.). Catálogo de plantas e fungos do Brasil. Vol. 1. Rio de Janeiro: Instituto de Pesquisas Jardim Botânico do Rio de Janeiro. <http://books.scielo.org/id/z3529/12>
- Moura, C.W.N., W.R. Almeida, A.A. Santos, J.C. Andrade Junior, A.M. Alves and K.L. Moniz-Brito. 2014a. Polyphysaceae (Dasycladales, Chlorophyta) in Todos os Santos Bay, Bahia, Brazil. *Acta Botanica Brasilica* 28(2): 147–164. doi: [10.1590/S0102-33062014000200002](https://doi.org/10.1590/S0102-33062014000200002)
- Moura, C.W.N., J.M.C. Nunes, M.T. Fujii, M.B.B. Barros-Barreto, O. Necchi Jr., S.M.P.B Guimarães, S.M.B. Pereira, M.F. Oliveira-Carvalho, M.C. Oliveira, M.A.O. Figueiredo, R.G. Bahia, G.M. Amado Filho, G.M. Lyra, P.B. Jesus, A.A. Santos, G.N. Santos, I.O. Costa, M.C. Henriques, S. Khader and V. Cassano. 2014b. Rhodophyceae. Lista de espécies da flora ---do Brasil. Jardim Botânico do Rio de Janeiro. Accessed at <http://floradobrasil.jbrj.gov.br/jabot/floradobrasil/FB99625>, 28 May 2014.
- Nunes, J.M.C. 1998. Catálogo de algas marinhas bentônicas do Estado da Bahia, Brasil. *Acta Botanica Malacitana* 23: 5–21. [http://www.biolveg.uma.es/abm/Volumenes/vol23/23\\_Nunes.pdf](http://www.biolveg.uma.es/abm/Volumenes/vol23/23_Nunes.pdf)
- Nunes, J.M.C. 2005. Rodofíceas marinhas bentônicas do Estado da Bahia, Brasil (PhD thesis). São Paulo: Instituto de Biociências da Universidade de São Paulo. 410 pp.
- Nunes, J.M.C. and S.M.P.B. Guimarães. 2008. Novas referências de rodofíceas marinhas bentônicas para o litoral brasileiro. *Biota Neotropica* 8: 89–100. <http://www.scielo.br/pdf/bn/v8n4/ao8v8n4.pdf>
- Oliveira, E.C., P.A. Horta, C.E. Amancio and C.L. Sant'Anna. 1999. Algas e angiospermas marinhas bênticas do litoral brasileiro: diversidade, exploração e conservação. Brasília: Agência Nacional do Petróleo, Gás Natural e Biocombustíveis. Accessed at [http://www.brasil-rounds.gov.br/round5/arquivos\\_r5/guias/perfuracao/5round/refere/plantas\\_marinhas.pdf](http://www.brasil-rounds.gov.br/round5/arquivos_r5/guias/perfuracao/5round/refere/plantas_marinhas.pdf), 28 May 2014.
- Oliveira Filho, E.C. and Y. Ugadim. 1974. New references of benthic marine algae to Brazilian flora. *Boletim Botânica da Universidade de São Paulo* 2: 71–91. <http://www.revistas.usp.br/bolbot/article/download/57616/60671>
- Oliveira Filho, E.C. and Y. Ugadim. 1976. A survey of the marine algae of Atol das Rocas (Brazil). *Phycologia* 15(1): 41–44. doi: [10.2216/i0031-8884-15-1-41.1](https://doi.org/10.2216/i0031-8884-15-1-41.1)
- Pereira, S.M.B. 2002. Desenvolvimento e situação atual do conhecimento das macroalgas marinhas das regiões Nordeste e Norte, pp. 117–121, in: E.L. Araujo, A.N. Moura, E.S.B. Sampaio, L.M.S. Gestinari and J.M.T. Carneiro (eds.). Biodiversidade, conservação e uso sustentável da flora do Brasil. Recife: Brasil/ Imprensa Universitária.
- Santos, A.A. and C.W.N. Moura. 2011. Additions to the epiphytic macroalgae flora of Bahia and Brazil. *Phytotaxa* 28: 53–64.
- Széchy, M.T.M. and J.C. Paula. 2010. Phaeophyceae. pp. 404–408, in: R.C. Forzza et al. (org.). Catálogo de plantas e fungos do Brasil. Vol. 1. Rio de Janeiro: Instituto de Pesquisas Jardim Botânico do Rio de Janeiro. <http://books.scielo.org/id/z3529/12>
- Surati, R. and S.M.P.B. Guimarães. 2007. *Nitophyllum adhaerens* (Ceramiales, Delesseriaceae), a new occurrence in the Brazilian coast. *Hoehnea* 37(3): 335–340. doi: [10.1590/S2236-89062007000300004](https://doi.org/10.1590/S2236-89062007000300004)
- Torrano-Silva, B.N. and E.C. Oliveira. 2013. Macrophytobenthic flora of the Abrolhos Archipelago and the Sebastião Gomes Reef, Brazil. *Continental Shelf Research* 70: 150–158. doi: [10.1016/j.csr.2013.09.019](https://doi.org/10.1016/j.csr.2013.09.019)
- Villaça, R., A.C. Fonseca, V.K. Jensen and B. Knoppers. 2010. Species composition and distribution of macroalgae on Atol das Rocas, Brazil, SW Atlantic. *Botanica Marina* 53: 113–122. doi: [10.1515/BOT.2010.013](https://doi.org/10.1515/BOT.2010.013)
- Villaça, R.C., A.G. Pedrini, S.M.B. Pereira and M.A.O. Figueiredo. 2006. Flora Marinha Bentônica das Ilhas Oceânicas Brasileiras, pp. 105–146, In: R.J.V. Alves, J.W.A. Castro (orgs.). Ilhas Oceânicas

Brasileiras. Da pesquisa ao Manejo. Brasília: Ministério do Meio Ambiente.

Wynne, M.J. 1997. *Nitophyllum adhaerens* sp. nov. (Delesseriaceae, Rhodophyta) from the Caribbean and Bermuda. *Cryptogamie, Algologie* 18: 211–221.

Wynne, M.J. 2011. A checklist of benthic marine algae of the tropical and subtropical western Atlantic: third revision. *Nova Hedwigia Beihefte* 140: 1–166.

**Authors' contribution statement:** CWNM, AMA, AAS, WRA, KLMB, ISO, NSF, IBO, GJPR wrote the text and collected the data.

**Received:** 17 July 2014

**Accepted:** 9 July 2015

**Academic editor:** Sandra Costa-Böddeker