



First record of genus *Imparfinis* from a northeastern coastal Brazilian river basin: *I. borodini* Mees & Cala, 1989 in Rio de Contas, Bahia

**Luisa Maria Sarmiento-Soares^{1,2*}, Heraldo A. Britski³, Marcia Anjos⁴, Angela M. Zanata⁵,
Ronaldo F. Martins-Pinheiro¹ and Marluce Galvão Barretto⁴**

1 Instituto Nacional da Mata Atlântica, Laboratório de Zoologia. Av. José Ruschi, 4, Centro, 29650-000 Santa Teresa, ES, Brazil

2 Programa de Pós-Graduação em Biologia Animal (PPGBAN), Universidade Federal do Espírito Santo (UFES), Campus de Goiabeiras, 29043-900 Vitória, ES, Brazil

3 Museu de Zoologia, Universidade de São Paulo, Caixa Postal 42494, 04218-970 São Paulo, SP, Brazil

4 Programa de Pós-Graduação em Genética, Biodiversidade e Conservação, Universidade Estadual do Sudoeste da Bahia (UESB), Campus Jequié, Jequié, BA, Brazil

5 Departamento de Zoologia, Instituto de Biologia, Universidade Federal da Bahia, Campus de Ondina, Rua Barão de Geremoabo s/n, 40170-290, Salvador, BA, Brazil

* Corresponding author. E-mail: luisa@nossosriachos.net

Abstract: *Imparfinis borodini* Mees & Cala, 1989 is recorded in low population abundance in the upper Paraná, Tocantins and São Francisco river drainages according to the scientific literature and available collection data, but it has not been previously recorded from a coastal Brazilian river. Herein, the geographic range of this species is extended to the Rio de Contas basin in northeastern Brazil. This represents the first record in a coastal system. Additionally, *I. borodini* was captured in the middle Rio São Francisco at Barreiras. Both new records are the first from Bahia state, Northeastern Brazil.

Key words: Atlantic Forest; Caatinga; Heptapteridae; Northeastern Brazil; Rio São Francisco; taxonomy

The catfish family Heptapteridae includes peculiar forms, with troglomorphic species such as *Rhamdiopsis krugi* Haseman, 1911, from Chapada Diamantina, Bahia state (Bockmann and Castro 2010), and miniature species, which besides small size (less than 20 mm in standard length), show reduction on laterosensory canals, bones and number of rays, as exemplified by *Horiomyzon retropinnatus* D.J. Stewart, 1986 from the upper Amazon (Weitzman and Vari 1988). Members of Heptapteridae are arranged in 24 genera that include 208 valid species (Eschmeyer and Fong 2015). The group is one of the most representative members of the order Siluriformes in small water bodies in South America.

Although the family is recognized as monophyletic (Bockmann and Guazelli 2003), it lacks external morphological attributes that allow easy recognition, being

instead identified by a group of general anatomical features present in other Neotropical families (Lundberg and McDade 1986; Lundberg et al. 1991; Bockmann 1998). In spite of the unquestionable family monophyly, the taxonomic status of some genera remains less precise and such situation includes *Imparfinis* Eigenmann & Norris, 1910, as stated by Ortega-Lara et al. (2011). On the other hand, the *Nemuroglanis* subclade is a well-corroborated group within heptapterids (Ferraris 1988; Bockmann 1994), recognized on the basis of 16 synapomorphies, mostly associated with the Weberian complex and fin supports. This clade comprises *Acentronichthys* Eigenmann & Eigenmann 1889, *Cetopsorhamdia* Eigenmann & Fisher 1916, *Chasmocranus* Eigenmann 1912, *Heptapterus* Bleeker 1858, *Horiomyzon* Stewart 1986, *Imparfinis* Eigenmann & Norris 1900, *Mastiglanis* Bockmann 1994, *Nannoglanis* Boulenger 1847, *Nemuroglanis* Eigenmann & Eigenmann 1889, *Pariolius* Cope 1872, *Phenacorhamdia* Dahl 1961, *Rhamdioglanis* Ihering 1907, *Rhamdiopsis* Haseman 1911, and *Taunayia* Miranda-Ribeiro 1918 (Lundberg and McDade 1986; Bockmann and Ferraris 2005). *Phreatobius* Goeldi 1905, stated as a member of heptapterid clade *Nemuroglanis* by the above mentioned authors, was recently recognized as a separate clade recognized within the superfamily Pimelodoidea. In spite of its highly distinct morphology, such features do not provide strong evidence for family-level placement (Sullivan et al. 2013).

The genus *Imparfinis* is a poorly diagnosed genus, with generic recognition based on an array of osteological features, as presented in Bockmann (1998) and also the presence of a dark band along lateral line. Additional

external features for generic recognition include branchiostegal rays 6–8, usually 7; branched rays on pectoral fin 7–10, usually 8 or 9; anal fin bearing 11–14 rays, usually 12–13 rays; caudal fin with 7 branched rays on upper lobe and 8 in lower lobe (Bockmann 1998). The presence of conspicuous bars on flanks is observed in some *Imparfinis* species, including *I. borodini* Mees & Cala, 1989. Although the number of bars varies between the species, such a feature deserves further investigation for generic assignment, since it is also observed in *Chasmocranus* species.

The genus includes 19 species (Eschmeyer and Fricke 2015), and is one of the most broadly distributed within the family Heptapteridae, from streams in Costa Rica to the Paraná and Uruguay river basins in Argentina, and to both sides of the Andean cordillera (Ortega-Lara et al. 2011). The easternmost record is that of *I. borodini*, for the São Francisco River basin.

Imparfinis borodini was described in 1989 from Franca, Rio Grande basin, upper Paraná system, São Paulo, on the basis of a single specimen (AMNH 8639). The name *I. borodini* was proposed in face of a homonymy case, in substitution of *I. longicauda* Borodin (1927), preoccupied by its senior homonym *I. longicauda* Boulenger (1887), that was described from the Rio Bobonaza basin, upper Rio Pastaza drainage, Ecuador (Mees and Cala 1989: 387).

The species is characterized by the elongate, slender body, ending in a very long falcate caudal fin obliquely asymmetric; long adipose fin; and 4 to 5 transverse bars on flanks (Borodin 1927). This species has fins devoid of spines, with 7 to 8 branched rays on pectoral and anal fins, 6 branched rays on upper caudal lobe and 7 on lower; eyes in dorsal position; low body depth (10–12% in standard length [SL]); and presence of a dark stripe along the lateral line (LMSS, pers. obs.).

Currently, *I. borodini* is recorded for upper Tocantins, upper Paraná and São Francisco river drainages. A single record is given to the upper Tocantins at Serra da Mesa River dam in Goiás state (Miranda and Mazzoni 2003). In the upper Paraná drainage, this species is widely distributed, with records for the Brazilian states of Goiás (Pavanelli et al. 2007), São Paulo (Rondineli and Braga 2010; Oyakawa and Menezes 2011), Paraná (Galves et al. 2007; Cunico et al. 2009), and Mato Grosso (collection data). According to the literature, the species occurs in low population abundance, and is found in third and fourth order streams (e.g., Pinto 2009; Takahashi 2010). For the Rio São Francisco basin the species was recorded in Minas Gerais state, at its upper and middle stretches, and also in the Distrito Federal (Species Link; Figures 1 and 2). Otherwise, no literature records indicate its occurrence for the




Figure 1. Records of *Imparfinis borodini* (red dots) in Brazil. Additional records in Bahia are indicated by green rectangles. See Figure 2 for additional information on Bahia records. The major Brazilian river basins are indicated by green and dark blue and are the Paraná, Tocantins and São Francisco basins, respectively. The coastal Atlantic drainage is indicated by light blue. Bahia State is outlined in yellow.




Figure 2. Records of *Imparfinis borodini* in Bahia State, Brazil. The basins of the Rio São Francisco and Rio de Contas are green and light pink, respectively.

sub-middle and lower stretches of the Rio São Francisco in Bahia state.

Recent samplings in the Rio de Contas basin revealed the presence of *Imparfinis borodini*, representing the first record of the species from this drainage (Figures 2 and 3). The Northeastern Mata Atlântica freshwater ecoregion, as stated by Abell et al. (2008), includes all coastal river basins between the Rio Itabapoana (Rio de Janeiro and Espírito Santo states) in the south and Rio Sergipe (Sergipe state) in the north. The whole area is delimited on the west by the Rio São Francisco watershed along the Espinhaço mountain range. The record of *I. borodini* in the Rio de Contas basin constitutes the

first occurrence of the genus in the Northeastern Mata Atlântica freshwater ecoregion (Figure 2; ecoregion 328 of Abell et al. 2008).

The Rio de Contas basin, situated in central-south Bahia and the largest basin completely within in the state, occupies about 10.2% of the state's territory. Near its headwaters, in Jequié municipality, the Rio de Contas is dammed for energy production by the Barragem de Pedra, which is controlled by the Companhia Hidro Elétrica do São Francisco. Downstream from that barrier the main tributaries of Rio de Contas are Rio Jequezinho and the Rio Preto do Criciúma. The Rio Preto do Criciúma a left margin tributary in the middle



Figure 3. *Imparfinis borodini*, MBML 30327, 68.9 mm SL, Rio Preto do Criciúma, Rio de Contas basin, Jitaúna, Bahia, Brazil. Lateral and dorsal views.

stretch of the Rio de Contas, in the municipality of Jitaúna, drain a transition zone between the Caatinga and Atlantic Forest biomes. It is situated between Jequié and Jitaúna municipalities and was the object of research on fish communities (e.g., Barreto and Xavier 2003; Xavier 2003; Oliveira 2012; Souza 2015).

One specimen of *Imparfinis borodini* was captured in the Rio Preto do Criciúma (Figure 4). The stretch of the Rio Preto do Criciúma where *Imparfinis borodini* was found is shallow (ca. 20 cm to 1 m deep), with rapids over rocks, sand and silt, and pools about 2.5 m deep and 13 m wide. Waters are clear, warm (21–31°C) and oxygenated (7.8 mg/l). Water current between pools and rapids were 0.36–0.55 m/s (Barreto and Xavier 2003). Marginal vegetation consists in grass, and remnants of riparian forest. Along the Rio Preto do Criciúma granite blocks were quarried at the end of the 20th century. Nowadays, the edges of the river are occupied by familiar agriculture, mainly horticulture.



Figure 4. Collecting locality of *Imparfinis borodini* at Rio Preto da Criciúma, downstream of the Rio de Pedras dam, Rio de Contas basin, in Jitaúna near border with Jequié, Bahia, Brazil.

The other specimens of *I. borodini* were captured in two stretches of the Rio de Ondas (Figure 5), a tributary of the Rio Grande, which is itself a tributary of the middle portion of the São Francisco River. The sampled stretch (ca. 1.2 m deep) was relatively fast running and clear and predominantly rocky with sand and pebbles. The remaining marginal vegetation with shrubs and trees of Buriti (*Mauritia flexuosa*). Various anthropic disturbances are found along the river, including water extraction for irrigation and supplying the cities of Barreiras and Luís Eduardo Magalhães, and for recreation (Moraes 2003). *Imparfinis borodini* was found associated with vegetation debris, near the riverbank in a flooded stretch with a sequence of cascades (ca. 40–60 cm high).

This species was recorded in syntopy with *Astyanax* sp., *Hypostomus* sp., *Pareiorhaphis* sp., *Parotocinclus jimi* Garavello 1977, and *Geophagus* sp. in the rio Preto do Criciúma, and with *Astyanax fasciatus* (Cuvier 1819), *Bryconops* aff. *affinis*, *Cichlasoma sanctifranciscense* Kullander 1983, *Myelus micans* (Lütken 1875), and *Serrapinnus heterodon* (Eigenmann 1915) in the Rio de Ondas. Some of these species have a preference for well-oxygenated waters in rapids.

Imparfinis borodini was defined as of Least Concern, and not threatened with extinction in the national Brazilian Red List (Brazil, Ministério do Meio Ambiente 2014). Particularly, the conservation status of *I. borodini* is uncertain in the state of Bahia, based on the limited knowledge of its geographic distribution. The species is so far known from two areas, a tributary of the Rio de Contas, subject to permanent anthropic perturbation, and a tributary of the Rio São Francisco, also with a series of anthropic alterations. However, given the absence of studies on population biology and geographical range of the species, we are unable to assess the effect of these putative threats on the conservation of *I. borodini* and on the maintenance of its populations within the




Figure 5. Collecting localities of *Imparfinis borodini* along the Rio de Ondas, Barreiras, Bahia, Brazil.

state of Bahia. Thus, *I. borodini* would be classified as data deficient (DD), albeit not assessed using the International Union for Conservation of Nature (IUCN) categories and criteria. Additional collecting efforts should be conducted in that region in order to better understand biological aspects and distribution of the species.

The species was not included among the freshwater fish species of Bahia, during the meetings for evaluation of endangered fauna of the state (Jucá Chagas et al. 2015).

A recent study of the species distributions of the Northeastern Mata Atlantica ecoregion and the fish fauna shared with adjacent ecoregions by Camelier and Zanata (2014), listed 28 species shared between Northeastern Mata Atlantica and the São Francisco ecoregion (without considering species that are widely distributed and taxonomically problematic species, such as *Characidium* spp., *Hypostomus* spp., *Trichomycterus* spp.). The discovery of new occurrences, such as those of *I. borodini*, and the lack of knowledge of the distributions of species within taxonomically difficult groups of the above genera attests for the incomplete taxonomic knowledge of the fish diversity of the ecoregion. Taxonomic revision of these problematic species groups and samplings efforts are encouraged to better understand the composition and distribution of species throughout Northeastern Mata Atlantica river systems.

Regarding the species shared between the Northeastern Mata Atlantica and the São Francisco ecoregions, most of these occur in rivers, such as the Rio Itapicuru and the Rio Paraguaçu, in the northern portion of the Northeastern Mata Atlantica. Camelier and Zanata (2014) recorded three of those species in the Rio São Francisco basin and a group of rivers of the Northeastern Mata Atlantica ecoregion that includes Rio de Contas, named Central-South group (*Cichlasoma sanctifranciscense*, *Pamphorichthys hollandi* (Henn 1916), and *Phenacogaster franciscoensis* Eigenmann 1911). *Imparfinis borodini* represents the fourth species shared by the Rio de Contas and São Francisco basins. According to Camelier and Zanata 2014, the southernmost limit of distribution of these shared species is the Rio de Contas. South of the Rio de Contas, the taxonomic affinity and the presence of shared species between the eastern Brazilian basins and the Rio São Francisco basin clearly decreases.

Geomorphological evidence indicates an ancient drainage connection among eastern basins and the Rio São Francisco basin. Cenozoic captures of tributaries of the ancient São Francisco drainage are evidenced by the large number of wind gaps, which are valleys where water once flowed (Saadi 1998). Dry former watercourses bordering the Espinhaço mountain range,

are testimonies of old connections between the São Francisco and coastal basins (King 1956).

The Espinhaço mountain chain represents the most extensive and continuous elevation within Brazilian territory (Almeida-Abreu and Renger 2002). It was formed during the Mesoproterozoic Era as a product of cover sequence over the adjacent stable eastern São Francisco craton, partially reworked by the Brasiliano Araçuaí Fold Belt (Almeida-Abreu 1995; Almeida et al. 2000). During the Mesozoic and Cenozoic eras erosive events, denudation cycles, and blocks tilting occurred. During this time, a connection may have occurred between the Rio de Contas and São Francisco basins that would explain the shared fish fauna. However, such stream captures are not yet fully comprehended. Further investigation into past river connections and their timing are needed for a better understanding of faunal connections.

Ribeiro (2006) applied the term “Pattern C” to distributions that reflect the most recent exchanges between the upland crystalline shield rivers and the adjacent coastal drainages, recognized on the basis of shared species, and, in some cases, of hybrid zones. According to Camelier and Zanata (2014) six species are currently known to occur only in the Rio São Francisco basin and basins of the north portion of Northeastern Mata Atlantica: *Hyphessobrycon micropterus* (Eigenmann 1915), *Lepidocharax burnsi* Ferreira, Menezes & Quaggio-Grassiotto 2011, *Oligosarcus argenteus* Günther 1864, *Prochilodus costatus* Valenciennes 1850, *Steindachnerina elegans* (Steindachner 1875), and *Phenacogaster franciscoensis*. These species appear to exemplify “Pattern C” as defined by Ribeiro (2006). Most of these species have a somewhat broad distribution, occurring also in some of the coastal drainages north to the Rio São Francisco, and also in the Rio Paraná basin. In turn, the authors also state that only *Acestrorhynchus lacustris* (Lütken 1875), *Cetopsorhamdia iheringi* Schubart & Gomes 1959, *Hemigrammus marginatus* Ellis 1911, *Hyphessobrycon bifasciatus* Ellis 1911, *Piabina argentea* Reinhardt 1867, *Prochilodus vimboides* Kner 1859, *Serrapinnus heterodon* (Eigenmann 1915), *S. piaba* (Lütken 1875), and *Trachelyopterus striatulus* (Steindachner 1877) are shared between the drainages of the Northeastern Mata Atlantica ecoregion and the Rio Paraná basin (LMSS and AMZ, pers. obs.). On the other hand, *Imparfinis borodini* is shared between both São Francisco (ecoregion 327), Northeastern Mata Atlantica (ecoregion 328), Upper Paraná (344 ecoregion) and Tocantins-Araguaia (ecoregion 324) freshwater ecoregions (numbers in parentheses *sensu* Abell et al. 2008).

Biogeographical explanations aside, the *I. borodini* population in the Rio de Contas is known only by a single specimen that was found by us. It is interesting to note that Borodin (1927) stated in the original description of

the species that the adipose fin does not reach the caudal fin and this condition can be verified in his illustration (Figure 3 of Borodin, 1927) and in the photograph of the holotype (AMNH 8639; available at ACSI 2015). The rio de Contas specimen has the adipose fin almost connected to the caudal fin. Further investigations regarding the population of *I. borodini* in Rio de Contas are desirable, once additional specimens are found.

Comparative material examined. Brazil: Bahia. Jitaúna: MBML 10327, 1 (68.9 mm SL), Rio Preto do Criciúma, downstream from Barragem das Pedras dam, near border with Jequié, Rio de Contas basin, 13°57'55.9" S, 039°05'26.8" W, 267 above sea level (a.s.l.), M. Anjos and L.A.S. Santos, 12 January 2015.

Barreiras: MBML 10729, 12 (61.6–102.2 mm SL), lower stretch of Rio de Ondas, upstream of Barreiras, Rio Grande sub-basin, Rio São Francisco basin, 12°07'47" S, 045°02'06" W, M.C. Moreira and E.R.R.S. Castro. UFBA 6162, 1 (98.3 mm SL), Rio de Ondas on its confluence with Rio das Pedras, Rio Grande sub-basin, Rio São Francisco basin, 677 a.s.l., 12°14'21" S, 045°17'07" W. R. Burger and J. A. Reis, 12 February 2009.

Additional material in Species Link. Brazil: Bahia. São Desidério: UNT 09543, 2, Rio dos Angicos, Rio Corrente sub-basin, São Francisco basin, 12°42'25" S, 045°50'44" W, A. Akama and A.B. Soares, 25 October 2008.

Roda Velha: UNT 10048, 1, Rio Roda Velha, in Roda Velha de Baixo village, Rio Grande sub-basin, São Francisco basin, 11°59'00" S, 048°39'34" W, A.B. Soares, E.F. Oliveira and V.C.O. Marto, 11 October 2010.

ACKNOWLEDGEMENTS

We are indebted to Rafael Burger and José Amorim Reis for information on *Imparfinis borodini* in Barreiras. We also thank Elis Regina Rodrigues de Souza Castro for making available specimens from rio de Ondas. This study received financial support from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, PCI- E1 grant to LMSS (process # 465023/2014-2). Permission for collecting specimens was granted by ICMBio.

LITERATURE CITED

- Abell, R., M.L. Thieme, C. Revenga, M. Bryer, M. Kottelat, N. Bogutskaya, B. Coad, N. Mandrak, S.C. Balderas, W. Bussing, M.L.J. Stiassny, P. Skelton, G.R. Allen, P. Unmack, A. Naseka, R. Ng, N. Sindorf, J. Robertson, E. Armijo, J.V. Higgins, J.J. Heibel, E. Wikramanayake, D. Olson, H.L. López, R.E. Reis, J.G. Lundberg, M.H. Sabaj-Pérez and P. Petry. 2008. Freshwater ecoregions of the world: a new map of biogeographic units for freshwater biodiversity conservation. *BioScience* 58(5): 403–414. doi: <http://dx.doi.org/10.1641/B580507>
- ACSI- All Catfish Species Inventory Project. 2015. World Wide Web electronic publication. Accessed at <http://acsi.acnatsci.org/base/index.html> 23 December 2015.
- Almeida, F.F.M., B.B.B. Neves and C.D.R. Carneiro. 2000. The origin and evolution of the South American Platform. *Earth-Science Reviews* 50: 77–111. <http://www.sciencedirect.com/science/journal/00128252>
- Almeida-Abreu, P.A. 1995. O Supergrupo Espinhaço: O rifte, a bacia e o orógeno. *Geonomos* 3: 1–18.
- Almeida-Abreu, P.A. and F.E. Renger. 2002. Serra do Espinhaço Meridional: Um Orógeno de Colisão do Mesoproterozóico. *Revista Brasileira de Geociências* 32(1): 1–14.
- Barretto, M. G. and A.T. Xavier. 2003. Composição e abundância de uma comunidade de peixes de dois trechos do rio Preto do Criciúma, Jequié, Jitauna/BA; p. 243, in: XV Encontro de Ictiologia, 2003, Rumos da Ictiologia Brasileira, São Paulo.
- Bockmann, F.A. 1994. Description of *Mastiglanis asopos*, a new pimelodid catfish from northern Brazil, with comments on phylogenetic relationships inside the subfamily Rhamdiinae (Siluriformes, Pimelodidae). *Proceedings of the Biological Society of Washington* 107(4): 760–777. <http://biodiversitylibrary.org/page/35515405>
- Bockmann, F.A. 1998. Análise filogenética da família Heptapteridae (Teleostei, Ostariophysi, Siluriformes) e redefinição de seus gêneros [Doctoral thesis]. São Paulo: Universidade de São Paulo, Instituto de Biociências. 614 pp. Abstract at <http://www.bv.fapesp.br/pt/bolsas/92751/analise-filogenetica-da-familia-heptapteridae-teleostei-ostariophysi-siluriformes-com-uma-redefi/>
- Bockmann, F.A. and R.M.C. Castro. 2010. The blind catfish from the caves of Chapada Diamantina, Bahia, Brazil (Siluriformes: Heptapteridae): description, anatomy, phylogenetic relationships, natural history, and biogeography. *Neotropical Ichthyology* 8(4): 673–706. doi: [10.1590/S1679-62252010000400001](https://doi.org/10.1590/S1679-62252010000400001)
- Bockmann, F.A. and C.J. Ferraris Jr. 2005. Systematics of the Neotropical Catfish Genera *Nemuroglanis* Eigenmann and Eigenmann 1889, *Imparaleis* Schultz 1944, and *Medemichthys* Dahl 1961 (Siluriformes: Heptapteridae). *Copeia* 2005(1): 124–137. <http://www.jstor.org/stable/4098627>
- Bockmann, F.A. and G.M. Guazzelli. 2003. Family Heptapteridae; pp. 407–433, in: R.E. Reis, S.O. Kullander and C.J. Ferraris Jr. (eds.). Check list of the freshwater fishes of South and Central America. Porto Alegre, Edipucrs..
- Borodin, N.A. 1927. Some new catfishes from Brazil. *American Museum Novitates*, 266: 1–8. <http://hdl.handle.net/2246/4196>
- Boulenger, G.A. 1887. An account of the fishes collected by Mr. C. Buckley in eastern Ecuador. *Proceedings of the Zoological Society of London* 27: 274–283. doi: <http://dx.doi.org/10.1111/j.1096-3642.1887.tb02962.x>
- Brazil, Ministério do Meio Ambiente. Portarias Nos. 443, 444, 445, de 17 de Dezembro de 2014, Diário Oficial da União–Seção 1, 245, 110. 18 December 2014. Imprensa Nacional. <http://pesquisa.in.gov.br/imprensa/jsp/visualiza/index.jsp?data=18/12/2014&journal=1&pagina=110&totalArquivos=144>.
- Camelier, P. and A.M. Zanata. 2014. Biogeography of freshwater fishes from the Northeastern Mata Atlantica freshwater ecoregion: distribution, endemism, and area relationships. *Neotropical Ichthyology* 12(4): 683–698. doi: [10.1590/1982-0224-20130228](https://doi.org/10.1590/1982-0224-20130228)
- Cunio, A.M., W.J. Graça, A.A. Agostinho, W.M. Domingues and J.D. Latini. 2009. Fish, Maringá Urban Streams, Pirapó river drainage, upper Paraná river basin, Paraná state, Brazil. *Check List* 5(2): 273–280.
- Eigenmann, C.H. and A.A. Norris. 1900. Sobre alguns peixes de S. Paulo, Brazil. *Contribuições do Laboratório Zoológico da Universidade de Indiana*, n. 33. *Revista do Museu Paulista* 4: 349–362.
- Eschmeyer, W.N. and R. Fricke (eds.). [2015]. Catalog of fishes: genera, species, references. Accessed at <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>, 12 September 2015.
- Eschmeyer, W.N. and J.D. Fong. [2015]. Species by family/subfamily. Accessed at <http://researcharchive.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>, 12 December 2015.

- Ferraris Jr., C.J. 1988. Relationships of the Neotropical catfish genus *Nemuroglanis*, with a description of a new species (Osteichthys: Siluriformes: Pimelodidae). *Proceedings of the Biological Society of Washington* 101(3): 509–516. <http://biodiversitylibrary.org/page/34646200>
- Froese, R. and D. Pauly. 2015. FishBase. World Wide Web electronic publication. Version 04/2015. Accessed at <http://www.fishbase.org/summary/Imparfinis-borodini.html>, 23 December 2015.
- Galves, W., O.A. Shibatta and F.C. Jerrep. 2007. Fish, Taquara river basin, northern of the state of Paraná, Brazil. *Check List* 3(3): 253–259 doi: [10.15560/3.3.253](https://doi.org/10.15560/3.3.253)
- Jucá Chagas, R., F.A.C. Sampaio, A.T. da Silva and S. Campiolo. 2015. Avaliação do estado de conservação dos peixes de água doce do estado da Bahia, Brasil. In: XXI Encontro Brasileiro de Ictiologia, R-0023, Recife. Sociedade Brasileira de ictiologia.
- King, L.C. 1956. Geomorfologia do Brasil Oriental. *Revista Brasileira de Geografia* 18(2): 1–147. http://biblioteca.ibge.gov.br/visualizacao/periodicos/115/rbg_1956_v18_n2.pdf
- Lundberg, J.G. and L.A. McDade. 1986. On the South American catfish *Brachyrhamdia imitator* Myers (Siluriformes, Pimelodidae), with phylogenetic evidence for a large intrafamilial lineage. *Notulae Naturae of the Academy of Natural Sciences of Philadelphia* 463: 1–24.
- Lundberg, J.G., A.H. Bornbusch and F. Mago-Leccia. 1991. *Gladioglanis conquistador* n. sp., from Ecuador with diagnoses of the subfamilies Rhamdiinae Bleeker and Pseudopimelodinae n. subf. (Siluriformes, Pimelodidae). *Copeia* 1991(1): 190–209. doi: [10.2307/1446263](https://doi.org/10.2307/1446263)
- Mees, G.F. and P. Cala. 1989. Two new species of *Imparfinis* from northern South America (Pisces, Nematognathi, Pimelodidae). *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen, (Series C, Biological and Medical Sciences)* 92(3): 379–394.
- Miranda, J. C. and R. Mazzoni. 2003. Composição da ictiofauna de três riachos do alto rio Tocantins - GO. *Biota Neotropica* 3(1): 1–11. doi: [http://dx.doi.org/10.1590/S1676-06032003000100005](https://doi.org/10.1590/S1676-06032003000100005)
- Moraes, L.S. 2003. Diagnóstico de uso e ocupação da bacia do rio de Ondas – Barreiras – BA [Masters Dissertation]. Brasília: Universidade Católica de Brasília, Programa de Mestrado em Planejamento e Gestão Ambiental. 170 pp. Accessed at http://www.bdt.d.uec.br/tede/tde_busca/arquivo.php?codArquivo=78, 26 September 2015.
- Oliveira, I.A. 2012. Análise da diversidade do complexo “*Geophagus*” *brasiliensis* (Perciformes, Cichlidae) de bacias costeiras da Bahia a partir de dados citogenéticos e morfométricos. [Masters Dissertation]. Jequié: Universidade Estadual do Sudoeste da Bahia, Programa de Pós Graduação em Genética, Biodiversidade e Conservação. 118 pp. List accessed at http://www2.uesb.br/ppg/ppggbc/?page_id=182, 27 September 2015.
- Ortega-Lara, A., N. Milani, C. DoNascimento, F. Villa-Navarro and J.A. Maldonado-Ocampo. 2011. Two new trans-Andean species of *Imparfinis* Eigenmann & Norris, 1900 (Siluriformes: Heptapteridae) from Colombia. *Neotropical Ichthyology* 9(4): 777–793. doi: [10.1590/S1679-62252011000400009](https://doi.org/10.1590/S1679-62252011000400009)
- Oyakawa, O.T. and N.A. Menezes. 2011. Checklist dos peixes de água doce do Estado de São Paulo, Brasil. *Biota Neotropica* 11(1): 1–14. doi: [http://dx.doi.org/10.1590/S1676-06032011000500002](https://doi.org/10.1590/S1676-06032011000500002)
- Pavanelli, C.S., W.J. Graça, C.H. Zawadzki, H.A. Britski, A.P. Vidotti, G.S. Avelino and S. Veríssimo. 2007. Fishes from the Corumbá Reservoir, Paranaíba River drainage, upper Paraná River basin, state of Goiás, Brazil. *Check List* 3(1): 58–64. doi: [http://dx.doi.org/10.15560/3.1.58](https://doi.org/10.15560/3.1.58)
- Pinto, T.L.F. 2009. Preferência alimentar por insetos aquáticos em espécies de peixes de um riacho tropical [Masters dissertation]. Botucatu: Universidade Estadual Paulista, Instituto de Biociências. 102 pp. Accessed at http://www.ibb.unesp.br/posgrad/teses/zoologia_me_2009_tamara_pinto.pdf, 26 September 2015.
- Ribeiro, A.C. 2006. Tectonic history and the biogeography of the freshwater fishes from the coastal drainages of eastern Brazil: an example of faunal evolution associated with a divergent continental margin. *Neotropical Ichthyology* 4(2): 225–246. doi: [http://dx.doi.org/10.1590/S1679-62252006000200009](https://doi.org/10.1590/S1679-62252006000200009)
- Rondineli, G.R. and F.M.S. Braga. 2010. Reproduction of the fish community of Passa Cinco Stream, Corumbataí River sub-basin, São Paulo state, Southeastern Brazil. *Brazilian Journal of Biology* 70(1): 181–188. doi: [http://dx.doi.org/10.1590/S1519-69842010000100025](https://doi.org/10.1590/S1519-69842010000100025)
- Saadi, A. 1998. A geomorfologia da Serra do Espinhaço em Minas Gerais e de suas margens. *Geonomos* 3: 41–63.
- Souza, C.R. 2015. Marcadores moleculares na análise da diversidade do complexo de peixes *Geophagus brasiliensis* da Bahia. [Masters Dissertation]. Jequié: Universidade Estadual do Sudoeste da Bahia, Programa de Pós Graduação em Genética, Biodiversidade e Conservação. 54 pp. List accessed at http://www2.uesb.br/ppg/ppggbc/?page_id=181, 27 September 2015.
- SpeciesLink. 2015. Sistema de Informação integrado com dados primários de coleções científicas. Centro de Referência em Informação Ambiental (CRIA). Accessed at <http://www.splink.org.br>, 30 December 2015.
- Sullivan, J.P., J. Muriel-Cunha and J.G. Lundberg. 2013. Phylogenetic relationships and molecular dating of the major groups of catfishes of the Neotropical superfamily Pimelodoidea (Teleostei, Siluriformes). *Proceedings of the Academy of Natural Sciences of Philadelphia* 162: 89–110. doi: [http://dx.doi.org/10.1635/053.162.0106](https://doi.org/10.1635/053.162.0106)
- Takahashi, E.L.H. 2010. Ictiofauna do Córrego Rico, bacia do rio Mogi Guaçu, alto Paraná. [Doctoral Thesis]. Jaboticabal: Universidade Estadual Paulista, Centro de Aquicultura. 71 pp. Accessed at http://www.caunesp.unesp.br/publicacoes/dissertacoes_teses/teses/Tese%20Erico%20Luis%20Hoshiba%20Takahashi.pdf, 27 September 2015.
- Xavier, A.T. 2003. Composição e distribuição de uma comunidade de peixes de dois trechos do Rio Preto do Criciúma, Jequié/Jituaia/BA. Trabalho de Conclusão de Curso, Graduação em Ciências Biológicas, Universidade Estadual do Sudoeste da Bahia, UESB, Campus Jequié. 73 pp.
- Weitzman, S.H. and R.P. Vari. 1988. Miniaturization in South American Freshwater fishes, an overview and discussion. *Proceedings of the Biological Society of Washington* 101(2): 444–465. <http://biodiversitylibrary.org/page/34646129>

Author contributions: LMSS, AMZ wrote the text, HAB identified the species, RFMP provided the map, MA, MGB collected specimens and described collection site at rio de Contas, all authors reviewed the text.

Received: 28 September 2015

Accepted: 3 January 2016

Academic editor: Tiago P. Carvalho