NOTES ON GEOGRAPHIC DISTRIBUTION

Check List 12(3): 1897, 10 June 2016 doi: http://dx.doi.org/10.15560/12.3.1897 ISSN 1809-127X © 2016 Check List and Authors

Check List the journal of biodiversity de

biodiversity data

## The rediscovery of *Brownlowia tersa* (L.) Kosterm. (Malvaceae), from the Andaman Islands, India — a Near Threatened mangrove species

## P. Ragavan<sup>1\*</sup>, Alok Saxena<sup>2</sup>, P.M. Mohan<sup>1</sup>, R.S.C. Jayaraj<sup>3</sup> and K. Ravichandran<sup>4</sup>

- 1 Department of Ocean studies and Marine Biology, Pondicherry University, Brookshabad Campus, Port Blair-744112, Andaman and Nicobar Islands, India
- 2 Indira Gandhi National Forest Academy, Dehradun, New Forest Post, Uttarakhand-248006, India
- 3 Rain Forest Research Institute, Jorhat-785010, Assam

- 4 Department of Environment and Forest, Andaman and Nicobar Administration, Port Blair-744102, Andaman and Nicobar Islands, India
- Corresponding author. E-mail: pandivan.ragavan@gmail.com

Abstract: Brownlowia tersa, described by C.E. Parkinson from the Andaman Islands during the 1920s, was never recollected. The present report concerns its recent collection from North Andaman and Middle Andaman Islands with updated nomenclature, detailed description and distribution notes. It is considered a near threatened species due to habitat loss.

Key words: Andaman Island; Brownlowia tersa; India; rediscovery

The genus Brownlowia Roxb., of the family Malvaceae, is comprised of about 30 species and is widely distributed in Southeast Asia through Malaysia and the Pacific Islands (Tomlinson 1986). Two species, Brownlowia tersa and B. argentata, are known from mangrove communities and are classified as true mangrove species (Duke 1992; Giesen et al. 2006; Polidoro et al. 2010). Brownlowia tersa was first reported from the Andaman and Nicobar Islands (ANI) from Dhani Khari Creek, Middle Andaman, by Parkinson (1923, as B. lanceolata). Subsequently, B. tersa was reported by Sahni (1958), Das and Dev Ray (1989), Dagar et al. (1991), Dagar and Singh (1999), Debnath (1999) and Debnath (2004) but without locality data and detailed description. Debnath (1999) and Debnath (2004) noted that about 80 years earlier, B. tersa reportedly was abundant nearer to the large creeks of Middle Andaman and Dhanikahri Creek but had been rarely observed there recently. Venu et al. (2006) also noted that there were no recent collections of *B. tersa* from the ANI and that it may have been extirpated from the ANI. From these publications it is apparent that the precise locality of *B. tersa* in the ANI remained unknown despite many authors mentioning its distribution there. Here, we discuss taxonomy and distribution of *B. tersa* in the ANI, India.

Recent floristic study on species diversity and distribution of mangroves of the ANI revealed the occurrence of B. tersa in Shyamkund Creek (Middle Andaman) and Kalighat Creek (North Andaman) (Figure 1). Specimens were collected for assessing the numeric and multistate attributes of a wide range of vegetative and reproductive morphological characters and were identified using standard references (Parkinson 1923; Tomlinson 1986; Giesen et al. 2006). Collected specimens were also compared with electronic resources such as the Angiosperm Phylogeny Website (Stevens 2001), Tropicos of Missouri Botanical Garden, REMIB (World Biodiversity Information Network), Kew Herbarium Database, Global Plants Initiative, ePIC (electronic Plant Information Centre 2002), and IPNI (The International Plant Names Index 2012). Voucher material of B. tersa was deposited at the National Botanical Collection of Andaman and Nicobar Regional Centre, Botanical Survey of India, Port Blair.

Brownlowia tersa (L.) Kosterm. Kosterm, A.J.G.H. 1959. Penerbitan Madj. enget. Indonesia 1: 73. (Figures 2-9)

Brownlowia beccarii (Mast.) Pierre — Pierre, J.B.L.188. Fl. Forest. Cochinch. t. 130.

Brownlowia lanceolata Benth. - Bentham, G. 1861. J. Proc. Linn. Soc., Bot. 5(Suppl. 2): 57.

Dialycarpa beccarii Mast. — Mast. 1875. Linn. Soc., Bot. 14: 506. Glabraria tersa L. — Linnaeus, C. 1771. Mant. Pl. 2: 276.



Figure 1. Map showing study area. Red spots indicate localities of Brownlowia tersa.

**Specimens examined.** India, Andaman and Nicobar Islands, Middle Andaman, Shyamkund Creek, 13 March 2014 (12°28'33.86" N, 092°50'37.6" E), P. Ragavan, PBL 30987 and 30988.

Description. Shrub: spreading, height to 2 m (Figure 2). Roots: no above ground roots. Branches: grey and smooth, small braches often pubescent; Leaves: lanceolate (Figure 3), apex pointed, base rounded,  $5-20 \times 2-5$  cm, ratio of length to width >2, underside slivery grey (Figure 4), above glossy green in colour; petiole small, 1–2 cm long, grey in colour. *Inflorescences*: terminal or axial, few flowered, up to 4 cm long (Figure 8). Mature flower bud: rounded, 0.4–0.6 cm long (Figure 7); calyx, brown, 0.5 cm long, 3-5 lobed, bell shaped (Figure 6); petals 5, yellowish or pink in colour (Figure 5),  $0.3 \times 0.2$  cm, apex rounded; stamens numerous with yellow anther (Figure 4). *Mature fruits*: woody capsule, heart shaped, up to 1.5 cm long, grayish green in colour, covered with small, brown warts (Figure 9), single seeded.

**Distribution.** Rare; North and Middle Andaman islands.

**Habitat and ecology.** Found on the soft mud of intertidal estuarine banks along with *Rhizophora* and *Bruguiera* species; forms dense thickets.

**Phenology.** Flowering February to March; fruiting April to July

**Conservation status.** Categorized as Near Threatened by the IUCN (Kathiresan et al. 2010).

This study reports the occurrence of *B. tersa* from the ANI after lapse of 92 years. *Brownlowia tersa* can be readily recognized in the field by its brown-scaly twigs, lanceolate leaves with dull silvery under surface and pear-shaped, 2-valved fruits. *Brownlowia tersa* distinguished from *B. argentata* by its shrubby habitus and lanceolate leaves, (vs. medium sized tree and cordate leaves in *B. argentata*). It often grows along the banks in stands and remains nearly half submerged during high tide, withstanding the tidal surge by its



Figures 2–9. Morphological characters of *Brownlowia tersa*. 2: Habitat. 3: Lanceolate leaves. 4: Silvery grey underside of the leaf. 5: Flower with numerous stamens and petals. 6: Calyx. 7: Flower buds. 8: Inflorescences. 9: Mature fruits.

intricate root system. Globally, *B. tersa* is considered as a true mangrove species (Duke 1992; Giesen et al. 2006; Polidoro et al. 2010), but it has not been added to the mangrove flora of India in some important national and international status reports (Kathiresan and Bingham 2001; Kathiresan and Rajendran 2005; Anonymous 2008; Mandal and Naskar 2008). In ANI only, Sahni (1958) and Debnath (2004) classified it as true mangrove species but without locality data.

*Brownlowia tersa* is categorized as Near Threatened by the IUCN (Kathiresan et al. 2010; Polidoro et al. 2010). This species is threatened by habitat loss from coastal development, erosion, and the construction of shrimp and fish ponds throughout its range (Kathiresan et al. 2010). It is common in Southeast Asian countries: Bangladesh, Brunei, Cambodia, Indonesia Malaysia, Myanmar, Singapore, Philippines and Thailand (Giesen et al. 2006). In India its distribution is restricted to the east coast and the ANI; it is not known from the west coast. Along the east coast it is common in West Bengal and Odisha but rare in the Godavari estuary of Andhra Pradesh (Venu et al. 2006; Kathiresan 2010; Bhatt et al. 2011).

This study provides the only specific locality data of *B. tersa* in the Andaman Islands in many years. Among the 51 sites surveyed in the ANI, *B. tersa* was found at only two sites (Figure 1), and it is now rare in these islands. *Brownlowia tersa* has not been reported from the Nicobar Islands. In the Nicobar Islands, mangroves are found in narrow stretches because the creeks do not penetrate as far inland as in the Andaman Islands. In addition, mangroves of the Nicobar Islands were affected badly by the tsunami of 26 December 2004 (Ramachandran et al. 2005; Sridhar et al. 2006). The absence of *B. tersa* from the Nicobar Islands may be because of these factors.

*Brownlowia tersa* has long been used as a traditional folk remedy for diarrhoea, dysentery, wounds and boils. Roots have been found to possess significant antibacterial ability. Leaves possess anti-inflammatory, antioxidant, analgesic and anti-diarrhoeal activities (Hossain et al. 2013). A wide range of volatile phenolic components has also been isolated from various parts of *B. tersa* (Atanu et al. 2012).

Populations of *B. tersa* are experiencing severe loss at the range margins due to human activities and coastal development. This is an endangered species in India (Gopal and Chauhan 2006), so immediate and effective conservation measures should be taken for their protection and propagation.

## **ACKNOWLEDGEMENTS**

We are extremely grateful to the Principal Chief Conservator of Forests, Andaman and Nicobar Islands for his guidance and for ensuring necessary support in the field.

## LITERATURE CITED

- Anonymous. 2008. Mangroves for the future: national strategy and action plan, India (revised draft). Ministry of Environment and Forest, Government of India. 63 pp.
- Atanu, M.S.H., I. Ashab, M.S.I Howlader, M.R. Sarkar, K.M Muhsinin Mahbub and S.M.M. Lina. 2012. Phytochemical, anti-bacterial, anti-diarrhoeal and analgesic activity of *Brownlowia tersa* (Linn.). International Journal of Pharmaceutical and Biomedical Research 3: 157–161
- Bhatt, J.R., J.D. Macintosh, T.S. Nayar, C.N. Pandey and B.P. Nilratna (eds.). 2011. Towards conservation and Management of Mangrove ecosystems in India. New Delhi: Ministry of Envirionment and Forest. 278 pp.
- Dagar, J.C. and N.T. Singh. 1999. Plant resources of the Andaman and Nicobar Islands. Introduction, general features, vegetation and floristic elements. Dehra Dun: Bishen Singh Mahendra Pal Singh. Vol. 1: 252 pp.; vol. 2: 900 pp.
- Dagar, J.C., A.D. Mongia and A.K. Bandhyopadhyay. 1991. Mangroves of Andaman and Nicobar Islands. New Delhi: Oxford and IBH. 166 pp.
- Das, A. K. and M. K. Dev Roy. 1989. A general account of the mangrove fauna of Andaman and Nicobar Islands. Fauna and conservation areas: 4. Kolkata: Zoological Survey of India. 173 pp.
- Debnath, H.S. 1999. Tiliaceae; pp. 175–182, in P.K. Hajra, P.S.N. Raoand and V. Mudgal (eds). Flora of Andaman and Nicobar Islands. Calcutta: Botanical Survey of India.
- Debnath, H.S. 2004. Mangroves of Andaman and Nicobar islands; taxonomy and ecology. A community profile. Dehradun: Bishen Singh Mahendra Pal Singh. 133 pp.
- Duke, N.C. 1992. Mangrove floristics and biogeography; pp. 63–100, in: A.I. Robertson and D.M. Alongi (eds.). Tropical mangrove ecosystems. Washington, DC: American Geophysical Union.
- Giesen, W., S. Wulffraat, M. Zieren and L. Scholten. 2006. Mangrove guidebook for Southeast Asia. Bangkok: RAP Publication. 769 pp.
- Gopal, B. and M. Chauhan. 2006. Biodiversity and its conservation in the Sundarban mangrove ecosystem. Aquatic Sciences 68: 338–354. doi: 10.1007/s00027-006-0868-8
- Hossain, H., I.A. Jahan, S.I. Howlader, J.H. Shilpi, S.K. Dey and A. Hira. 2013. Anti-inflammatory and antioxidant activities of ethanolic leaf extract of *Brownlowia tersa* (L.) Kosterm. Oriental Pharmacy and Experimental Medicine 13: 181–189. doi: 10.1007/ s13596-013-0109-3
- IPNI. 2015. The international plant names index. Accessed at http://www.ipni.org, 10 August 2015.
- Kathiresan, K. 2010. Globally threatened mangrove species in India. Current Science 98: 1551.
- Kathiresan, K. and B.L. Bingham. 2001. Biology of mangroves and mangrove ecosystems. Advances in Marine Biology 40: 81–251.
- Kathiresan, K. and N. Rajendran. 2005. Mangrove ecosystems of the Indian Ocean region. Indian Journal of Marine Science 34: 104–113.
- Kathiresan, K., S.G. Salmo IIIE.S. Fernando, J.R. Peras, S. Sukardjo, T. Miyagi, J. Ellison, N.E. Koedam, Y. Wang, J. Primavera, O. JinEong, J. Wan-Hong Yong and V. Ngoc Nam. 2010. *Brownlowia tersa*. The IUCN Red List of threatened species 2010: e.T178835A7621783. Accessed on 11 September 2015. doi: 10.2305/IUCN.UK.2010-2.RLTS.T178835A7621783.en
- Mandal, R.N. and K.R. Naskar. 2008. Diversity and classification of Indian mangroves: a review. Tropical Ecology 49: 131–146.
- Missouri Botanical Garden. 2015. Tropicos.org. Accessed at http:// www.tropicos.org, 10 August 2015.
- Parkinson, C.E. 1923. A forest flora of the Andaman Islands. Dehradun: Bishen Singh and Mahendrapal Singh. 325 pp.

- Polidoro, B.A., K.E. Carpenter, L. Collins, N.C. Duke, A.M. Ellison et al. 2010. The loss of species: mangrove extinction risk and geographic areas of global concern. PLoS ONE 5: e10095. doi: 10.1371/journal.pone.0010095
- Ramachandran, S., S. Anitha, V. Balamurugan, K. Dharanirajan, K.E. Vendhan, M.I.P. Divien, A.S. Vel, I.S. Hussain and A. Udayaraj. 2005. Ecological impact of tsunami on Nicobar Islands (Camorta, Katchal, Nancowry and Trinkat). Current Science 89: 195–200.
- Royal Botanic Gardens, Kew. 2002. Electronic Plant Information Centre. Accessed at http://epic.kew.org/epic/, 10 August 2015.
- Sahni, K.C. 1958. Mangrove forests in the Andaman and Nicobar Islands. Indian Forester 84: 554–562.
- Sridhar, R., T. Thangaradjou, L. Kannan, A. Ramachandran and S. Jayakumar. 2006. Rapid assessment on the impact of tsunami on mangrove vegetation of the Great Nicobar Island. Journal of the Indian Society of Remote Sensing 34(1): 89–93. doi: 10.1007/ BF02990750.

- Stevens, P.F. 2001. Angiosperm phylogeny website. University of Missouri and Missouri Botanical Garden. Version 12. Accessed at http://www.mobot.org/MOBOT/research/APweb/, 10 August 2015.
- Tomlinson, P.B. 1986. The botany of mangroves. Cambridge: Cambridge University Press. 419 pp.
- Venu, P., W. Arisdasan, C.R. Magesh and T. Sathyamurthy. 2006. Brownlowia tersa (L.) Kosterm. (Tiliaceae) in India. Rheedea 16: 111–114.

**Author contributions:** PR and PMM identified the specimens, AS and KR collected the data and PR and RSCJ wrote text.

Received: 24 November 2015 Accepted: 5 May 2016 Academic editor: Nik Fadzly