



Bat assemblage in a karstic area from northern Brazil: seven new occurrences for Tocantins state, including the first record of *Glyphonycteris sylvestris* Thomas, 1896 for the Cerrado

Saulo Felix¹, Roberto Leonan M. Novaes^{2*}, Renan F. Souza³ and Leonardo S. Avilla¹

1 Universidade Federal do Estado do Rio de Janeiro, Instituto de Biociências. Av. Pasteur 458, CEP 22290-240, Rio de Janeiro, RJ, Brazil

2 Universidade Federal do Rio de Janeiro, Instituto de Biologia. Av. Carlos Chagas Filho 373, CEP 21941-902, Rio de Janeiro, Brazil

3 Universidade do Estado do Rio de Janeiro, Programa de Pós-Graduação em Ecologia e Evolução. Av. São Francisco Xavier 534, CEP 20550-013, Rio de Janeiro, RJ, Brazil

* Corresponding author. E-mail: robertoleonan@gmail.com

Abstract: The Cerrado, the second largest morphoclimatic area of South America, has many limestone outcrops with caves. However, studies of the bat fauna in karstic environments in the Cerrado are scarce. We present an inventory of bats in a karstic Cerrado area in the Tocantins state. We used mist-nets to sample caves, savannas, deciduous forests, and periurban environments. We captured 516 bats of 30 species, revealing that the study area is one of richest for bat species in the Brazilian Cerrado. Seven new occurrences of bat species were recorded for the Tocantins state, and we report the first record of *G. sylvestris* from the Cerrado biome. We recorded 21 species simultaneously at one cave, Gruta dos Moura, which is the highest species richness of bats for a single cave in the Neotropics.

Keywords: Chiroptera; conservation; community; Mammalia; Neotropical; range extension; savanna

INTRODUCTION

The Cerrado is the second largest morphoclimatic domain of South America, located in the central portion of the continent (Klink and Machado 2005). It has open vegetation (savanna) and semi-deciduous forest, with dry and mesic habitats (Oliveira-Filho and Ratter 2002). Human activities have drastically transformed part of the Cerrado, and 55% of its original territory was converted in pastures and agricultural fields (Klink and Machado 2005). Its high biological diversity and increasing rates of destruction make it a globally important biodiversity hotspot (Myers et al. 2000).

This morphoclimatic domain is one of the Neotropical ecoregions with lesser knowledge of its wildlife (Cavalcanti and Joly 2002) and biodiversity studies are imminently needed. In this aspect, species surveys are

essential to understand regional patterns of biological diversity (Soulé and Wilcox 1980) and can generate important data required for biodiversity conservation plans (Bernard et al. 2011).

An important geological aspect of the Cerrado is the abundance of limestone outcrops and the existence of large caves complexes (Jansen et al. 2012) that are essential roosts for many bat species (Kunz 1982). However, studies of the bat fauna in the karstic environments of the Cerrado are scarce (Trajano and Gimenez 1998; Bredt et al. 1999; Esbérard et al. 2005). Bernard et al. (2011) considered the Cerrado to be the second least studied of the Brazilian biomes for bats, and only 6% of its territory was minimally sampled. Given the knowledge gaps in karstic areas, we inventoried bat species in a karstic Cerrado area in Tocantins state. Here, we present a list of bat species for two caves, new records of bats for Tocantins, and conservation considerations.

MATERIALS AND METHODS

Study sites

The study area is within the priority area for the conservation of Cerrado biodiversity (Cavalcanti and Joly 2002). Our sampling points are in the municipality of Aurora do Tocantins (12°42' S, 046°24' W, 470 m above sea level), located in the southeastern Tocantins state, northern Brazil. Aurora do Tocantins is within the Paranã watershed, a subbasin of Tocantins River, and situated in the Paranã Valley. The watershed is bounded by the Serra Geral de Goiás to the east and the Serra Geral do Paranã to the west. The study area has remnants of Cerrado vegetation, including deciduous forest and savanna (Oliveira-Filho and Ratter 2002).

The study area has several massive limestone outcrops. Among the 533 municipalities listed by the




Figure 1. Study area in Aurora do Tocantins, northern Brazil, in the context of the Cerrado (in gray).

Sociedade Brasileira de Espeleologia (2014), Aurora do Tocantins has the eighth highest number of caves (99 records), although many caves in the area are not mapped and officially registered. This makes Aurora do Tocantins one of the most important karstic areas of the Brazilian Cerrado.

Data collection and analysis

We sampled four sites (Site 1: 12°34'54"S, 046°30'59"W; Site 2: 12°35'31"S, 046°30'56"W; Site 3: 12°40'04"S, 046°28'04"W; Site 4: 12°42'46"S, 046°24'19"W) composed of remnants of semi-deciduous forest and savanna in a matrix of open areas (pastures) and agricultural fields (Figure 1; Table 1). In January 2012, we performed eight sampling nights using eight to ten mist-nets (average of 8.8 nets/night, SD $\pm$ 0.8) (Zootech®, 9×3

m, 20 mm mesh). The mist-nets remained open all night (from sunset to sunrise), inspected at 30-minute intervals, and placed along trails within forest and savanna remnants. Bats were marked using small holes in wing membranes, a temporary marking technique (Bonaccorso and Symthe 1972).

The bats were identified from external characters (e.g., length of forearm, pelage coloration patterns, morphology of ears and tragus, etc.) and craniodental characters, following the literature descriptions (e.g., Simmons and Voss 1998; Lim and Engstrom 2001; Gardner 2008; Reis et al. 2007, 2013). We collected at least one individual of each species, and others that generated identification doubts. Vouchers were deposited in the mammalian collection of Museu Nacional, Universidade Federal do Rio de Janeiro (Appendix, Table A1). All procedures with animals followed the recommendations by Sikes et al. (2011). The captures and specimen collections were authorized by SISBIO/IBAMA (authorization number 4028-1/28717). The nomenclature of bats followed Nogueira et al. (2014).

The sampling effort was calculated following Straube and Bianconi (2002). We calculated the capture efficiency dividing the number of captures by the sampling effort in each locality sampled. The species richness estimation was made using Chao-1 method (Magurran

Table 1. Sampling sites of community structure of bats from karstic Cerrado of Tocantins state, northern Brazil.

Localities	Coordinates	Remnant size (ha)	Sampling effort (m ² .h)
Site 1	12°34'54" S, 046°30'59" W	350	14,256
Site 2	12°35'31" S, 046°30'56" W	700	2,856
Site 3	12°40'04" S, 046°28'04" W	400	2,856
Site 4	12°42'46" S, 046°24'19" W	4	2,856

2004) and the rarefaction curve with 95% confidence interval were performed using the software PAST 3.0 (Hammer et al. 2001).

To determine the composition of bat species occurring simultaneously in caves of Aurora of Tocantins, we chose two caves to sample: Gruta dos Moura (12°34'54"S, 046°30'59"W) and Gruta do Urso (12°34'56"S, 046°30'59"W). These caves were chosen for their ease of access and because of knowledge by one of us (LSA) of their interiors from previous studies (e.g., Castro et

al. 2013; Rodrigues et al. 2014; Soibelzon et al. 2015; see these studies for a basic description of caves). These caves are separated by approximately 500 m and have no connection with each other. We set mist-nets in the entrance of the caves before nightfall (16:30 h). The mist-nets remained open for three hours after sunset. To determine which species use the cave as a roost, only individuals captured exiting the caves were considered. Captures made in caves were not considered in the quantitative analysis of community structure.

Table 2. Bats from karstic Cerrado of Tocantins state, northern Brazil, including number of captures per sampling site, total number of capture (N) and relative abundance (%).

Taxa	Site 1	Site 2	Site 3	Site 4	N	%
Emballonuridae						
<i>Peropteryx macrotis</i> (Wagner, 1843)	5	0	0	0	5	1.0
Phyllostomidae						
Micronycterinae						
<i>Micronycteris sanborni</i> Simmons, 1996	4	1	0	0	5	1.0
<i>Micronycteris schmidtorum</i> Sanborn, 1935	1	0	0	0	1	0.2
Desmodontinae						
<i>Desmodus rotundus</i> (É. Geoffroy, 1810)	189	2	6	0	197	38.1
<i>Diphylla ecaudata</i> Spix, 1823	3	0	0	0	3	0.6
Phyllostominae						
<i>Chrotopterus auritus</i> (Peters, 1856)	6	0	0	0	6	1.2
<i>Mimon bennettii</i> (Gray, 1838)	1	0	1	0	2	0.4
<i>Phylloderma stenops</i> Peters, 1865	3	0	0	0	3	0.6
<i>Phyllostomus discolor</i> Wagner, 1843	1	0	0	0	1	0.2
<i>Phyllostomus elongatus</i> (É. Geoffroy, 1810)	5	0	0	0	5	1.0
<i>Phyllostomus hastatus</i> (Pallas, 1767)	15	0	1	4	20	3.8
<i>Trachops cirrhosus</i> (Spix, 1823)	9	0	0	0	9	1.7
Glossophaginae						
<i>Anoura caudifer</i> (É. Geoffroy, 1818)	2	0	0	2	4	0.8
<i>Glossophaga soricina</i> (Pallas, 1766)	29	2	0	1	32	6.2
Lonchophyllinae						
<i>Lionycteris spurrelli</i> Thomas, 1913	4	0	0	0	4	0.8
<i>Lonchophylla dekeyseri</i> Taddei, Vizotto & Sazima, 1983	8	0	0	0	8	1.5
Carollinae						
<i>Carollia perspicillata</i> (Linnaeus, 1758)	36	2	0	21	59	11.4
Glyphoncterinae						
<i>Glyphoncteris sylvestris</i> Thomas, 1896	1	0	0	0	1	0.2
Stenodermatinae						
<i>Artibeus planirostris</i> Spix, 1823	34	21	32	20	107	20.7
<i>Chiroderma villosus</i> Peters, 1860	0	0	1	0	1	0.2
<i>Platyrrhinus lineatus</i> (É. Geoffroy, 1810)	12	0	0	0	12	2.3
<i>Sturmira lilium</i> (É. Geoffroy, 1810)	4	0	6	0	10	1.9
Mormoopidae						
<i>Pteronotus parnellii</i> (Gray, 1843)	1	0	0	0	1	0.2
Noctilionidae						
<i>Noctilio leporinus</i> (Linnaeus, 1758)	1	0	0	0	1	0.2
Furipteridae						
<i>Furipterus horrens</i> (F. Cuvier, 1828)	5	0	0	0	5	1.0
Molossidae						
<i>Molossops temminckii</i> (Burmeister, 1854)	5	0	1	0	6	1.2
<i>Molossus molossus</i> (Pallas, 1766)	0	0	0	2	2	0.4
Vespertilionidae						
<i>Eptesicus diminutus</i> Osgood, 1915	0	1	1	0	2	0.4
<i>Eptesicus furinalis</i> (d'Orbigny & Gervais, 1847)	1	0	0	0	1	0.2
Total	385	29	52	50	516	100

RESULTS

Assemblage structure

We captured 516 specimens of 30 species: one species of Emballonuridae, 22 species of Phyllostomidae, one species of Mormoopidae, one species of Noctilionidae, one species of Furipteridae, two species of Molossidae, and two species of Vespertilionidae (Table 2).

The total sampling effort was 22.824 m²·h and the total capture efficiency was 0.022 bats/m²·h. The Chao-1 index estimated 45 species as the maximum richness (SD ± 1.0), which is an indication that with increased capture effort more species will be recorded for the region. The rarefaction curve also indicates that more species will be recorded by increasing the capture effort (Figure 2). The family with the most number of captures was Phyllostomidae, with 95% of the captures.




Figure 2. Rarefaction curve for the bat community in Aurora do Tocantins, northern Brazil.

Desmodus rotundus (É. Geoffroy, 1810) was the most abundant species with 197 captures (38%), followed by *Artibeus planirostris* Spix, 1823 and *Carollia perspicillata* (Linnaeus, 1758), with 107 (21%) and 60 captures (12%), respectively. No recapture was made.

New records

We provide seven new occurrence records to Tocantins state: *Peropteryx macrotis* (Wagner, 1843), *Anoura caudifer* (É. Geoffroy, 1818), *Chrotopterus auritus* (Peters, 1856), *Diphylla ecaudata* Spix, 1823, *Glyphoncycteris sylvestris* Thomas, 1896, *Mimon bennettii* (Gray, 1838) and *Chiroderma villosum* Peters, 1860 (Figure 3). Previously in the same locality, *Furipterus horrens* (F. Cuvier, 1828) and *Phyllostomus elongatus* (E. Geoffroy, 1810) were already reported as first occurrences for Tocantins state (Novaes et al. 2012, 2014).

The capture of *G. sylvestris* is the first record of this species for the Cerrado domain and expands the species' distribution by approximately 1,500 km south of the Amazonian populations and more than 850 km north of the Atlantic Forest populations (Figure 4). Occurrence data of this species in South America are shown in Table 3.

Caves survey

Through direct observations, mist-netting at caves exits, and manual captures, we recorded 22 species occupying two caves, with a great difference in species richness between them: Gruta dos Moura has 21 species (including 11 exclusive species) and Gruta do Urso has 11 species (with one species recorded only in this cave; Table 4). We compiled the bat richness in 139 caves from seven countries in the Neotropics (Brazil, Bolivia, Colombia, Cuba, Honduras, Mexico, and Venezuela). The species richness in caves ranged from zero in Buena



Figure 3. New records of bats for Tocantins state, northeastern Brazil. **A.** *Peropteryx macrotis*. **B.** *Anoura caudifer*. **C.** *Chrotopterus auritus*. **D.** *Diphylla ecaudata*. **E.** *Mimon bennettii*. **F.** *Chiroderma villosum*. **G.** *Glyphoncycteris sylvestris*, including a new record for the Cerrado. Photos by Roberto Leonan Morim Novaes.




Figure 4. Records of *Glypomyotis sylvestris* (dots) in South America, including a new record from Cerrado (gray) from Tocantins, Northern Brazil (star).

Vista Cave, Mexico, to 17 at Gruta da Judite, Brazil (Appendix, Table A2). Our literature review indicates Gruta dos Moura as the richest cave for bat species in the Neotropics.

DISCUSSION

Assemblage structure and new records

Bat inventories in the Cerrado presented a richness ranging from three to 39 species (e.g., Gonçalves and Gregorin 2004; Bordignon 2006; Zortéa and Alho 2008; Cunha et al. 2009; Bezerra and Marinho-Filho 2010; Ferreira et al. 2010; Zortéa et al. 2010; Cunha et al. 2011; Gregorin et al. 2011; Sousa et al. 2013; Muylaert et al. 2014; Shapiro and Bordignon 2014). However,

this richness can be considerably higher in areas with caves (Bredt et al. 1999; Esbérard et al. 2005; Gregorin et al. 2011). Brunet and Medellín (2001) indicate that species richness tends to be higher in karstic areas compared with environments without caves. Probably, the high species richness of bats in Aurora do Tocantins is due to limestone caves, which increase the availability of roosts, making possible the coexistence of a large number of species (Trajano 1995).

Bat species richness in Aurora do Tocantins is higher than the average found in other Cerrado localities, only behind Estação Ecológica Serra Geral de Tocantins, with 39 species (Gregorin et al. 2011). However, the difference between those areas can be explained by the

difference in sampling effort employed, with the effort being greater in Serra Geral do Tocantins (Gregorin et al. 2011). The capture efficiency obtained by us was

Table 4. Bats captured in two caves in Aurora do Tocantins, northern Brazil: presence (X) and absence (–).

Species	Moura Cave	Urso Cave
<i>Peropteryx macrotis</i>	X	X
<i>Glossophaga soricina</i>	X	X
<i>Lionycteris spurrelli</i>	X	–
<i>Lonchophylla dekeyseri</i>	X	–
<i>Desmodus rotundus</i>	X	X
<i>Diphylla ecaudata</i>	X	X
<i>Carollia perspicillata</i>	X	X
<i>Chrotopterus auritus</i>	X	X
<i>Glyphonycteris sylvestris</i>	X	–
<i>Micronycteris sanborni</i>	X	–
<i>Micronycteris schmidtorum</i>	X	–
<i>Mimon bennettii</i>	X	X
<i>Phylloderma stenops</i>	X	–
<i>Phyllostomus discolor</i>	X	–
<i>Phyllostomus elongatus</i>	X	–
<i>Phyllostomus hastatus</i>	X	X
<i>Trachops cirrhosus</i>	X	–
<i>Artibeus planirostris</i>	–	X
<i>Platyrrhinus lineatus</i>	X	X
<i>Furipterus horrens</i>	X	X
<i>Noctilio leporinus</i>	X	–
<i>Pteronotus parnellii</i>	X	–
Total	21	11

0.022 bats/m²·h, also considered high for Cerrado areas, with minimum values of 0.002 bats/m²·h in the Serra Geral do Tocantins (Gregorin et al. 2011) and maximum around 0.032 bats/m²·h in the Aporé-Sucuriú Complex of Mato Grosso do Sul (Bordignon 2006).

This study recorded seven new occurrences of species for Tocantins state, increasing the state's species diversity to 60 (Novaes et al. 2012; Reis et al. 2013; Novaes et al. 2014; Lapenta and Bueno 2015). Other than *G. sylvestris*, occurrences of six widely distributed species in the Brazilian Cerrado were expected in Tocantins and required only formalization with voucher deposits. Our report of *G. sylvestris* is the first from the Cerrado. This species is rare throughout its range and has two disjunct populations: one from Costa Rica to the Amazon basin, occupying areas east of the Andes, Venezuela, Colombia, French Guiana, Suriname, Peru, and Trinidad and Tobago, and another population occurring in the Atlantic Forest in southeastern Brazil (Williams and Genoways 2008). However, our record of this species from Aurora do Tocantins, in a deciduous forest area inserted into the Cerrado, is midway between the two populations. More studies are needed to determine if the population of *G. sylvestris* found in Aurora do Tocantins is relictual and has no functional connection with other populations. However, the incipient distributional information about *G. sylvestris* may result from misidentifications. According

Table 3. Occurrences of *Glyphonycteris sylvestris* in South America.

Locations	Coordinates	Authors
Salazar Trace, Trinidad and Tobago	10°07'N, 61°47'W	Goodwin and Greenhall (1961)
Brokopondo, Brownsweig, Suriname	05°00'N, 54°59'W	Williams and Genoways (1980)
Paracou, Sinnamary, French Guiana	05°16'N, 52°55'W	Simmons and Voss (1998)
Saül, French Guiana	03°38'N, 53°13'W	Brosset and Dubost (1968)
Iwokrama Forest, Potaro-Siparuni, Guyana	04°32'N, 59°05'W	Lim and Engstrom (2001)
Cueva del Guácharo, Monagas, Venezuela	10°07'N, 63°29'W	Linares (1969)
San Juan de Manapiare, Amazonas, Venezuela	05°15'N, 66°05'W	Lim and Engstrom (2001)
Socorré, Alto Rio Sinú, Bolívar, Colombia	Not mentioned	Simmons (1996)
Vereda Las Confusas, Antioquia, Colombia	05°59'N, 74°48'W	Morales-Martínez and Suárez-Castro (2014)
Estación Puerto Abeja, Caquetá, Colombia	00°04'N, 72°26'W	Montenegro and Romero-Ruiz (2000)
Sangay National Park, Morona Santiago, Ecuador	02°05'S, 78°09'W	Tirira et al. (2016)
Cordillera Vilcabamba, Cuzco, Peru	11°35'S, 73°54'W	Koopman (1978)
Ilha de Maracá, Roraima, Brazil	03°25'N, 61°39'W	Robinson (1998)
Montanhas do Tumucumaque, Amapá, Brazil	01°36'N, 52°29'W	Martins et al. (2011)
FLONA do Amapá, Amapá, Brazil	01°17'N, 51°35'W	Martins et al. (2011)
Rio Iratapuru, Amapá, Brazil	00°16'N, 53°06'W	Martins et al. (2011)
Guamá, Pará, Brazil	01°26'S, 48°25'W	Handley (1967)
Estação Científica Ferreira Penna, Pará, Brazil	01°44'S, 51°27'W	Marques-Aguiar et al. (2009)
80 km N of Manaus, Amazonas, Brazil	02°24'S, 59°43'W	Bernard (2001)
Alter do Chão, Pará, Brazil	02°30'S, 54°57'W	Bernard and Fenton (2002)
FLONA Tapajós, Pará, Brazil	03°21'S, 45°57'W	Castro-Arellano et al. (2007)
Aurora do Tocantins, Tocantins, Brazil	12°34'S, 46°31'W	Present study
Antônio Dias, Minas Gerais, Brazil	19°44'S, 42°37'W	Tavares et al. (2010)
ESEC dos Caetetus, São Paulo, Brazil	22°22'S, 49°40'W	Pedro et al. (2001)
Pedra Branca, Rio de Janeiro, Brazil	22°56'S, 43°26'W	Dias et al. (2003)
Juréia-Itatins, São Paulo, Brazil	24°24'S, 47°01'W	Gimenez and Ferrarezzi (2004)
PETAR, Alto Ribeira, São Paulo, Brazil	24°34'S, 48°41'W	Trajano (1982)
Parque Nacional do Iguaçu, Paraná, Brazil	25°43'S, 54°25'W	Sekiama et al. (2001)

to Nogueira et al. (2007), this species can be misidentified as bats of the genera *Carollia* and *Trinycteris*. Therefore, it is possible that taxonomic review of museum collections of *Glyphonotis*, *Carollia*, and *Trinycteris* specimens may yield new occurrence data for *G. sylvestris*.

We recorded other rare species, including *Micronycteris schmidtorum* Sanborn, 1935, *Phylloderma stenops* Peters, 1865, and *Lonchophylla dekeyseri* Taddei, Vizotto & Sazima, 1983. The last species is endemic to the Cerrado and Caatinga ecoregions of South America and is classified as Near Threatened (Sampaio et al. 2008). This shows the importance of conservation of the caves and the surrounding savanna and forest habitats of the Aurora do Tocantins region for the maintenance of bat diversity.

Caves surveys

Studies on bats in caves are rare throughout the Neotropics, precluding a robust data comparison. However, according to our results, Gruta dos Moura, with 21 species, has the highest richness of bat species of any cave in the Neotropical region. Surveys in Brazil revealed species richness ranged from one to 17 species of bats per cave (e.g., Esbérard et al. 1998; Trajano and Gimenez 1998; Bredt et al. 1999; Esbérard et al. 2005). Studies in other Neotropical countries indicate a maximum 12 species per cave (Appendix, Table A2). Arita (1996) found from one to 12 species per cave in a survey conducted in 36 caves in Mexico. Also in Mexico, Brunet and Medellín (2001) studied bat diversity of 20 caves recorded from zero to 11 species per cave. Molinari et al. (2012) found from four to five species in four Venezuelan caves. Siles et al. (2007) reported six to seven species in three caves in Bolivia. In Colombia, one to 10 species were reported in 13 different caves (Muñoz-Saba et al. 1999; Pérez-Torres et al. 2015). Two caves in Honduras had their richnesses estimated from three to five species (Divoll and Buck 2013), and from a cave in Cuba six species were recorded (Mora et al. 2002).

Despite the little knowledge on bat fauna in caves, some studies suggest a trend of positive correlation between the size of the cave and the number of species (Arita 1996; Brunet and Medellín 2001). Probably, this species–area relationship, where larger caves tend to have greater species richness of bats, is explained by greater humidity values and the presence of speleothems and cavities (Brunet and Medellín 2001).

Conservation considerations

Decree 6,640 of 7 November 2008 relaxed the laws to protect Brazilian caves. It represents a retreat in the legislation that protects the natural heritage and allows for the complete destruction of caves that have economic interest. The most plausible result of this loss of protection is an increase in the rate of biodiversity

loss (Novaes 2012; Trajano 2013). Moreover, the Gruta dos Moura cave and the surrounding savanna vegetation are within in an area of intensive agricultural activity and not protected by conservation units. Therefore, although it has the highest richness of bat species found in a single cave in Neotropics, this cave is susceptible on anthropogenic exploration, which may cause its complete destruction. Some caves of this karstic complex already have signs of anthropogenic disturbance, such as damaged speleothems and dumped garbage in the cave entrances. We also found signs of hunting (ammunition for firearms) at the entrance to Gruta do Urso.

Considering the expansiveness of the karstic complex of Aurora do Tocantins, we believe that the richness and abundance of bats may be greater than that reported here. Bats contribute in a crucial way to the maintenance of essential ecological services, such as seed dispersal and insect population control (Wilson 1973; Kalka et al. 2008; Kunz et al. 2011; Puig-Montserrat et al. 2015).

In Aurora do Tocantins, the destruction of caves and their surrounding vegetation may do damage to local agriculture by the loss of pest-control bat species (see Boyles et al. 2011). The loss of these caves and their bats would be a serious setback for the conservation of the remaining Cerrado and its biodiversity. Paleontological studies have shown the importance of these caves for the fossil record (Oliveira et al. 2011; Castro et al. 2013; Rodrigues et al. 2014; Soibelzon et al. 2015).

With the exception of this first study of bats from Aurora do Tocantins, there is no other information about the extant wildlife found in the municipality, making it difficult to build a case for classifying the area as a priority for conservation, which would allow its legal protection as a conservation unit.

ACKNOWLEDGEMENTS

Cristal Sauwen, Gabriella Jacob and Frederico Bonisoni Pêgo contributed to the fieldwork; UNIRIO provided logistical support. We appreciate all technical and logistic support provided during expeditions by Prefeitura de Aurora do Tocantins, Mr. Wagner Moura and the Sociedade Brasileira de Espeleologia. We thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (401812/2010-3, Edital MCT/CNPq No. 32/2010–Fortalecimento da Paleontologia Nacional / Edital 32/2010–Faixa B and 552975/2011, Apoio a Projetos de Pesquisa / Chamada MCTI/CNPq No. 23/2011–Apoio Técnico para Fortalecimento da Paleontologia Nacional); the Fundação Carlos Chagas Filho de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ) for the MSc and PhD studentship awarded to S. Felix and R.F. Souza, respectively; and the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the MSc studentship to R.L.M. Novaes.

LITERATURE CITED

- Arita, H.T. 1996. The conservation of cave-roosting bats in Yucatán, Mexico. *Biological Conservation* 76: 177–185. doi: [10.1016/0006-3207\(95\)00105-0](https://doi.org/10.1016/0006-3207(95)00105-0)
- Bernard, E. 2001. Species list of bats (Mammalia, Chiroptera) of Santarém area, Pará state, Brazil. *Revista Brasileira de Zoologia* 18(2): 455–463. doi: [10.1590/S0101-81752001000200016](https://doi.org/10.1590/S0101-81752001000200016)
- Bernard, E. and M.B. Fenton. 2002. Species diversity of bats (Mammalia: Chiroptera) in forest fragments, primary forests, and savannas in Central Amazonia, Brazil. *Canadian Journal of Zoology* 80: 1124–1140. doi: [10.1139/z02-094](https://doi.org/10.1139/z02-094)
- Bernard, E., L.M.S. Aguiar and R.B. Machado. 2011. Discovering the Brazilian bat fauna: a task for two centuries? *Mammal Review* 41: 23–39. doi: [10.1111/j.1365-2907.2010.00164.x](https://doi.org/10.1111/j.1365-2907.2010.00164.x)
- Bezerra, A.M.R. and J.S. Marinho-Filho. 2010. Bats of the Paranã River valley, Tocantins and Goiás states, Central Brazil. *Zootaxa* 2725: 41–56.
- Bonaccorso, F.J. and N. Smythe. 1972. Punch-marking bats: an alternative to banding. *Journal of Mammalogy* 53: 389–390. doi: [10.2307/1379186](https://doi.org/10.2307/1379186)
- Bordignon, M.O. 2006. Diversidade de morcegos (Mammalia, Chiroptera) do Complexo Aporé-Sucuriú, Mato Grosso do Sul, Brasil. *Revista Brasileira de Zoologia* 23: 1002–1009. doi: [10.1590/S0101-81752006000400004](https://doi.org/10.1590/S0101-81752006000400004)
- Boyles J.G., P.M. Cryan, G.F. McCracken and T.H. Kunz. 2011. Economic importance of bats in agriculture. *Science* 332: 41–42. doi: [10.1126/science.1201366](https://doi.org/10.1126/science.1201366)
- Bredt, A., W. Uieda and E.D. Magalhães. 1999. Morcegos cavernícolas da região do Distrito Federal, centro-oeste do Brasil (Mammalia, Chiroptera). *Revista Brasileira de Zoologia* 16: 731–770. doi: [10.1590/S0101-81751999000300012](https://doi.org/10.1590/S0101-81751999000300012)
- Brosset, A. and G. Dubost. 1968. Chiroptères de la Guyane Française. *Mammalia* 31: 583–594.
- Brunet, A.K. and R.A. Medellín. 2001. The species-area relationship in bat assemblages of tropical caves. *Journal of Mammalogy* 82: 1114–1122. doi: [10.1644/1545-1542\(2001\)082<1114:tsarib>2.0.CO;2](https://doi.org/10.1644/1545-1542(2001)082<1114:tsarib>2.0.CO;2)
- Burmeister, H. 1854. Systematische uebersicht der Thiere Brasiliens, welchewährend einer reise durch die Provinzen von Rio de Janeiro und Minas Geräes gesammelt oder beobachtet wurden von Dr. Hermann Burmeister. Säugethiere (Mammalia). Berlin: Georg Reimer. 342 pp.
- Castro, M.C., L.S. Avilla, M.L. Freitas and A.A. Carlini. 2013. The armadillo *Propraopus sulcatus* (Mammalia: Xenarthra) from the late Quaternary of northern Brazil and a revised synonymy with *Propraopus grandis*. *Quaternary International* 317: 80–87. doi: [10.1016/j.quaint.2013.04.032](https://doi.org/10.1016/j.quaint.2013.04.032)
- Castro-Arellano, I., S.J. Presley, L.N. Saldanha, M.R. Willig and J.M. Wunderle Jr. 2007. Effects of reduced impact logging on bat biodiversity in *terra firme* forest of lowland Amazonia. *Biological Conservation* 138: 269–285. doi: [10.1016/j.biocon.2007.04.025](https://doi.org/10.1016/j.biocon.2007.04.025)
- Cavalcanti, R.B. and C.A. Joly. 2002. Biodiversity and conservation priorities in the Cerrado region; pp. 351–367, in: P.S. Oliveira and R.J. Marquis (eds.). *Ecology and natural history of a Neotropical savanna*. New York: Columbia University Press.
- Cunha, N.L., E. Fischer, L.F.A. Carvalho and A.F. Santos. 2009. Bats of Buraco das Araras Natural Reserve, southwestern Brazil. *Biota Neotropica* 9(4): 189–195. doi: [10.1590/S1676-06032009000400019](https://doi.org/10.1590/S1676-06032009000400019)
- Cunha N.L., E. Fischer and C.F. Santos. 2011. Bat assemblage in savanna remnants of Sonora, central-western Brazil. *Biota Neotropica* 11(2): 197–201. doi: [10.1590/S1676-06032011000300017](https://doi.org/10.1590/S1676-06032011000300017)
- Cuvier, F. 1828. Description d'un nouveau genre de chauvesouris sous le nom de Furie. *Mémoires du Muséum d'Histoire Naturelle*, Paris 16: 149–155.
- Dias, D., S.S.P. Silva and A.L. Peracchi. 2003. Ocorrência de *Glyphoncteris sylvestris* Thomas (Chiroptera, Phyllostomidae) no estado do Rio de Janeiro, sudeste do Brasil. *Revista Brasileira de Zoologia* 20(2): 363–366. doi: [10.1590/S0101-81752003000200030](https://doi.org/10.1590/S0101-81752003000200030)
- Divoll, T.J. and D.G. Buck. 2013. Noteworthy field observations of cave roosting bats in Honduras. *Mastozoologia Neotropical* 20: 149–151.
- d'Orbigny, A. and P. Gervais. 1847. Mammifères; pp. 1–32, in: A. d'Orbigny (ed.) *Voyage dans l'Amérique méridionale* (le Brésil, la République orientale de l'Uruguay, la République Argentine, la Patagonie, la République du Chili, la République de Bolivie, la République du Pérou), exécuté pendant les années 1826, 1827, 1828, 1829, 1830, 1831, 1832 et 1833. Volume 4. Paris: Chez Pitois-Levrault et Cie. doi: [10.5962/bhl.title.85973](https://doi.org/10.5962/bhl.title.85973)
- Esbérard, C.E.L., L.F.S. Martins, R.C. Cruz, L.M. Costa, M.S. Nunes, E.M. Luz and A.S. Chagas. 1998. Aspectos da biologia de *Lonchorhina aurita* no Estado do Rio de Janeiro (Mammalia: Chiroptera: Phyllostomidae). *Bioikos* 11(1): 46–49.
- Esbérard, C.E.L., J.A. Motta and C. Perigo. 2005. Morcegos cavernícolas da Área de Proteção Ambiental (APA) Nascentes do Rio Vermelho, Goiás. *Revista Brasileira de Zoociências* 7(3): 311–325.
- Ferreira, C.M.M., E. Fischer and A. Pulchério-Leite. 2010. Fauna de morcegos em remanescentes urbanos de Cerrado em Campo Grande, Mato Grosso do Sul. *Biota Neotropica* 10: 155–160. doi: [10.1590/S1676-06032010000300017](https://doi.org/10.1590/S1676-06032010000300017)
- Gardner, A.L. 2008. *Mammals of South America*. Volume 1, marsupials, xenarthrans, shrews, and bats. Chicago: University of Chicago Press. 669 p.
- Geoffroy St.-Hilaire, É. 1810. Sur les Phyllostomes et les mégadermes. *Annales du Muséum National d'Histoire Naturelle* 15: 157–198.
- Geoffroy St.-Hilaire, É. 1818. Sur de nouvelles chauve-souris, sous le nom de Glossophages. *Mémoires du Muséum d'Histoire Naturelle*, Paris 4: 411–418.
- Gimenez, E.A. and H. Ferrarezzi. 2004. Diversidade de morcegos no sudeste da Mata Atlântica; pp. 214–330, in: O.A.V. Marques and W. Duleba (eds.). *Estação Ecológica Juréia-Itatins: ambiente físico, flora e fauna*. Ribeirão Preto: Editora Holos.
- Gonçalves, E. and R. Gregorin. 2004. Quirópteros da Estação Ecológica da Serra das Araras, Mato Grosso, Brasil, com o primeiro registro de *Artibeus gnomus* e *A. anderseni* para o Cerrado. *Lundiana* 5(3): 143–149.
- Goodwin, G.G. and A.M. Greenhall. 1961. A review of the bats of Trinidad and Tobago: descriptions, rabies infection, and ecology. *Bulletin of the American Museum of Natural History* 122: 187–302. <http://hdl.handle.net/2246/1270>
- Gray, J.E. 1838. A revision of the genera of bats (Vespertilionidae), and the description of some new genera and species. *Magazine of Zoology and Botany* 2: 483–505.
- Gray, J.E. 1843. Letter addressed to the curator. *Proceedings of the Zoological Society of London*. London: Academic Press. 50 pp.
- Gregorin, R., E. Gonçalves, C.C. Aires and A.P. Carmignotto. 2011. Morcegos (Mammalia: Chiroptera) da Estação Ecológica Serra Geral do Tocantins: composição específica e considerações taxonômicas. *Biota Neotropica* 11(4): 299–311. doi: [10.1590/S1676-06032011000100028](https://doi.org/10.1590/S1676-06032011000100028)
- Hammer, Ø., D.A.T. Harper and P.D. Ryan. 2001. PAST: Paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1): 1–9. http://palaeo-electronica.org/2001_1/past/issue1_01.htm
- Handley, C.O. 1967. Bats of the canopy of an Amazonian forest. *Atas do Simpósio sobre a biota Amazônica* 5(1): 211–215.
- Jansen, D.C., L.F. Cavalcanti and H.S. Lamblém. 2012. Mapa de potencialidade de ocorrência de cavernas no Brasil, na escala 1:2.500.000. *Revista Brasileira de Espeleologia* 2(1): 42–57.
- Kalka, M.B., A.R. Smith and E.K.V. Kalko. 2008. Bats limit anthropods and herbivory in a tropical forest. *Science* 320: 71. doi: [10.1126/science.1154444](https://doi.org/10.1126/science.1154444)

- 10.1126/science.1153352
- Klink, C.A. and R.B. Machado. 2005. Conservation of the Brazilian Cerrado. *Conservation Biology* 19(5): 707–713. doi: [10.1111/j.1523-1739.2005.00702.x](https://doi.org/10.1111/j.1523-1739.2005.00702.x)
- Koopman, K.F. 1978. Zoogeography of Peruvian bats with special emphasis on the role of the Andes. *American Museum Novitates* 2651: 1–33. <http://hdl.handle.net/2246/2965>
- Kunz, T.H. 1982. Roosting ecology of bats; p. 1–55, in: T.H. Kunz (ed.). *Ecology of bats*. New York: Plenum Publishing.
- Kunz, T.H., E.B. Torrez, T. Lobova and T.H. Fleming. 2011. Ecosystem services provided by bats. *Annals of New York Academy of Science* 1223: 1–38. doi: [10.1111/j.1749-6632.2011.06004.x](https://doi.org/10.1111/j.1749-6632.2011.06004.x).
- Lapenta, M.J. and A.A. Bueno. 2015. Checklist of bats (Mammalia, Chiroptera) from Tocantins and Bahia, Brazil: a gradient from Cerrado, Caatinga and Atlantic Forest. *Check List* 11(4): 1673. doi: [10.15560/11.4.1673](https://doi.org/10.15560/11.4.1673)
- Lim, B.K. and M.D. Engstrom. 2001. Species diversity of bats (Mammalia: Chiroptera) in Iwokrama Forest, Guyana, and the Guianan subregion: implications for conservation. *Biodiversity and Conservation* 10: 613–657. doi: [10.1023/A:1016660123189](https://doi.org/10.1023/A:1016660123189)
- Linares, O.J. 1969. Nuevos murciélagos para la fauna de Venezuela en el Museo de Historia Natural La Salle. *Memorias de la Sociedad de Ciencias Naturales La Salle* 29: 37–42.
- Linnaeus, C. 1758. *Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis*. Holmiae: Impensis Direct, Laurentii Salvii. 824 pp. doi: [10.5962/bhl.title.542](https://doi.org/10.5962/bhl.title.542)
- Magurran, A.E. 2004. *Measuring biological diversity*. Oxford: Blackwell. 256 pp.
- Marques-Aguiar, S., M.V.D. Aguila, G.F.S. Aguiar, L.N. Saldanha, M.M.B. Rocha and J.S. Silva Jr. 2009. Morcegos (Mammalia: Chiroptera) da Estação Científica Ferreira Penna; pp. 649–671, in: P.L.B. Lisboa (org.). *Caxiuanã: desafios para a conservação de uma Floresta Nacional na Amazônia*. Belém: Museu Paraense Emílio Goeldi.
- Martins, A.C.M., E. Bernard, R. Gregorin and W.A.S. Silva. 2011. Filling data gaps on the diversity and distribution of Amazonian bats (Chiroptera): the case of Amapá, Easternmost Brazil. *Zoologia* 28(2): 177–185. doi: [10.1590/S1984-46702011000200004](https://doi.org/10.1590/S1984-46702011000200004)
- Montenegro, O.L. and M. Romero-Ruiz. 2000. Murciélagos del sector sur de la serranía de Chiribiquete, Caquetá, Colombia. *Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales* 23: 641–649.
- Molinari, J., J.M. Nassar, A. García-Rawlings and R.J. Márquez. 2012. Singularidad biológica e importancia socioeconómica de los murciélagos cavernícolas de la península de Paraguaná, Venezuela, con propuestas para su conservación. *Revista de Ecología Latinoamericana* 17(1): 1–40.
- Mora, E.C., S. Macías, D. Rojas, A. Rodríguez, I. Quiñonez, A. García, A. Cádiz and B. Boburg. 2002. Aplicación de métodos bioacústicos y convencionales en la caracterización de la comunidad de murciélagos de la Cueva del Indio, Tapaste, La Habana, Cuba. *Revista de Biología* 16: 159–166.
- Morales-Martínez, D.M. and A.F. Suárez-Castro. 2014. New records for *Glyphoncteris* Thomas, 1896 (Chiroptera: Phyllostomidae) from Colombia. *Check List* 10(3): 639–544. doi: [10.15560/10.3.639](https://doi.org/10.15560/10.3.639)
- Muylaert, R.L., R.C. Teixeira, L. Hortenci, J.R. Estêvão, P.K. Rogeri and M.A.R. Mello. 2014. Bats (Mammalia: Chiroptera) in a Cerrado landscape in São Carlos, southeastern Brazil. *Check List* 10: 287–291. doi: [10.15560/10.2.287](https://doi.org/10.15560/10.2.287)
- Myers, N, R.A. Mittermeier, C.G. Mittermeier, G.A.B. Fonseca and J. Kent. 2000. Biodiversity hotspots for conservation priorities. *Nature* 403: 853–858. doi: [10.1038/35002501](https://doi.org/10.1038/35002501)
- Muñoz-Saba, Y., H.F. López-Arévalo and A.G. Cadena. 1999. Aportes al conocimiento de la ecología de los murciélagos de los afloramientos de mármoles y calizas, sector de Río Claro (Antioquia, Colombia). *Revista de la Academia Colombiana de Ciencias* 23: 651–658.
- Nogueira, M.R., A.L. Peracchi and R. Moratelli. 2007. Subfamília Phyllostominae; pp. 61–98, in: N.R. Reis, A.L. Peracchi, W.A. Pedro and I.P. Lima (eds.). *Morcegos do Brasil*. Londrina: Editora da Universidade Estadual de Londrina.
- Nogueira, M.R., I.P. Lima, R. Moratelli, V.C. Tavares, R. Gregorin and A.L. Peracchi. 2014. Checklist of Brazilian bats, with comments on original records. *Check List* 10(4): 808–821. doi: [10.15560/10.4.808](https://doi.org/10.15560/10.4.808)
- Novaes, R.L.M. 2012. Morcegos e cavernas: história escondida de evolução, conservação e preconceito. *Ciência Hoje* 49: 40–44.
- Novaes, R.L.M., R.F. Souza, S. Felix, C. Sauwen, G. Jacob and L.S. Avilla. 2012. New record of *Furipterus horrens* (Cuvier, 1828) (Mammalia, Chiroptera) from the Cerrado of Tocantins state with a compilation of the known distribution within Brazil. *Check List* 8(4): 1359–1361. doi: [10.15560/8.6.1359](https://doi.org/10.15560/8.6.1359)
- Novaes, R.L.M., R.F. Souza, S. Felix, C. Sauwen, G. Jacob and L.S. Avilla. 2014. Occurrence of *Phyllostomus elongatus* (É. Geoffroy, 1810) (Mammalia, Chiroptera) in the Cerrado of Tocantins and a compilation of its Brazilian distribution. *Check List* 10(2): 213–216. doi: [10.15560/10.1.213](https://doi.org/10.15560/10.1.213)
- Oliveira, E., P. Villa-Nova, F.J. Goin, and L.S. Avilla. 2011. New marsupial (Mammalia, Didelphimorphia) from the Quaternary of northern Brazil: a distinct lineage within the Didelphoidea? *Zootaxa* 3041: 51–62.
- Oliveira-Filho, A.T. and J.A. Ratter. 2002. Vegetation physiognomies and woody flora of the Cerrado biome; pp. 91–120, in: P.S. Oliveira and R.J. Marquis (eds.). *The Cerrados of Brazil: ecology and natural history of a Neotropical savanna*. New York: Columbia University Press.
- Osgood, W.H. 1915. New mammals from Brazil and Peru. *Publications of Field Museum of Natural History, Zoological Series* 10: 187–198. doi: [10.5962/bhl.title.2556](https://doi.org/10.5962/bhl.title.2556)
- Pallas, P.S. 1766. *Miscellanea zoologica*. Hagae Comitum: Apud Petrum van Cleef. xii + 224 pp. doi: [10.5962/bhl.title.69851](https://doi.org/10.5962/bhl.title.69851)
- Pallas, P.S. 1767. *Spicilegium zoologicum, quibus novae et obscurae animalium species iconibus, descriptionibus atque commentariis illustrantur*. Fasciculus tertius. Berlin: Gottl, August, Lange. 35 pp. doi: [10.5962/bhl.title.39832](https://doi.org/10.5962/bhl.title.39832)
- Pedro, W.A., F.C. Passos and B.K. Lim. 2001. Morcegos (Chiroptera, Mammalia) da Estação Ecológica dos Caetetus, estado de São Paulo. *Chiroptera Neotropical* 7(2): 136–140.
- Pérez-Torres, J., D. Martínez-Medina, M. Peñuela-Sangado, M.C. Ríos-Blanco, S. Estrada-Villegas and L. Martínez-Luque. 2015. Macaregua: the cave with the highest bat richness in Colombia. *Check List* 11(2): 1616. doi: [10.15560/11.2.1616](https://doi.org/10.15560/11.2.1616)
- Peters, W. 1856. Ueber die systematische Stellung der Gattung Mormops Leach und über die classification der Phyllostomata so wie über eine neue Art der Gattung Vampyrus, von welcher hier ein kurzer Bericht gegeben wird. *Monatsberichte der Königlichen Preufs. Akademie der Wissenschaften zu Berlin* 1857: 409–415.
- Peters, W. 1860. Eine neue Gattung von Flederthieren, Chiroderma villosus, aus Brasilien. *Monatsberichte der Königlichen Preufs. Akademie der Wissenschaften zu Berlin* 1861: 747–755.
- Peters, W. 1865. Über die zu den Vampyri gehörigen Flederthiere und über die natürliche Stellung der Gattung Antrozous. *Monatsberichte der Königlichen Preufs. Akademie der Wissenschaften zu Berlin* 1866: 503–525.
- Puig-Montserrat, X., I. Torre, A. López-Baucells, E. Guerrieri, M.M. Monti, R. Ràfols-García, X. Ferrer, D. Gisbert and C. Flaquer. 2015. Pest control service provided by bats in Mediterranean rice paddies: linking agroecosystems structure to ecological functions. *Mammalian Biology* 80(3): 237–245. doi: [10.1016/j.mambio.2015.03.008](https://doi.org/10.1016/j.mambio.2015.03.008)
- Reis, N.R., A.L. Peracchi, W.A. Pedro and I.P. Lima. 2007. Morcegos do

- Brasil. Londrina: Editora da Universidade Estadual de Londrina. 253 pp.
- Reis, N.R., M.N. Fregonezi, A.L. Peracchi and O.A. Shibatta. 2013. Morcegos do Brasil: guia de campo. Rio de Janeiro: Editora Technical Books. 252 pp.
- Robinson, F. 1998. The bats of the Ilha de Maracá; pp. 165–187, in: W. Milliken and J.A. Ratter (eds.). Maracá: the biodiversity and environment of an Amazonian rainforest. Chichester: John Wiley & Sons.
- Rodrigues, S., L.S. Avilla, L. Soibelzon and C. Bernardes. 2014. Late Pleistocene carnivores (Carnivora: Mammalia) from a cave sedimentary deposit in northern Brazil. *Anais da Academia Brasileira de Ciências* 86(4): 1641–1655. doi: [10.1590/0001-3765201420140314](https://doi.org/10.1590/0001-3765201420140314)
- Sampaio, E., B.K. Lim and S. Peters. 2008. *Lonchophylla dekeyseri*. The IUCN Red List of threatened species. Version 2015.2. Accessed at <http://www.iucnredlist.org/>, 23 July 2015.
- Sanborn, C.C. 1935. New mammals from Guatemala and Honduras. *Publications of Field Museum of Natural History, Zoological Series* 20(11): 81–85. <http://www.biodiversitylibrary.org/item/20960>
- Sekiam, M.L., N.R. Reis, A.L. Peracchi and V.J. Rocha. 2001. Morcegos do Parque Nacional do Iguaçu, Paraná (Chiroptera, Mammalia). *Revista Brasileira de Zoologia* 18(3): 749–754. doi: [10.1590/S0101-81752001000300011](https://doi.org/10.1590/S0101-81752001000300011)
- Shapiro, J.T. and M.O. Bordignon. 2014. Bat (Chiroptera) assemblages in three Cerrado fragments of Mato Grosso do Sul, southwestern Brazil. *Check List* 10: 1380–1386. doi: [10.15560/10.6.1380](https://doi.org/10.15560/10.6.1380)
- Sikes, R.S., W.L. Gannon and Animal Care and Use Committee of the American Society of Mammalogists. 2011. Guidelines of the American Society of Mammalogists for the use of wild mammals in research. *Journal of Mammalogy* 92: 235–253. doi: [10.1644/10-MAMM-F-355.1](https://doi.org/10.1644/10-MAMM-F-355.1)
- Siles, L., A. Muñoz and L.F. Aguirre. 2007. Bat diversity in three caves in a montane forest of Bolivia. *Ecotropica* 13: 67–74.
- Simmons, N.B. 1996. A new species of *Micronycteris* (Chiroptera: Phyllostomidae) from northeastern Brazil, with comments on phylogenetic relationships. *American Museum Novitates* 3158: 1–34. <http://hdl.handle.net/2246/3679>
- Simmons, N.B. and R.S. Voss. 1998. The mammals of Paracou, French Guiana: a Neotropical lowland rainforest fauna, Part 1. Bats. *Bulletin of the American Museum of Natural History* 237: 1–219. <http://hdl.handle.net/2246/386>
- Sociedade Brasileira de Espeleologia. 2014. Cadastro Nacional de Cavernas. Accessed at <http://www.cavernas.org.br/cnc/Home/Index>, 25 July 2014.
- Soibelzon, E., L.S. Avilla and M. Castro. 2015. The cingulates (Mammalia: Xenarthra) from the late Quaternary of northern Brazil: fossil records, paleoclimates and displacements in America. *Quaternary International* 377: 118–125. doi: [10.1016/j.quaint.2015.02.052](https://doi.org/10.1016/j.quaint.2015.02.052)
- Soulé, M.E. and B.A. Wilcox. 1980. Conservation biology: an evolutionary-ecological perspective. Sunderland: Sinauer and Associates. 395 pp.
- Sousa, R.F., P.C. Venere and K.C. Faria. 2013. Bats in forest remnants of the Cerrado savanna of eastern Mato Grosso, Brazil. *Biota Neotropica* 13(3): 236–241. doi: [10.1590/S1676-06032013000200023](https://doi.org/10.1590/S1676-06032013000200023)
- Spix, J. de. 1823. *Simiarum et Vespertilionum brasiliensium species novae*. Monachii: Typis Francisci Seraphi Hybschmann. 72 p.
- Straube, F.C. and G.V. Bianconi. 2002. Sobre a grandeza e a unidade utilizada para estimar esforço de captura com utilização de redes-de-neblina. *Chiroptera Neotropical* 8(2): 150–152.
- Taddei V.A., L.D. Vizotto and I. Sazima. 1983. Uma nova espécie de *Lonchophylla* do Brasil e chave para identificação das espécies do gênero (Chiroptera, Phyllostomidae). *Ciência e Cultura* 35: 625–629.
- Tavares, V.C., L.M.S. Aguiar, F.A. Perini, F.C. Falcão and R. Gregorin. 2010. Bats of the state of Minas Gerais, southeastern Brazil. *Chiroptera Neotropical* 16(1): 675–705.
- Thomas, O. 1896. On new small mammals from the Neotropical Region. *Annals and Magazine of Natural History* 18(106): 301–314. doi: [10.1080/00222939608680459](https://doi.org/10.1080/00222939608680459)
- Thomas, O. 1913. A new genus of glossophagine bat from Colombia. *Annals and Magazine of Natural History* 12(69): 270–271. doi: [10.1080/00222931308693397](https://doi.org/10.1080/00222931308693397)
- Tirira, D.G., M.A. Camacho, N. Tinoco, M.F. Solórzano and S.F. Burneo. 2016. Genus *Glyphonycteris* Thomas, 1896 (Mammalia: Chiroptera) in Ecuador: first confirmed record of *G. sylvestris* Thomas, 1896 and a geographical review to *G. daviesi* (Hill, 1965). *Check List* 12(5): 1965. doi: [10.15560/12.5.1965](https://doi.org/10.15560/12.5.1965)
- Trajano, E. and E.A. Gimenez. 1998. Bat community in a cave from eastern Brazil, including a new record of *Lionycteris* (Phyllostomidae, Glossophaginae). *Studies on Neotropical Fauna and Environment* 33: 69–75.
- Trajano, E. 1982. New records of bats from southeastern Brazil. *Journal of Mammalogy* 63: 529–531. doi: [10.2307/1380462](https://doi.org/10.2307/1380462)
- Trajano, E. 1985. Ecologia de populações de morcegos cavernícolas em uma região cárstica do sudeste do Brasil. *Revista Brasileira de Zoologia* 2(5): 255–320. doi: [10.1590/S0101-81751984000100001](https://doi.org/10.1590/S0101-81751984000100001)
- Trajano, E. 1995. Protecting caves for the bats or bats for the caves? *Chiroptera Neotropical* 1(1): 19–22.
- Trajano, E. 2013. Variações anuais e infra-anuais em ecossistemas subterrâneos: implicações para estudos ambientais e preservação de cavernas. *Revista de Biologia* 10: 1–7.
- Wagner, J.A. 1843. Diagnosen neuer Arten brasilischer Handflügler; pp. 365–368, in: G. von A.F.A. Wiegmann and W.F. Erichson (eds.). *Archiv für Naturgeschichte* 9(1). Berlin: Nicolai. <http://biodiversitylibrary.org/page/13703071>
- Williams, S.L. and H.H. Genoways. 1980. Results of the Alcoa Foundation-Suriname Expeditions. II. Additional records of bats (Mammalia: Chiroptera) from Suriname. *Annals of Carnegie Museum* 49: 213–236.
- Williams, S.L. and H.H. Genoways. 2008. Subfamily Phyllostominae; pp. 255–299, in: A.L. Gardner (ed.). *Mammals of South America*, volume 1: marsupials, xenarthrans, shrews, and bats. Chicago and London: University of Chicago Press.
- Wilson, D.E. 1973. Bat faunas: a trophic comparison. *Systematic Zoology* 22: 14–29. doi: [10.2307/2412374](https://doi.org/10.2307/2412374)
- Zortéa, M. and C.J.R. Alho. 2008. Bat diversity of a Cerrado habitat in central Brazil. *Biodiversity and Conservation* 17: 791–805. doi: [10.1007/s10531-008-9318-3](https://doi.org/10.1007/s10531-008-9318-3)
- Zortéa, M., F.R. Melo, J.C. Carvalho and Z.D. Rocha. 2010. Morcegos da Bacia do rio Corumbá, Goiás. *Chiroptera Neotropical* 16(2): 611–617.

Author contributions: SF, RLMN and RFS collected the data; SF, RLMN and LSA delineated the study, RLMN and RFS made analysis, all authors wrote the text.

Received: 2 June 2016

Accepted: 12 October 2016

Academic editor: Sergio Solari

APPENDIX

Table A1. Vouchers of bats from Aurora Tocantins deposited in the Museu Nacional, Universidade Federal do Rio de Janeiro, Brazil (MN).

Family	Species	Voucher number
Emballonuridae	<i>Pteropteryx macrotis</i>	MN81194, MN81196, MN81220, MN81221
Phyllostomidae	<i>Anoura caudifer</i>	MN81187, MN81188
	<i>Glossophaga soricina</i>	MN81183, MN81193, MN81199, MN81200, MN81202, MN81203, MN81209, MN81211, MN81212, MN81214, MN81215, MN81216, MN81218
	<i>Lionycteris spurrelli</i>	MN81195, MN81205, MN81217
	<i>Lonchophylla dekeyseri</i>	MN81184, MN81190, MN81213
	<i>Desmodus rotundus</i>	MN78108, MN78109, MN78110, MN78111
	<i>Diphylla ecaudata</i>	MN78112, MN78113
	<i>Carollia brevicauda</i>	MN81197, MN81208, MN81219
	<i>Carollia perspicillata</i>	MN81191, MN81198, MN81210
	<i>Chrotopterus auritus</i>	MN78107
	<i>Glyphoncteris sylvestris</i>	MN78118
	<i>Micronycteris sanborni</i>	MN81185, MN81201
	<i>Micronycteris schmidtorum</i>	MN81206
	<i>Mimmon bennettii</i>	MN78119, MN78120
	<i>Phylloderma stenops</i>	MN81186, MN81189, MN81192
	<i>Phyllostomus discolor</i>	MN78387
	<i>Phyllostomus elongatus</i>	MN78388, MN78389, MN78390
	<i>Phyllostomus hastatus</i>	MN78383, MN78384, MN78385, MN78386
	<i>Trachops cirrhosus</i>	MN78122, MN78123
	<i>Artibeus planirostris</i>	MN78396, MN78397, MN78398
	<i>Chiroderma villosus</i>	MN78393
	<i>Platyrrhinus lineatus</i>	MN78394, MN78397
	<i>Sturnira lilium</i>	MN78391, MN78392
Furipteridae	<i>Furipterus horrens</i>	MN78114, MN78115, MN78116, MN78117
Noctilionidae	<i>Noctilio leporinus</i>	MN78124
Mormoopidae	<i>Pteronotus parnellii</i>	MN78121
Molossidae	<i>Molossops temminckii</i>	MN78399, MN78400, MN78401, MN78402
	<i>Molossus molossus</i>	MN81222
Vespertilionidae	<i>Eptesicus diminutus</i>	MN81182, MN81204
	<i>Eptesicus furinalis</i>	MN81207

Table A2. Caves compiled by literature review, including species richness of bats and author of the record.

Country	State/district/ department	Municipality	Cave	No. of species	Source
Brazil	Tocantins	Aurora do Tocantins	Gruta dos Moura	21	Present study
	Tocantins	Aurora do Tocantins	Gruta do Urso	11	Present study
	Goiás	Posse	Caverna dos Revolucionários	6	Esbérard et al. 2005
	Goiás	Posse	Caverna Asa Branca	8	Esbérard et al. 2005
	Goiás	Buritópolis	Gruta da Fazenda Bananal	8	Esbérard et al. 2005
	Goiás	Buritópolis	Gruña Meândrica	3	Esbérard et al. 2005
	Goiás	Buritópolis	Gruta da Judite	17	Esbérard et al. 2005
	Goiás	Buritópolis	Lapa da Fazenda Extrema	9	Esbérard et al. 2005
	Goiás	Buritópolis	Lapa da Lapa	11	Esbérard et al. 2005
	Goiás	Mambaí	Caverna Nova Esperança	6	Esbérard et al. 2005
	Goiás	Mambaí	Gruta da Fazenda Arroz	5	Esbérard et al. 2005
	Goiás	Mambaí	Caverna Ventura	8	Esbérard et al. 2005
	Goiás	Mambaí	Gruta Borá	4	Esbérard et al. 2005
	Goiás	Mambaí	Gruta do Landim	4	Esbérard et al. 2005
	Goiás	Damianópolis	Lapa da Fazenda Buritizinho	7	Esbérard et al. 2005
	Goiás	Padre Bernardo	Gruta Morro	13	Bredt et al. 1999
	Goiás	Padre Bernardo	Toca da Gameleira	8	Bredt et al. 1999
	Goiás	Padre Bernardo	Gruta das Orquídeas	4	Bredt et al. 1999
	Distrito Federal	Brazlândia	Gruta do Sal	8	Bredt et al. 1999
	Distrito Federal	Brazlândia	Gruta Fenda II	8	Bredt et al. 1999
	Distrito Federal	Brazlândia	Gruta da Barriguda	8	Bredt et al. 1999
	Distrito Federal	Brazlândia	Gruta Labirinto da Lama	5	Bredt et al. 1999
	Distrito Federal	Brazlândia	Gruta Dois Irmãos	9	Bredt et al. 1999

Continued

Table A2. Continued.

Country	State/district/ department	Municipality	Cave	No. of species	Source
	Distrito Federal	Brazlândia	Gruta Muralha	2	Bredt et al. 1999
	Distrito Federal	Sobradinho	Gruta da Saúva	7	Bredt et al. 1999
	Distrito Federal	Sobradinho	Gruta dos Morcegos	3	Bredt et al. 1999
	Distrito Federal	Sobradinho	Gruta Kipreste	3	Bredt et al. 1999
	Distrito Federal	Sobradinho	Gruta Mogi	7	Bredt et al. 1999
	Distrito Federal	Sobradinho	Gruta Boca de Lobo	1	Bredt et al. 1999
	Distrito Federal	Planaltina	Gruta Dança dos Vampiros	9	Bredt et al. 1999
	Distrito Federal	Planaltina	Gruta Água Rasa	8	Bredt et al. 1999
	Distrito Federal	Paranoá	Gruta Volks Club	8	Bredt et al. 1999
	Distrito Federal	Paranoá	Toca Mata da Anta	3	Bredt et al. 1999
	Distrito Federal	Ceilândia	Toca do Falcão	6	Bredt et al. 1999
	Distrito Federal	Ceilândia	Fenda do Barreiro	5	Bredt et al. 1999
	Minas Gerais	Itacarambi	Caverna Olhos d'Água	13	Trajano and Gimenez 1998
	Rio de Janeiro	Cantagalo	Caverna Santana	7	Esbérard et al. 1998
	São Paulo	Iporanga	Gruta Águas Quentes	4	Trajano 1985
	São Paulo	Iporanga	Caverna Água Suja	8	Trajano 1985
	São Paulo	Iporanga	Caverna Alambari de Baixo	5	Trajano 1985
	São Paulo	Iporanga	Gruta Alambari de Cima	1	Trajano 1985
	São Paulo	Iporanga	Complexo das Areias	4	Trajano 1985
	São Paulo	Iporanga	Toca Berta Funda	4	Trajano 1985
	São Paulo	Iporanga	Gruta do Betari	5	Trajano 1985
	São Paulo	Iporanga	Gruta do Chapéu	2	Trajano 1985
	São Paulo	Iporanga	Gruta do Córrego Seco	4	Trajano 1985
	São Paulo	Iporanga	Gruta do Grilo	3	Trajano 1985
	São Paulo	Iporanga	Abismo da Gurutuvai	3	Trajano 1985
	São Paulo	Iporanga	Abismo da Hipotenusai	3	Trajano 1985
	São Paulo	Iporanga	Gruta do Jeremias	2	Trajano 1985
	São Paulo	Iporanga	Gruta da Laje Branca	2	Trajano 1985
	São Paulo	Iporanga	Gruta do Macaco	2	Trajano 1985
	São Paulo	Iporanga	Gruta do Morro Preto	4	Trajano 1985
	São Paulo	Iporanga	Gruta do Couto	2	Trajano 1985
	São Paulo	Iporanga	Gruta Ouro Grosso	3	Trajano 1985
	São Paulo	Iporanga	Abismo da Passoca	2	Trajano 1985
	São Paulo	Iporanga	Caverna Santana	3	Trajano 1985
	São Paulo	Iporanga	Santana IV	4	Trajano 1985
	São Paulo	Iporanga	Gruta do Sítio Novo	2	Trajano 1985
	São Paulo	Iporanga	Sumidouro do David	2	Trajano 1985
	São Paulo	Iporanga	Gruta do Zezo	2	Trajano 1985
	São Paulo	Ribeira	Caverna Tiraprosa	5	Trajano 1985
	São Paulo	Ribeira	Toca do Porco	4	Trajano 1985
	São Paulo	Apiá	Gruta Calcáreo Branco	3	Trajano 1985
	São Paulo	Apiá	Gruta dos Vieira	4	Trajano 1985
	Paraná	Adrianópolis	Gruta São João	4	Trajano 1985
Mexico	Yucatán	-	Ruins of Kabah	1	Arita 1996
	Yucatán	-	Zorro Cave	1	Arita 1996
	Yucatán	-	Cave "A"	1	Arita 1996
	Yucatán	-	Actun Chac-Xix	1	Arita 1996
	Yucatán	-	Aguacate Cave	1	Arita 1996
	Yucatán	-	Actun Kan-Lol	1	Arita 1996
	Yucatán	-	Actun Dzontot	1	Arita 1996
	Yucatán	-	Actun Kab	1	Arita 1996
	Yucatán	-	Actun X-Maasit	1	Arita 1996
	Yucatán	-	Actun On	2	Arita 1996
	Yucatán	-	Porcupine Cave	2	Arita 1996
	Yucatán	-	La Doña Blanca Cave	2	Arita 1996
	Yucatán	-	Actun Tolok	2	Arita 1996
	Yucatán	-	Guayaba y Aguacate	2	Arita 1996
	Yucatán	-	Acanceh Cave	2	Arita 1996
	Yucatán	-	Iguana Cave	2	Arita 1996
	Yucatán	-	Actun Chunkunab	2	Arita 1996

Continued

Table A2. Continued.

Country	State/district/ department	Municipality	Cave	No. of species	Source
	Yucatán	-	Actun Maas	2	Arita 1996
	Yucatán	-	Kabahchen Cave	2	Arita 1996
	Yucatán	-	Actun Sitz	2	Arita 1996
	Yucatán	-	Chocantes Cave	2	Arita 1996
	Yucatán	-	Cave "B"	2	Arita 1996
	Yucatán	-	Bejucos Cave	3	Arita 1996
	Yucatán	-	Actun Oxpehol	3	Arita 1996
	Yucatán	-	Ramonal y Naranja	3	Arita 1996
	Yucatán	-	Roble Cave	4	Arita 1996
	Yucatán	-	Flor de Mayo Cave	4	Arita 1996
	Yucatán	-	Ramonal Cave	4	Arita 1996
	Yucatán	-	Ruins of Mayapan	4	Arita 1996
	Yucatán	-	Cinco de Mayo Cave	5	Arita 1996
	Yucatán	-	Bat Cave	7	Arita 1996
	Yucatán	-	Hoctun Cave	8	Arita 1996
	Yucatán	-	Actun Sabak-ha	8	Arita 1996
	Yucatán	-	Tzab-Nah Cave	9	Arita 1996
	Yucatán	-	Actun Spukil	9	Arita 1996
	Yucatán	-	Actun Lol-Tun	12	Arita 1996
	Puebla	-	La Escondida	1	Brunet and Medellín 2001
	Puebla	-	El Escorpion	1	Brunet and Medellín 2001
	Puebla	-	La Herradura	1	Brunet and Medellín 2001
	Puebla	-	Buenavista	no	Brunet and Medellín 2001
	Puebla	-	El Manantial	1	Brunet and Medellín 2001
	Puebla	-	El Tigre	1	Brunet and Medellín 2001
	Puebla	-	Mision Imposible	2	Brunet and Medellín 2001
	Puebla	-	La Salamandra	6	Brunet and Medellín 2001
	Puebla	-	San Martin	5	Brunet and Medellín 2001
	Puebla	-	El Arenal	4	Brunet and Medellín 2001
	Puebla	-	El Sapo	4	Brunet and Medellín 2001
	Puebla	-	Waterfall	1	Brunet and Medellín 2001
	Puebla	-	Nacho Sosa	2	Brunet and Medellín 2001
	Puebla	-	Los Copales	3	Brunet and Medellín 2001
	Puebla	-	La Ciudad	6	Brunet and Medellín 2001
	Puebla	-	El Coyote	3	Brunet and Medellín 2001
	Puebla	-	La Piramide	3	Brunet and Medellín 2001
	Puebla	-	Virgen	7	Brunet and Medellín 2001
	Puebla	-	El Tigre Grande	1	Brunet and Medellín 2001
	Puebla	-	Cuevade Las Vegas	11	Brunet and Medellín 2001
Honduras	Tegucigalpa	Masca	Piedra Cocha	3	Divoll and Buck 2013
	Tegucigalpa	San Juancito	Cueva1	5	Divoll and Buck 2013
Bolivia	Cochabamba	-	Repechón cave A	7	Siles et al. 2007
	Cochabamba	-	Repechón cave B	7	Siles et al. 2007
	Cochabamba	-	Repechón cave C	6	Siles et al. 2007
Venezuela	Paranaguá	Falcón	Cueva del Guano	5	Molinari et al. 2012
	Paranaguá	Falcón	Cuevade Piedra Honda	5	Molinari et al. 2012
	Paranaguá	Falcón	Cuevade Jacuque	4	Molinari et al. 2012
	Paranaguá	Los Taques	Cueva N° 2 del Pico	4	Molinari et al. 2012
Cuba	La Habana	San José de Las Lajas	Cuevadel Indio	6	Mora et al. 2002.
Colombia	Santander	Curití	Macaragua cave	10	Pérez-Torres et al. 2015
	Antioquia	Río Claro	Caverna de Ilusiones IIIb	1	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Corinto	4	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Ilusiones IIIa	3	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Ilusiones IV	5	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Tambores	5	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Chontaduro	1	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Mauro I	4	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	Caverna Mauro II	2	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	El Condor	3	Muñoz-Saba et al. 1999
	Antioquia	Río Claro	La Danta	2	Muñoz-Saba et al. 1999