



# List of *Culicoides* biting midges (Diptera: Ceratopogonidae) from the state of Amazonas, Brazil, including new records

Emanuelle de Sousa Farias<sup>1,2</sup>, Jessica Feijó Almeida<sup>1,3</sup> and Felipe Arley Costa Pessoa<sup>1\*</sup>

1 Instituto Leônidas e Maria Deane, Fiocruz Amazônia, Laboratório de Ecologia de Doenças Transmissíveis na Amazônia, Rua Terezina, N° 476, Adrianópolis, CEP 69.057-070, Manaus, AM, Brazil

2 Programa de Pós-graduação em Biodiversidade e Saúde, Instituto Oswaldo Cruz (IOC/FIOCRUZ)

3 Programa de Pós-graduação em Condições de Vida e Situações de Saúde na Amazônia, Instituto Leônidas e Maria Deane (ILMD/FIOCRUZ)

\* Corresponding author. E-mail: [felipe.pessoa@fiocruz.br](mailto:felipe.pessoa@fiocruz.br)

**Abstract:** *Culicoides* are vectors of pathogenic agents that infect humans and other animals. Here, we provide a list of *Culicoides* from the state of Amazonas and also document new records from Presidente Figueiredo Municipality, Amazonas, Brazil. We provide a map of recorded species and a wing atlas for identification. The *Culicoides* fauna of Amazonas is now known to include 89 known species that belong to seven subgenera, 10 informal species groups, and one ungrouped species. We record nine species of *Culicoides* (*C. aldomari*, *C. batesi*, *C. brownie*, *C. flavivenulus*, *C. franklini*, *C. guamai*, *C. paramaruim*, *C. pusilloides* and *C. tidwelli*) for the first time from Amazonas state. *Culicoides brownie* and *C. tidwelli* are reported for the first time from Brazil.

**Key words:** biodiversity, culicomorpha, hematophagous insects, Amazon Basin

## INTRODUCTION

*Culicoides* Latreille, 1809, or biting midges, are known vectors of pathogenic nematodes, protozoa, and viruses that cause infection in humans and other animals. In addition to their vectorial importance, these midges are pernicious due to their bites (Mellor et al. 2000). There are 1,355 species described worldwide (Borkent 2015); 116 species inhabit Brazil and 80 have been recorded in the state of Amazonas (Farias et al. 2016; Santarém and Felipe-Bauer 2016). A survey conducted in Presidente Figueiredo Municipality recorded species previously unknown in the state of Amazonas. Here, we revise the list *Culicoides* species occurring in Amazonas state and record the nine new species records for the state, and two new records for Brazil.

## MATERIAL AND METHODS

We compiled a list of *Culicoides* species (Table 1),

and mapped each species in Amazonas state, based on Borkent and Spinelli (2007), Farias et al. (2016), Castellón and Felipe-Bauer (2015), and Santarém and Felipe-Bauer (2016). The municipalities in Amazonas state where there are records of *Culicoides* are mapped in Figure 1. We collected additional records during an entomological study in a rural settlement of Rio Pardo, Presidente Figueiredo Municipality, Amazonas, Brazil. This survey was carried out between June and August 2010, using CDC light traps in forest fragments and peridomiciles (shelters for domestic animals and cultivation). Specimens of *Culicoides* were preserved in 70% ethanol, dissected, and slide-mounted in Canada balsam using the techniques described by Wirth and Marston (1968). We identified our specimens using various identification keys, as follows: Wirth and Blanton (1959); Spinelli et al. (1993); Felipe-Bauer et al. (2013); and Santarém et al. (2015). Subgeneric classification of *Culicoides* species was based on Borkent and Spinelli (2007) and Borkent (2015). Some voucher specimens are deposited at the Laboratory of Infectious Disease Ecology in Amazon (ILMD/FIOCRUZ, Amazônia; Table 1). Microphotographs of the wings were obtained using a digital system adapted to an optical microscope with a digital camera.

## RESULTS

The *Culicoides* fauna of Amazonas state is comprised of 89 species belonging to seven subgenera, 10 informal species groups, and one ungrouped species (Table 1). In our study, *C. brownei* Spinelli, 1993 and *C. tidwelli* Spinelli, 1993 are recorded for the first time in Brazil. Another seven species were found for the first time in Amazonas state, as follows: *C. aldomari* Felipe-Bauer & Trindade, 2013; *C. batesi* Wirth & Blanton, 1973; *C. flavivenulus* Costa Lima, 1937; *C. franklini* Spinelli, Greiner & Wirth, 1993; *C. guamai* Wirth & Blanton, 1973; *C.*

**Table 1.** *Culicoides* species recorded to Amazonas state, Brazil.

| Subgenus/group                   | Species                   | Municipalities /Coordinates                                                                                                                                                                                                                                                                                                                                                                                                  | Voucher number |
|----------------------------------|---------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
| <i>Anilomyia / covagarciai</i>   | <i>Culicoides efferus</i> | Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W)                                                                                                                                                                                                                                                                                                                                                                              | ***            |
| <i>Avaritia</i>                  | <i>C. pusilloides</i> *   | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-8         |
|                                  | <i>C. pusillus</i>        | Manacapuru (03°23'46.71" S, 060°37'56.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Santa Isabel do Rio Negro (00°24'49.74" S, 65°01'00.60" W)                                                                                                                                                                                                                                                            | 0001-10        |
| <i>Diphaomyia</i>                | <i>C. freitasi</i>        | Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W)                                                                                                                                                                                                                                                                                                                                                                              | ***            |
| <i>Haematomyidium</i>            | <i>C. crucifer</i>        | Manaus (03° 5'39.60" S, 059°59'14.09" W)                                                                                                                                                                                                                                                                                                                                                                                     | ***            |
|                                  | <i>C. debilipalpis</i>    | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Novo Airão (02°42'43.17" S, 060°56'48.13" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W), Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W)                                                             | 0001-11        |
|                                  | <i>C. denisae</i>         | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                     | ***            |
|                                  | <i>C. glabrior</i>        | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                                                 | 0001-12        |
|                                  | <i>C. hoffmani</i>        | ***                                                                                                                                                                                                                                                                                                                                                                                                                          | ***            |
|                                  | <i>C. insinuatus</i>      | Barcelos (00°36'59.92" S, 062°55'54.82" W); Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Manacapuru (03°23'46.71" S, 060°37'56.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Santo Antônio do Içá (03°02'39.40" S, 069°05'49.03" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W)                                                                                            | 0004-1         |
|                                  | <i>C. spurius</i>         | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                        | 0005-2         |
|                                  | <i>C. todatangae</i>      | Manacapuru (03°23'46.71" S, 060°37'56.84" W), Manaus (03°00'32.04" S, 059°56'49.84" W)                                                                                                                                                                                                                                                                                                                                       | 0005-1         |
|                                  | <i>C. youngi</i>          | ***                                                                                                                                                                                                                                                                                                                                                                                                                          | ***            |
|                                  | <i>C. neoparaensis</i>    | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                     | ***            |
| <i>Haematomyidium/ paraensis</i> | <i>C. paraensis</i>       | Barcelos (00°36'59.92" S, 062°55'54.82" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" S, 067°35'36.0" W), Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W)                                                                                                            | 0001-13        |
|                                  | <i>C. quasiparaensis</i>  | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W)                                                                                                                                                                                                                                              | 0005-6         |
|                                  | <i>C. batesi</i> *        | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-2         |
| <i>Hoffmania /guttatus</i>       | <i>C. bimaculatus</i>     | ***                                                                                                                                                                                                                                                                                                                                                                                                                          | ***            |
|                                  | <i>C. brasilianum</i>     | ***                                                                                                                                                                                                                                                                                                                                                                                                                          | ***            |
|                                  | <i>C. brownei</i> **      | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-3         |
|                                  | <i>C. coutinhoi</i>       | Manaus (03°03'00.97" S, 060°01'29.92" W), Parintins (02°38'15.99" S, 056°43'46.29" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                               | 0001-13        |
|                                  | <i>C. diabolicus</i>      | Coari (04°10'50.95" S, 063°09'30.77" W), Manacapuru (03°23'46.71" S, 060°37'56.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" S, 67°35'36.0" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W)                                                                                                                        | 0001-14        |
|                                  | <i>C. filariferus</i>     | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Rio Preto da Eva (02°41'55.06" S, 059°42'01.01" W)                                                                                                                                                                                                                                                                                                                  | 0001-15        |
|                                  | <i>C. flavivenulus</i> *  | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-4         |
|                                  | <i>C. foxi</i>            | Barcelos (00°36'59.92" S, 062°55'54.82" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" S, 067°35'36.0" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                      | 0001-16        |
|                                  | <i>C. franklini</i> *     | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-5         |
|                                  | <i>C. fusipalpis</i>      | Barcelos (00°36'59.92" S, 062°55'54.82" W), Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                      | 0001-17        |
|                                  | <i>C. ignacioi</i>        | Manaus (03°03'00.97" S, 060°01'29.92" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                           | 0001-18        |
|                                  | <i>C. insignis</i>        | Barcelos (00°36'59.92" S, 062°55'54.82" W), Coari (02°13'14.38" S, 063°49'40.72" W), Manacapuru (03°23'46.71" S, 060°37'56.84" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Santo Antônio do Içá (03°02'39.40" S, 069°05'49.03" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Tabatinga (04°13'53.83" S, 069°56'12.98" W), Tefé (03°21'05" S, 064°42'53" W) | 0001-19        |
|                                  | <i>C. lutzi</i>           | Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Santo Antônio do Içá (03°02'39.40" S, 069°05'49.03" W)                                                                                                                                                                                                                                                                    | 0001-20        |
|                                  | <i>C. ocumarensis</i>     | Benjamin Constant (04°24'51.48" S, 069°39'33.05" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                 | 0001-21        |
|                                  | <i>C. paraignacioi</i>    | Manaus (02°45'31.15" S, 061°00'17.82" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                             | 0001-22        |
|                                  | <i>C. paramaruim</i> *    | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-7         |
|                                  | <i>C. plaumanni</i>       | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                      | 0001-23        |

Continued

Table 1. Continued.

| Subgenus/group                | Species                    | Municipalities /Coordinates                                                                                                                                                                                                                                                                                                                                                                                                                    | Voucher number |
|-------------------------------|----------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
|                               | <i>C. pseudodiabolicus</i> | Barcelos (00°36'59.92" S, 062°55'54.82" W), Coari (02°13'14.38" S, 063°49'40.72" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Novo Airão (02°42'43.17" S, 060°56'48.13" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W), Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W), Tefé (03°21'05" S, 064°42'53" W) | 0001-24        |
|                               | <i>C. ruizi</i>            | Manacapuru (03°23'46.71" S, 060°37'56.84" W)                                                                                                                                                                                                                                                                                                                                                                                                   | ***            |
|                               | <i>C. tidwelli</i> **      | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                                        | 0001-9         |
|                               | <i>C. travassosi</i>       | ***                                                                                                                                                                                                                                                                                                                                                                                                                                            | ***            |
| <i>Hoffmania / hylas</i>      | <i>C. aitkeni</i>          | Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                                                                                                                                                                                                               | 0002-2         |
|                               | <i>C. baniwa</i>           | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                                                              | 0001-25        |
|                               | <i>C. heliconiae</i>       | Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W)                                                                                                                                                                                                                                                                                                                                                                                    | ***            |
|                               | <i>C. hylas</i>            | Barcelos (00°36'59.92" S, 062°55'54.82" W), Coari (02°13'14.38" S, 063°49'40.72" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Airão (05°14'19.59" S, 060°57'03.77" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W), Santa Isabel do Rio Negro (00°24'49.74" S, 65°01'00.60" W), Tefé (03°21'05" S, 064°42'53" W)                                                   | 0001-26        |
|                               | <i>C. palpalis</i>         | ***                                                                                                                                                                                                                                                                                                                                                                                                                                            | ***            |
|                               | <i>C. polyptori</i>        | São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. pseudoheliconiae</i> | Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                                                                                                                                                                                                               | ***            |
|                               | <i>C. verecundus</i>       | Barcelos (00°36'59.92" S, 062°55'54.82" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                   | 0001-27        |
| <i>Mataemyia</i>              | <i>C. aldomari</i> *       | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                                        | 0001-1         |
|                               | <i>C. bricenoi</i>         | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                     | 0001-27        |
|                               | <i>C. discrepans</i>       | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. felippebaueri</i>    | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. wallacei</i>         | Benjamin Cosntant (04°24'51.48" S, 069°39'33.05" W), Coari (02°13'14.38" S, 063°49'40.72" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                         | ***            |
| <i>Oecacta</i>                | <i>C. alahialinus</i>      | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
| Unplaced / <i>acotylus</i>    | <i>C. acotylus</i>         | Presidente Figueiredo (01°55'05.53" S, 059°28'30.23" W)                                                                                                                                                                                                                                                                                                                                                                                        | ***            |
| Unplaced / <i>carpenteri</i>  | <i>C. belemensis</i>       | Manaus (02°45'31.15" S, 061°00'17.82" W)                                                                                                                                                                                                                                                                                                                                                                                                       | 0002-1         |
|                               | <i>C. camposi</i>          | Barcelos (00°36'59.92" S, 062°55'54.82" W)                                                                                                                                                                                                                                                                                                                                                                                                     | ***            |
|                               | <i>C. carpenteri</i>       | Fonte Boa (02°48'01.79" S, 066°18'47.08" W), Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                          | ***            |
| Unplaced / <i>dasyophrus</i>  | <i>C. dasyophrus</i>       | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                     | 0001-28        |
| Unplaced / <i>eublepharus</i> | <i>C. eublepharus</i>      | Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                              | 0001-29        |
|                               | <i>C. propriipennis</i>    | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°03'00.97" S, 060°01'29.92" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W) & São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                          | 0001-30        |
|                               | <i>C. rangeli</i>          | Santa Isabel do Rio Negro (00°24'49.74" S, 065°01'00.60" W)                                                                                                                                                                                                                                                                                                                                                                                    | ***            |
| Unplaced / <i>fluvialis</i>   | <i>C. fluvialis</i>        | Manaus (03°03'00.97" S, 060°01'29.92" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                              | 0001-31        |
|                               | <i>C. leopoldoi</i>        | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                                                                   | 0001-32        |
|                               | <i>C. tetrathyris</i>      | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°03'00.97" S, 060°01'29.92" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                     | 0001-33        |
| Unplaced / <i>leoni</i>       | <i>C. benarrochi</i>       | Manaus (03°03'00.97" S, 060°01'29.92" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                    | 0001-34        |
|                               | <i>C. fieldi</i>           | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. glabellus</i>        | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), Tefé (03°21'05" S, 064°42'53" W)                                                                                                                                                                                                                                                                                                                                                      | 0001-35        |
|                               | <i>C. leoni</i>            | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
| Unplaced / <i>limai</i>       | <i>C. antunesi</i>         | ***                                                                                                                                                                                                                                                                                                                                                                                                                                            | ***            |
|                               | <i>C. galindoi</i>         | Manacapuru (03°23'46.71" S, 060°37'56.84" W), Manaus (03°03'00.97" S, 060°01'29.92" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                               | 0004-2         |
|                               | <i>C. limai</i>            | Manaus (03°03'00.97" S, 060°01'29.92" W), Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                    | 0001-36        |
|                               | <i>C. lopesi</i>           | Manacapuru (03°23'46.71" S, 060°37'56.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W)                                                                                                                                                                                                                                                                                                                                                  | 0004-3         |
|                               | <i>C. santanderi</i>       | São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. vernoni</i>          | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                                        | 0005-3         |
| Unplaced / <i>pachymerus</i>  | <i>C. caprilesi</i>        | São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                                                                                                                       | ***            |
|                               | <i>C. pachymerus</i>       | Manaus (03°03'00.97" S, 060°01'29.92" W), São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                                                                                                                                                                                                             | ***            |
| Unplaced / <i>reticulatus</i> | <i>C. aureus</i>           | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                                                                                                                                                                                                                                                       | 0005-4         |
|                               | <i>C. castelloni</i>       | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                                                                                                                                                                                                                                                        | CCER           |
|                               | <i>C. fittkaui</i>         | Coari (04°10'50.95" S, 063°09'30.77" W)                                                                                                                                                                                                                                                                                                                                                                                                        | ***            |

Continued

Table 1. Continued.

| Subgenus/group              | Species                     | Municipalities /Coordinates                                                                                                                                                                                        | Voucher number |
|-----------------------------|-----------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------|
|                             | <i>C. guamai</i> *          | Presidente Figueiredo (31°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-37        |
|                             | <i>C. hildebrandoi</i>      | Presidente Figueiredo (01°55'05.53" S, 059°28'30.23" W)                                                                                                                                                            | 0003-1         |
|                             | <i>C. irregularis</i>       | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-38        |
|                             | <i>C. kuripako</i>          | São Gabriel da Cachoeira (01°03'44.5" N, 067°35'36.0" W)                                                                                                                                                           | ***            |
|                             | <i>C. lanei</i>             | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                           | ***            |
|                             | <i>C. paucienfuscatus</i>   | Coari (04°10'50.95" S, 063°09'30.77" W), Manaus (03°00'32.04" S, 059°56'49.84" W), Novo Aripuanã (05°26'03.28" S, 060°57'05.70" W), Novo Airão (02°42'43.17" S, 060°56'48.13" W), Tefé (03°21'05" S, 064°42'53" W) | 0005-5         |
|                             | <i>C. profundus</i>         | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-39        |
|                             | <i>C. pseudoreticulatus</i> | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-40        |
|                             | <i>C. rhombus</i>           | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-41        |
| Unplaced / <i>stigmalis</i> | <i>C. fluviatilis</i>       | Presidente Figueiredo (01°48'01.66" S, 060°18'59.82" W)                                                                                                                                                            | 0001-42        |
| Unplaced / ungrouped        | <i>C. wokei</i>             | Manaus (03°03'00.97" S, 060°01'29.92" W)                                                                                                                                                                           | ***            |

\* New records in Amazon State; \*\* new records in Brazil; \*\*\*Specific location & voucher number not found, occurrence done only to state.

*paramaruim* Wirth & Blanton, 1973; and *C. pusilloides* Wirth & Blanton, 1955.

### Species composition

Subgenus *Anilomyia* Vargas, 1960

*Anilomyia* Vargas (1960): 37, as subgenus of *Culicoides*. Type species: *Culicoides covagarciai* Ortiz, by original designation.

***Culicoides efferus*** Fox, 1952

(Figures 2, 10A)

*Culicoides efferus* Fox (1952): 365. Peru

**Identification:** Eyes contiguous, bare; third palpal segment long and slender with small sensorial pit; wing as figured, with very restricted dark spots, costa extending to 0.67 of wing length, macrotrichia very scanty on distal fourth of wing; halter whitish; spermathecae two, pyriform, subequal. Male with ninth sternum without posterior excavation; ninth tergum rounded caudad with only faint vestiges of apicolateral processes, deeply cleft mesad at apex with a pair of pointed; aedeagus narrow basally, anterior sclerotized membrane covering slightly more than distal half of basal arch, distal peg and apical spherical tip very slender; parameres fused more than halfway to apices, the free portions very slender, pointed, hairy, the common basal portion (Wirth and Blanton 1959).

**Distribution:** Bolivia, Brazil (Acre, Amazonas, and Pará); Guatemala to Peru.

Subgenus *Avaritia* Fox, 1955

*Avaritia* Fox (1955): 218, as subgenus of *Culicoides*. Type species: *Ceratopogon obsoletus* Meigen, by original designation.

**\**Culicoides pusilloides*** Wirth & Blanton, 1955

(Figures 2, 10B)

*Culicoides pusilloides* Wirth and Blanton (1955): 104. Panama.

**Identification:** Medium-sized, dark brown; wing as figured; legs brown with pale bands; third segment palpus pit deep; spermathecae subspherical and very unequal. This species is differs from *C. pusillus*, from which it can be distinguished by second radial cell distinctly pale on

the apical half, while in *C. pusillus* the second radial cell is pale only on extreme distal (Wirth and Blanton 1959).

**Distribution:** Belize to Panama, Brazil (Acre and Amazonas) and Guatemala

**New records:** Brazil, Amazonas State, Presidente Figueiredo, Rio Pardo (01°48'S, 060°19' W), VII-2010, 0001-8 (1 female), CDC light trap, peridomicilies, F.A.C. Pessoa collector.

***Culicoides pusillus*** Lutz, 1913

(Figures 2, 10C)

*Culicoides pusillus* Lutz (1913): 52. Rio de Janeiro, Brazil.

**Identification:** Small size, bare; wing as shown in figure, without marginal pale spots, the very short, entirely dark second radial cell, blackish mesonotum with two faint darker vittae and relatively pale. This species is an exception among the related species of the subgenus *Culicoides* (*Avaritia*) due to the apex of the very short second radial cell entirely dark or only very slightly encroached on by the poststigmatic pale spot (Wirth and Blanton 1959).

**Distribution:** Florida, USA to Central and South America to northeastern Argentina and Brazil (Acre, Amapá, Amazonas, Pará, Bahia, Minas Gerais, Rio de Janeiro, São Paulo, and Santa Catarina).

Subgenus *Diphaomyia* Vargas, 1960

*Diphaomyia* Vargas (1960): 40. Type species: *Culicoides baueri* Hoffman, by original designation.


***Culicoides freitasi*** Wirth & Blanton, 1973

(Figures 2, 10D)

*Culicoides freitasi* Wirth and Blanton (1973): 434. -Pará, Brazil.

**Identification:** Eyes narrowly separated, bare; third palpal segment moderately swollen, with a round deep sensory pit; wing as figured and similar in pattern to *C. iriatei* Fox 1952, but differs due to two pale spots in the distal portion of the anal cell and only pale spot lying adjacent to the proximal portion of vein M<sub>1</sub>; halter pale; two spermathecae (Wirth and Blanton 1973).

**Distribution:** Brazil (Amazonas and Pará).



**Figure 1.** Maps showing the position of Amazonas state in Brazil and South America. Large image shows the municipalities where there were records of *Culicoides* (modified after SimpleMappr, <http://www.simplemappr.net>).

#### Subgenus *Haematomyidium* Goeldi, 1905

*Haematomyidium* Goeldi (1905): 137. Type species: *Haematomyidium paraensis* Goeldi, by original designation.

#### *Culicoides crucifer* Clastrier, 1968

(Figure 2)

*Culicoides crucifer* Clastrier (1968): 85. French Guiana.

**Identification:** A large species; third palpal segment slender with broad shallow pit; wing with distal pale spot in cell  $m_1$  lying far from wing margin, no pale spot present in cell  $m_2$  lying adjacent to midportion of mediocubital stem; aedeagus conspicuously cleft distad with two sharp laterally directed teeth on each side; parameres with simple twisted tips without fringing spines (Vitale et al. 1981).

**Distribution:** Brazil (Amazonas, Pará), French Guiana, Guyana, and Trinidad.

#### *Culicoides debilipalpis* Lutz, 1913

(Figures 2, 10E)

*Culicoides debilipalpis* Lutz (1913): 60. São Paulo, Brazil.

*Culicoides khalafi* Beck (1957): 104. Florida, USA.

*Culicoides ichesi* Ronderos and Spinelli (1995): 77. Argentina.

**Identification:** Third palpal segment stout, PR 2.00–2.60; flagellomere 8 longer than 9; wing as shown in figure, with distal pale spot in cell  $m_1$  separated from wing margin by a distance approximate to its length, macrotrichia distributed on distal two-thirds of wing, extending in at least two rows to base of cell  $m_2$ ; halter brown (Spinelli et al. 2005).

**Distribution:** USA (Maryland, Kentucky, and Nebraska, south to Louisiana and Florida), Guatemala and Belize to Argentina and Brazil (Acre, Amapá, Amazonas, Ceará, Goiás, Espírito Santo, Mato Grosso do Sul, Pará, Pernambuco, São Paulo, Santa Catarina, and Roraima).

#### *Culicoides denisae* Clastrier, 1971

(Figure 2)

*Culicoides denisae* Clastrier (1971): 290. French Guiana.

**Identification:** Eyes narrowly separated; third palpal segment moderately swollen, PR 2.0; AR 0.9; wing with distal light spot transverse in  $r_3$ ,  $r-m$  with light spot, presence of two bright spots in cell  $m_1$ , the second spot not touch the edge of the wing, without clear area on one-third of vein  $M_2$ , apices of veins  $M_1$  and  $M_2$  usually clear; light spot more or less confluent in anal cell. (Waller et al. 1990).

**Distribution:** Brazil (Amazonas, Pará) and French Guiana.

#### *Culicoides glabrior* Macfie, 1940

(Figures 2, 10F)

*Culicoides debilipalpis* var. *glabrior* Macfie (1940): 27. Guyana.

*Culicoides grahambelli* Forattini (1956a): 35. Panama.


**Identification:** Eyes bare, broadly contiguous; third segment broad, with a very broad, shallow sensory pit; fourth segment much broader than fifth; wing with transverse distal spot in cell  $r_3$ , only one distal spot in cell  $m_2$ , pale spots present behind medial fork and in front of mediocubital fork, a single transverse pale spot in apex of anal cell, extending nearly to wing margin; halter brown, the end of the knob slightly paler; spermathecae two, pyriform, subequal (Wirth and Blanton 1959).

**Distribution:** Brazil (Acre, Amazonas, Pará, and Roraima), Guyana, Honduras to Ecuador, Suriname, and Trinidad

#### *Culicoides hoffmani* Fox, 1946

*Culicoides hoffmani* Fox (1946): 251. Trinidad.

**Identification:** Eyes separated, the line of separation broad above but narrowed below, short interfacetal hairs; third segment very short and swollen, with a moderately large and deep sensory pit; distal pale spot in cell  $r_3$  moderately large, rounded; two pale spots in cell  $m_1$ ; one pale spot each in apices of cells  $m_2$ ,  $CuA_1$  and anal cell; a



**Figure 2.** Map of the distribution of *Culicoides* species in Amazonas state, Brazil. **A.** *Culicoides efferus*. **B.** *C. pusilloides*. **C.** *C. pusillus*. **D.** *C. freitasi*. **E.** *C. crucifer*. **F.** *C. debilipalpis*. **G.** *C. denisae*. **H.** *C. glabrior*. **I.** *C. insinuatus*. **J.** *C. spurius* (modified after SimpleMappr, <http://www.simplemappr.net>).

pale spot lying ahead of mediocubital fork and another pale spot lying behind medial fork; halter brown, the flat end of the knob whitish; spermathecae two, pyriform, slightly unequal (Wirth and Blanton 1959).

**Distribution:** Brazil (Amazonas), Belize, Costa Rica, Cayman Islands, and Jamaica to Trinidad.

***Culicoides insinuatus*** Ortiz & León, 1955

(Figures 2, 10G)

*Culicoides insinuatus* Ortiz and León (1955): 577. Ecuador.

**Identification:** Eyes broadly separated; third palpal segment broader with shallow sensory pit; wing as figured, much shorter and broader with the pale spot grayish rather than whitish, the halter dark brownish, a distinct pale spot present in front of mediocubital fork; distinct ventral lobe present on the male parameres (Wirth and Blanton 1973, in *C. todatangae*).

**Distribution:** Brazil (Acre, Amapá, Amazonas, Pará,

Roraima and São Paulo), Colombia, Ecuador, Guyana, Peru, and Trinidad.

***Culicoides spurius*** Wirth & Blanton, 1959

(Figures 2, 10H)

*Culicoides spurius* Wirth and Blanton (1959): 433. Panama.

**Identification:** Eyes broadly separated, third segment very short and swollen, with a large, deep sensory pit; wing with pale spot over r-m crossvein about as broad as long, poststigmatic pale spots in cell  $r_3$  small and rounded, well separated, the posterior one slightly larger and located slightly proximad of the anterior one; distal pale spot in cell  $r_3$  moderately large, rounded; two pale spots in cell  $m_1$ ,  $CuA_1$  and anal cell; definite pale spots present behind medial fork and in front of mediocubital fork (Wirth and Blanton 1959).

**Distribution:** Brazil (Acre, Amazonas and Roraima), Costa Rica, and Panama.

***Culicoides todatangae* Wirth & Blanton, 1973**

(Figures 3, 10I)

*Culicoides todatangae* Wirth and Blanton (1973): 447. Pará, Brazil.

**Identification:** Eyes narrowly separated, with long interfacetal hairs; third palpal segment long and slender, with a small sensory pit deeper than diameter of pore opening; wing as shown in figure; halter slightly infuscated; spermathecae two, oval with long slender necks, slightly unequal; halter slightly infuscated. This species resembles *C. insinuatus* both species having the pale sport in cell  $m_2$  in front of mid-portion of the mediocubital stem (Wirth and Blanton 1973).

**Distribution:** Brazil (Acre, Amapá, Amazonas, Pará, and Santa Catarina) and Trinidad.

***Culicoides youngi* Wirth & Barreto, 1978***Culicoides youngi* Wirth and Barreto (1978): 562. Colombia.

**Identification:** Eyes nearly contiguous; third segment short and swollen to tip, with broad, round, shallow, sensory pit opening by a slightly smaller pore; second radial cell dark to tip with moderately broad lumen, slightly tapering distally, pale spot over r-m crossvein transverse, poststigmatic pale spot in cell  $r_3$  not divided, continued and expanded around posterior margin of second radial cell; distal pale spot presente in cell  $r_3$ , cell  $m_1$  with two pale spots, cell  $m_2$  with distinct pale spot lying behind medial fork and indistinct spot in front of mediocubital fork and a pale spot at apex of cell near wing margin, cell  $CuA_1$  with large rounded pale spot in distal portion, anal cell with pale spot in distal portion, dark at base; vein  $M_1$  with distinct pale spot at wing margin (Wirth and Barreto 1978).

**Distribution:** Brazil (Amazonas) and Colombia.

***Culicoides neoparaensis* Tavares & Souza, 1978**

(Figure 3)

*Culicoides neoparaensis* Tavares and Souza (1978): 621. Rio de Janeiro, Brazil.

**Identification:** Wing with three pale spots in cell  $m_1$ , distal pale spot distinctly separated from the subapical pale spot; third segment palpus short and stout (PR 1.7); sclerotized ring short; third rudimentary spermatheca greatly elongate. Male: paramere with definite broad ventral swelling, without elongate lobe (Felippe-Bauer et al. 2003).

**Distribution:** Brazil (Amazonas, Rio de Janeiro, and Santa Catarina).

***Culicoides paraensis* (Goeldi, 1905)**

(Figures 3, 10J)

*Haematomyidium paraensis* Goeldi (1905): 137. Pará, Brazil.*Culicoides undecimpunctatus* Kieffer (1917): 307. Argentina.

**Identification:** Third segment palpus longer and more slender (PR 2.1–2.8); wing with three pale spots in cell  $m_1$ , distal pale spot distinctly separated from the subapical pale spot; third rudimentary spermatheca shorter;

sclerotized ring long and curved; male with paramere uniformly slender in midportion, with elongate lobe (Felippe-Bauer et al. 2003).

**Distribution:** USA to Argentina and Brazil (Acre, Amazonas, Pará, Roraima, Bahia, Ceará, Maranhão, Pernambuco, Mato Grosso do Sul, Espírito Santo, Minas Gerais, Rio de Janeiro, São Paulo, and Santa Catarina).

***Culicoides quasiparaensis* Clastrier, 1971**

(Figures 3, 10K)

*Culicoides quasiparaensis* Clastrier (1971): 286. French Guiana.

**Identification:** Eyes narrowly separated above, contiguous below, the interocular space wedge-shaped; third palpal segment broad (PR 1.7) with pit broad and shallow, rarely deep and opening by a smaller pore; wing as shown in figure with three pale spots in cell  $m_1$ ; distal pale spot distinctly separated from the subapical pale spot; spermathecae very unequal sized (Felippe-Bauer et al. 2003).

**Distribution:** Honduras, and El Salvador, south to Colombia, Peru, French Guiana, and Brazil (Acre, Amazonas, Pará, Rondônia, Roraima, and Maranhão).

**Subgenus *Hoffmania* Fox, 1948**

*Hoffmania* Fox 1948: 21 (as subgenus of *Culicoides*). Type species: *Culicoides inamollae* Fox & Hoffman, 1944 (= *Culicoides insignis* Lutz, 1913, by original designation).

***Culicoides aitkeni* Wirth & Blanton, 1968**

(Figures 5)

*Culicoides aitkeni* Wirth and Blanton (1968): 214. Trinidad and Tobago.

**Identification:** Third palpal segment with scattered sensilla; wing length 1.50 mm;  $r_3$  with pale spot present anterior to base of  $M_1$  and in apices of  $CuA_1$  and  $CuA_2$ , a single pale spot crossing second radial cell; mid and hind femur with subapical pale band; spermathecae with short, slender necks. (Felippe-Bauer et al. 2009).

**Distribution:** Brazil (Amazonas, Pará); Trinidad and Tobago.

***Culicoides baniwa* Felippe-Bauer, 2009**

(Figure 5, 10L)

*Culicoides baniwa* Felippe-Bauer in Felippe-Bauer et al. (2009): 852. Amazonas, Brazil.

**Identification:** This species is easily distinguished from the other species of the *hylas* group by the presence of double pale spots in  $r_3$ , one crossing second radial cell and the other in distal portion of  $r_3$ , by the absence of pale spot in front of base of  $M_1$  and by the mid femur dark to tip (Felippe-Bauer et al. 2009).


**Distribution:** Brazil (Amazonas) and Peru.

**\**Culicoides batesi* Wirth & Blanton, 1973**

(Figures 3, 10M)

*Culicoides batesi* Wirth and Blanton (1973): 426. Brazil -- Pará, Brazil.*Culicoides sanmartini* Wirth and Barreto (1978): 553. Colombia.

**Identification:** This species is small to medium-sized,



**Figure 3.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. totatanga*. **B.** *C. neoparaensis*. **C.** *C. paraensis*. **D.** *C. quasiparaensis*. **E.** *C. batesi*. **F.** *C. brownei*. **G.** *C. coutinhoi*. **H.** *C. diabolicus*. **I.** *C. filariferus*. **J.** *C. flavivenulus* (modified after SimpleMappr, <http://www.simplemappr.net>).

dark brown; eyes contiguous by a distance equal to diameter of 2–3.5 ommatidial facets; third segment palpus moderately stout, sensory pit irregular. *Culicoides batesi* is very similar to *C. lutzi*, from which it can be distinguished by the faintly darkened r-m crossvein, dark apex of vein CuA<sub>1</sub> (Broadly pale em *C. lutzi*); macrotrichia present on distal one-third of wing, and dark halter (Spinelli et al. 1993).

**Distribution:** Bolivia, Colombia, Ecuador, Brazil (Amazonas, Pará, and Rondônia), and Guatemala

**New records:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VI-2010, VII-2010, VIII-2010, 0001-2 (4 females), CDC light traps, forest fragments and peridomiciles, F.A.C. Pessoa collector.

***Culicoides bimaculatus* Floch & Abonnenc, 1942**

*Culicoides bimaculatus* Floch and Abonnenc (1942a): 3. French Guiana.

**Identification:** Eyes contiguous by a distance equal

to the diameter of 2 ommatidial facets; wing with r-m crossvein pale, one transverse distal pale spot in cell r<sub>3</sub> broadly reaching wing margin, two distal pale spots in cell m<sub>1</sub> (the second one very faint in two of the available specimens), apices of veins M<sub>1</sub>, M<sub>2</sub>, CuA<sub>1</sub> pale, apex of vein CuA<sub>2</sub> dark; spermathecae small, spherical, subequal with short necks. This species resembles *C. foxi* by virtue of the presence of a small blackish spot behind the apex of the second radial cell, and by the dark brown halter. Also, it can be distinguished from *C. foxi* by the stouter third palpal segment with irregular sensory pit, and by the pale r-m crossvein (Spinelli et al. 1993).

**Distribution:** Brazil (Amazonas, Pará) and French Guiana.

***Culicoides brasilianum* Forattini, 1956**

*Culicoides brasilianum* Forattini (1956b): 81. São Paulo, Brazil.

**Identification:** Eyes contiguous by a distance equal to the diameter of 1 ommatidial facet; third palpal

segment slightly broader just beyond middle, pit irregular; r-m crossvein pale, one transverse distal pale spot in cell  $r_3$  broadly meeting wing margin, two distal pale spots in cell  $m_1$ , apices of veins  $M_1$ ,  $M_2$  and  $CuA_1$  pale, apex of vein  $CuA_2$  dark; halter knob dark, pedicel pale; spermathecae pyriform, unequal. Male genitalia with apicolateral processes of ninth tergum small and close together; aedeagus with terminal papilla; parameres connected at bases by a short loop, main bodies stout, apex with minute fringing hairs. This species distinguished from the related species by the wing pattern with extensive pale areas; from *C. guttatus* it also differs in the pale r-m crossvein and dark halter, and from *C. ignacioi* by the pale r-m crossvein (Spinelli et al. 1993).

**Distribution:** Northeastern Argentina and Brazil (Amazonas, Pará, Pernambuco, Rio de Janeiro, São Paulo, and Santa Catarina).

**\**Culicoides brownie*** Spinelli, 1993

(Figures 3, 10N)

*Culicoides brownie* Spinelli et al. (1993): 24. Colombia.

**Identification:** Wing with second distal pale spot in cell  $m_1$  sometimes faint or absent. This species is very similar to *C. franklini*, from which it can be distinguished by the stouter third palpal segment with irregular pit, eyes contiguous for 2.5–3.0 ommatidial facets (by 4 or 5 facets in *C. franklini*), and by its dark halter (Spinelli et al. 1993).

**Distribution:** Colombia and Brazil (Amazonas).

**New records:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VIII-2010, 0001-3 (1 female), CDC light trap, peridomicily, F.A.C. Pessoa collector.

***Culicoides coutinhoi*** Barretto, 1944

(Figures 3, 10O)

*Culicoides coutinhoi* Barretto (1944): 96. São Paulo, Brazil.

**Identification:** Eyes contiguous by a distance equal to the diameter of 3 ommatidial facets; third palpal segment stout, with irregular sensory pit; wing as in figure. This species is similar to *C. pseudodiabolicus*, from which it can be distinguished by the stout palpus with irregular pit (subdivided in *C. pseudodiabolicus*), by the small apicolateral processes closer together, and by the terminal papilla of the aedeagus (Spinelli et al. 1993).

**Distribution:** Brazil (Amazonas, Pará, and São Paulo) and French Guiana

***Culicoides diabolicus*** Hoffman, 1925

(Figures 3, 10P)

*Culicoides diabolicus* Hoffman (1925): 294. Panama.

**Identification:** Eyes contiguous by a distance equal to diameter of 1.5–2.5 ommatidial facets; antenna brown, bases of flagellar segments 3–10 pale; third palpal segment slender, slightly broad in middle, pit irregular; wing as shown in figure, with r-m crossvein pale (faintly darkened in some topotypic specimens), one large,

transverse, distal pale spot in cell  $r_3$  broadly meeting wing margin, two distal pale spot in cell  $m_1$ , the distal one smaller, apices of veins  $M_1$  and  $M_2$  pale, apices of veins  $CuA_1$  and  $CuA_2$  dark; halter pale; spermathecae subspherical to ovoid, unequal (Spinelli et al. 1993).

**Distribution:** Mexico to Ecuador and Brazil (Amazonas, Roraima, Maranhão).

***Culicoides filariferus*** Hoffman, 1939

(Figures 3, 10Q)

*Culicoides filariferus* Hoffman (1939): 172. Mexico.

**Identification:** Eyes contiguous by a distance equal to the diameter of 1.5–2.0 ommatidial facets; third palpal segment slightly broader in middle, with a conspicuous distal extension, pit irregular; wing as shown in figure, r-m crossvein pale (a faint darkish line, but not a definite dark spot, sometimes present on anterior half); one distal transverse pale spot in cell  $r_3$  broadly meeting wing margin, two distal pale spot in cell  $m_1$  (the second one faint or absent in some specimens); apices of veins  $M_1$  and  $M_2$  pale, apices of veins  $CuA_1$  and  $CuA_2$  dark (a few specimens with a faint pale area at apex of vein  $CuA_2$ ); halter dark brown (Spinelli et al. 1993).

**Distribution:** Mexico (Veracruz and Chiapas) to northern Brazil (Amazonas and Pará).

**\**Culicoides flavivenulus*** Costa Lima, 1937

(Figures 3, 10R)

*Culicoides flavivenula* [sic] Costa Lima (1937): 418. Brazil.

**Identification:** Eyes contiguous by a distance equal to the diameter of 1–2 ommatidial facets; wing as shown in figure, r-m crossvein pale, one transverse distal pale spot in cell  $r_3$  reaching wing margin, only one distal pale spot in cell  $m_1$ , vein  $M_1$  slightly pale at apex; halter pale; spermathecae subspherical to pyriform, subequal. This species is similar to *C. lutzi*, from which it can be distinguished by the third palpal segment with a well-defined, round pit (irregular in *C. lutzi*); apices of veins  $M_2$  and  $CuA_1$  dark, broadly pale in *C. lutzi* (Spinelli et al. 1993).

**Distribution:** Panama, French Guiana, Trinidad, and Brazil (Amazonas, Pará, Bahia, Pernambuco, Mato Grosso do Sul, Espírito Santo, Rio de Janeiro, São Paulo, and Santa Catarina).

**New records:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VI-2010, VIII-2010, 0001-4 (3 females), CDC light traps, forest fragments and peridomicilies, F.A.C. Pessoa collector.

***Culicoides foxi*** Ortiz, 1950 (Figures 4, 10S)

*Culicoides foxi* Ortiz (1950a): 461. Puerto Rico.

**Identification:** Eyes contiguous for a distance equal to the diameter of 1 ommatidial facet; third palpal segment elongated, pit rounded and shallow, the extension beyond the pit slender; wing as shown in figure, r-m crossvein entirely an deeply infuscated, one large, transverse, distal pale spot in cell  $r_3$  broadly meeting wing

margin, two distal pale spots in cell  $m_1$ , apices of veins  $M_1$ ,  $M_2$  and  $CuA_1$  broadly pale, apex of vein  $CuA_2$  dark; halter knob deeply infuscated, pedicel pale; spermathecae subspherical to ovoid, unequal (Spinelli et al. 1993).

**Distribution:** Mexico to northeastern Argentina; Brazil (Amazonas, Pará, Roraima, Bahia, Maranhão, Pernambuco, Piauí, Goiás, Mato Grosso do Sul, Minas Gerais, Rio de Janeiro, São Paulo, and Paraná).

\****Culicoides franklini*** Spinelli, Greiner & Wirth, 1993 (Figures 4, 10T)

*Culicoides franklini* Spinelli et al. (1993): 45. Panama.

**Identification:** Eyes contiguous for a distance equal to the diameter of 2–2.5 ommatidial facet; third palpal segment stout, with shallow, irregular pit; wing as shown in figure. This species is easily distinguished from *C. pseudodiabolicus*, *C. coutinhoi*, and *C. brownei* by the broadly contiguous eyes, very short proboscis, and third palpal segment shorter than combined lengths of segments 4 and 5 (Spinelli et al. 1993).

**Distribution:** Mexico (Guerrero) to Bolivia and Brazil (Amazonas and Pará).

**New records:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VI-2010, VIII-2010, 0001-5 (5 females), CDC light traps, forest fragments and peridomicilios, F.A.C. Pessoa collector.

***Culicoides fusipalpis*** Wirth & Blanton, 1973 (Figures 4, 10U)

*Culicoides fusipalpis* Wirth and Blanton (1973): 435. Pará, Brazil

**Identification:** Eyes contiguous by a distance equal to the diameter of 2–2.5 ommatidial facets; third palpus segment fusiform, slightly swollen in midportion only, with scattered surface sensilla; wing as figured, r-m crossvein in a large pale spot, but vein slightly darkened, one large transverse distal pale spot in cell  $r_3$ , broadly meeting wing margin, one distal pale spot in cell  $m_1$ , apices of veins  $M_1$  and  $M_2$  narrowly pale, apices of veins  $CuA_1$  and  $CuA_2$  dark; halter pale; spermathecae subequal, subspherical with short sclerotized necks (Spinelli et al. 1993).

**Distribution:** El Salvador to Bolivia, Ecuador, French Guyana, Guyana, and Brazil (Acre, Amapá, Amazonas, Pará, Rondônia, Roraima, Bahia, Maranhão, and Rio de Janeiro).

***Culicoides heliconiae*** Fox & Hoffman, 1944 (Figure 5)

*Culicoides heliconiae* Fox and Hoffman (1944): 108. Venezuela.

**Identification:** Third palpal segment with scattered sensilla; apical pale band on mid femur, hind femur dark to tip; wing with distal pale spot in  $r_3$  narrow and transverse,  $r_3$  with pale spot present anterior to base of  $M_1$ , a single pale spot crosses the second radial cell, apices of  $CuA_1$  and  $CuA_2$  pale; spermathecae with short, slender necks (Felippe-Bauer et al. 2009).

**Distribution:** Belize to Venezuela, Ecuador, and Bra-

zil (Acre, Amazonas, and Pará); Grenada, and Trinidad and Tobago.

***Culicoides hylas*** Macfie, 1940 (Figures 5, 11A)

*Culicoides hylas* Macfie (1940): 26. Guyana.

**Identification:** Irregular sensory pit in the third palpal segment; wing as shown,  $r_3$  with pale spot present anterior to base of vein  $M_1$ , a single pale spot crossing the second radial cell, apices of  $CuA_1$  and  $CuA_2$  pale; mid femur with subapical pale band, hind femur dark to tip; spermathecae with short, slender necks (Felippe-Bauer et al. 2009).

**Distribution:** Mexico to Peru and Brazil (Amapá, Amazonas, Pará, Roraima, and Maranhão).

***Culicoides ignacioi*** Forattini, 1957 (Figures 4, 11B)

*Culicoides ignacioi* Forattini (1957): 215. São Paulo, Brazil.

*Culicoides saintjusti* Tavares and Ruiz (1980): 27. Rio de Janeiro, Brazil.

**Identification:** Eyes contiguous by a distance equal to the diameter of 1.5 ommatidial facets; third palpus segment broad in middle, with conspicuous irregular pit; wing as shown in figure; halter knob bark brown, pedicel pale. This species is similar to *C. fernandoi*, from which it can be distinguished by the mandible with 20–22 teeth (14–15 in *C. fernandoi*), vein  $R_3$  pale (infuscated on lower portion with a very small dark spot behind apex in *C. fernandoi*), distal pale spot in cell  $r_3$  large, transverse (crescent-shaped or subdivided in *C. fernandoi*) (Spinelli et al. 1993).

**Distribution:** Brazil (Acre, Amazonas, Pará, Roraima, Maranhão, Minas Gerais, Rio de Janeiro, and São Paulo) and Paraguay.

***Culicoides insignis*** Lutz, 1913 (Figures 4, 11C)

*Culicoides insignis* Lutz (1913): 51. Bahia and Rio de Janeiro, Brazil.

*Culicoides inamollae* Fox and Hoffman (1944): 110. Puerto Rico.

*Culicoides painteri* Fox (1946): 257. Puerto Rico.


**Identification:** Eyes contiguous by a distance equal to the diameter of 1.5 ommatidial facets; third palpal segment with definite, irregular pit; wing as shown in figure, r-m crossvein dark on anterior half; vein  $CuA_2$  dark up to the point where it turns abruptly forward to meet the costa; only one distal pale spot in cell  $m_1$ ; halter dark (Spinelli et al. 1993).

**Distribution:** USA, Mexico, Central America, and the Caribbean; Brazil (Amazonas, Pará, Bahia, Ceará, Paraíba, Pernambuco, Maranhão, Goiás, Mato Grosso, Mato Grosso do Sul, Espírito Santo, Minas Gerais, São Paulo, Rio de Janeiro, Rio Grande do Sul, and Santa Catarina).

***Culicoides lutzi*** Costa Lima, 1937 (Figures 4, 11D)

*Culicoides lutzi* Costa Lima (1937): 419. Pará, Brazil.

**Identification:** Brown species; eyes contiguous by a



**Figure 4.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. foxi*. **B.** *C. franklini*. **C.** *C. fusipalpis*. **D.** *C. ignacioi*. **E.** *C. insignis*. **F.** *C. lutzi*. **G.** *C. ocumarensis*. **H.** *C. paraignacioi*. **I.** *C. paramaruim*. **J.** *C. plaumanni* (modified after SimpleMappr, <http://www.simplemappr.net>).

distance equal to the diameter of 2–3 ommatidial facets; third palpal segment stout, with irregular pit; wing as shown in figure, crossvein r-m pale, one transverse distal pale spot in cell  $r_3$  reaching wing margin, only one distal pale spot in cell  $m_1$ , apices of veins  $M_1$ ,  $M_2$  and  $CuA_1$  broadly pale, apex of vein  $CuA_2$  dark; halter knob pale (Spinelli et al. 1993).

**Distribution:** Colombia and French Guiana; Brazil (Acre, Amazonas, Pará, Rondônia, Roraima, Maranhão, Goiás, Mato Grosso, Minas Gerais, Rio de Janeiro, São Paulo, Paraná, and Santa Catarina) and northeastern Argentina.

***Culicoides ocumarensis*** Ortiz, 1950

(Figures 4, 11E)

*Culicoides ocumarensis* Ortiz (1950b): 455. Venezuela.

**Identification:** Eyes contiguous by a distance equal to the diameter of 1.5–2.0 ommatidial facets; third palpal segment slightly broader in middle, pit irregular; wing

as shown in figure, r-m crossvein pale, one transverse distal pale spot in cell  $r_3$  broadly meeting wing margin, two distal pale spot in cell  $m_1$ , apices of veins  $M_1$  and  $M_2$  pale, apices of veins  $CuA_1$  and  $CuA_2$  dark; halter dark brown (Spinelli et al. 1993).

**Distribution:** Mexico to northern Brazil (Amazonas, Pará, and Rondônia).

***Culicoides palpalis*** Macfie, 1948

*Culicoides palpalis* Macfie (1948): 78. Mexico.

**Identification:** Third palpal segment with scattered sensilla; mid femur with apical pale band, hind femur dark to tip; wing with  $r_3$  shows a pale spot present anterior to base of  $M_1$ , a single pale spot crosses the second radial cell, apices of  $CuA_1$  and  $CuA_2$  dark; spermathecae with short, slender necks; male tergite 9 with very small papilliform process on posterior margin; fused portion of the parameres shows a width to length ratio of 1.75 (Felippe-Bauer et al. 2009).

**Distribution:** Mexico to Peru and Brazil (Amazonas).

***Culicoides paraignacioi* Spinelli, 1993**

(Figures 4, 11F)

*Culicoides paraignacioi* Spinelli in Spinelli et al. (1993): 66. Colombia.

**Identification:** Small wing length 1.14 mm, third palpal segment slender, with a rounded, small, shallow pit; wing as figured. This species is similar to *C. ignacioi*, which differs from the females of *C. paraignacioi* by being significantly larger (wing length 1.53 mm), and by having the third palpal segment stouter, with a conspicuous irregular sensory pit. The male genitalia of both species are also very similar, with apicolateral processes widely separated, apex of aedeagus truncated, and parameres broadly fused at base, but they differ in the shape of the ninth tergum, which is rounded distally in *C. ignacioi* and subquadrangular in *C. paraignacioi* (Spinelli et al. 1993).

**Distribution:** Belize, Costa Rica, Colombia, Guiana, and Brazil (Amazonas and Pará).

**\**Culicoides paramaruim* Wirth & Blanton, 1973**

(Figures 4, 11G)

*Culicoides paramaruim* Wirth and Blanton (1973): 443. –Pará, Brazil.

**Identification:** Medium-sized dark brown; wing as figured, r-m crossvein darkened, especially so on anterior half, vein CuA<sub>1</sub> infuscated, one transverse distal pale spot in cell r<sub>3</sub> not reaching wing margin; only one distal pale spot in cell m<sub>1</sub>; apices of veins M<sub>1</sub>, M<sub>2</sub>, CuA<sub>1</sub> and CuA<sub>2</sub> dark. This species differs from *C. maruim*, from which it can be distinguished by the third palpal segment stouter, while in *C. maruim* is long and slender (Spinelli et al. 1993).

**Distribution:** Brazil (Amazonas, Pará, and Maranhão).

**New record** — Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VII-2010, 0001-7 (1 female), CDC light trap, peridomicity, F.A.C. Pessoa collector.

***Culicoides plaumanni* Spinelli, 1993**

(Figures 4, 11H)

*Culicoides plaumanni* Spinelli in Spinelli et al. (1993): 69. Argentina.

**Identification:** Medium-sized, eyes contiguous by a distance equal to the diameter of 1.5–2.5 ommatidial facets; third palpal segment with irregular pit; wing as shown in figure, r-m crossvein dark on anterior half, one distal, somewhat crescent-shaped pale spot in cell r<sub>3</sub> narrowly meeting wing margin, two distal pale spots in cell m<sub>1</sub>, apices of veins M<sub>1</sub>, M<sub>2</sub> and CuA<sub>1</sub> pale, apex of vein CuA<sub>2</sub> dark; halter dark brown (Spinelli et al. 1993).

**Distribution:** Bolivia, Peru, Brazil (Amazonas, Pará, and Minas Gerais) and northeastern Argentina.

***Culicoides polypori* Wirth & Blanton, 1968**

*Culicoides polypori* Wirth and Blanton (1968): 212. Panama.

**Identification:** Third palpal segment with scattered sensilla; wing with r<sub>3</sub> with pale spot present anterior to base of M<sub>1</sub>, a single pale spot crosses the second radial cell, apices of CuA<sub>1</sub> and CuA<sub>2</sub> dark; mid and hind femur with subapical pale band; spermathecae with short, slender necks; male tergite 9 with a bilobed process on posterior margin; fused portion of the parameres slightly longer than basal width (Felippe-Bauer et al. 2009).

**Distribution:** Honduras to Colombia and Brazil (Amazonas).

***Culicoides pseudodiabolicus* Fox, 1946**

(Figures 5, 11I)

*Culicoides pseudodiabolicus* Fox (1946): 256. Trinidad.

**Identification:** Wing with contrasting pattern, r-m crossvein dark on anterior half, vein R<sub>3</sub> pale, one large transverse distal pale spot in cell r<sub>3</sub> broadly meeting wing margin, cell m<sub>1</sub> with two distal pale spots, apices of veins M<sub>1</sub>, M<sub>2</sub> and CuA<sub>1</sub> pale, apex of vein CuA<sub>2</sub> dark, scattered macrotrichia present at apices of cells r<sub>3</sub>, m<sub>1</sub>, and m<sub>2</sub>. Several species in the *guttatus* group are very similar to *C. pseudodiabolicus*, especially in wing; characteristic subdivided pit, pale halter, distinctly unequal spermathecae, and male aedeagus without terminal papilla (Spinelli et al. 1993).

**Distribution:** Mexico to Peru and Brazil (Acre, Amapá, Amazonas, Pará, Rondônia, Roraima, and Maranhão).

***Culicoides pseudoheliconiae* Felippe-Bauer, 2008**

(Figure 5)

*Culicoides pseudoheliconiae* Felippe-Bauer in Felippe-Bauer et al. (2008): 260. Peru.

**Identification:** Pale diffuse wing markings, rather than brightly contrasting; r<sub>3</sub> with pale spot present anterior to base of M<sub>1</sub>, pale spot that crosses the second radial cell subdivided in two separate spots, apices of CuA<sub>1</sub> and CuA<sub>2</sub> dark; third palpal segment with scattered sensilla; mid femur with apical pale band, hind femur dark to tip; spermathecae with short, slender necks (Felippe-Bauer et al. 2009).

**Distribution:** Peru (Madre de Dios and San Martín), Brazil (Amazonas), and Argentina (Misiones).


***Culicoides ruizi* Forattini, 1954**

(Figure 5)

*Culicoides ruizi* Forattini (1954a): 189. Goiás, Brazil

**Identification:** This species is very similar to *C. travassosi*, from which it can be distinguished by the stouter palpus, pale r-m crossvein, and vein CuA<sub>2</sub> without a small blackish spot just beyond apex (Spinelli et al. 1993).

**Distribution:** Colombia and Brazil (Amazonas, Pará, Maranhão, and Goiás).



**Figure 5.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. pseudodiabolicus*. **B.** *C. tidwelli*. **C.** *C. aitkeni*. **D.** *C. baniwa*. **E.** *C. heliconiae*. **F.** *C. hylas*. **G.** *C. pseudoheliconiae*; **H.** *C. verecundus*. **I.** *C. aldomari*. **J.** *C. bricenoi*. **K.** *C. ruizi* (modified after SimpleMappr, <http://www.simplemappr.net>).

**\**Culicoides tidwelli* Spinelli, 1993**

(Figures 5, 11J)

*Culicoides tidwelli* Spinelli in Spinelli et al. (1993): 74. Colombia.

**Identification:** Small wing as shown in figure; proximal antennal segments elongated, about twice as long as broad; halter knob pale (Spinelli et al. 1993). This species differs from *C. diabolicus* by the short and rounded palpal segment (definite extension behind tip; palpal segments 4 and 5 slender in *C. diabolicus*), sensilla coeloconica on flagellomeres additional sometimes present on 1, 5, 7, while in *C. diabolicus* the sensilla coeloconica on flagellomeres 1, 9–13 (Spinelli et al. 1993).

**Distribution:** Honduras to Colombia, Ecuador, and Brazil (Amazonas).

**New records:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VI-2010, VIII-2010, 0001-9 (8 females), CDC light traps, peridomicilies, F.A.C. Pessoa collector.

***Culicoides travassosi* Forattini, 1957**

*Culicoides travassosi* Forattini (1957): 198. Pará, Brazil

**Identification:** Third palpal segment slender, slightly broader beyond middle, pit subdivided; r-m crossvein dark, two distal pale spots in  $r_3$  (the usual one large, subdivided in some specimens, and a small second near wing tip), two large distal pale spots in cell m1, apices of veins  $M_1$ ,  $M_2$  and  $CuA_1$  very narrowly pale, apex of vein  $CuA_2$  dark; halter with dark brown knob, pedicel pale (Spinelli et al. 1993).

**Distribution:** Suriname and Brazil (Amazonas, Pará, and Mato Grosso).

***Culicoides verecundus* Macfie, 1948**

(Figures 5, 11K)

*Culicoides verecundus* Macfie (1948): 76. Mexico.

**Identification:** Third palpal segment contains scattered sensilla; mid femur with apical and hind femur

with subapical pale band; wing as shown in figure,  $r_3$  with pale spot present anterior to base of  $M_1$ , a single pale spot crossing second radial cell, apices of  $CuA_1$  and  $CuA_2$  dark; spermathecae with short, slender necks. Male tergite 9 with a prominent papilliform process on posterior margin; fused portion of the parameres longer than width; separate portion long and V-shaped at base (Felippe-Bauer et al. 2009).

**Distribution:** Mexico to Panama; Brazil (Amazonas, Pará, and Roraima).

#### Subgenus *Mataemyia* Vargas, 1960

*Mataemyia* Vargas (1960): 43, as subgenus of *Culicoides*. Type species: *Culicoides mojingaensis* Wirth & Blanton, 1953, by original designation.

#### \**Culicoides aldomari* Felippe-Bauer, Silva & Trindade, 2013 (Figures 5, 11L)

*Culicoides aldomari* Felippe-Bauer et al. (2013): 56. Pará, Brazil.

**Identification:** Medium-sized; eyes narrowly separated; flagellomeres 8 and 9 subequal, sensilla coeloconica on flagellomeres 1, 6–8, AR 0.77–0.89; PR 2.0–2.2; wing as figured. This species can be distinguished from *C. azureus* Wirth & Blanton, 1959, *C. barthi* Tavares & Souza, 1978, *C. felippebauerae* Spinelli, 2007, and *C. mojingaensis* Wirth & Blanton, 1953 all subgenus *Mataemyia*, due to the smaller wing, darker lumen of the second radial cell (pale lumen in the others species), by the small dark spot in  $r_3$ , located on the distal end of second radial cell (without defined dark spot in *C. barthi*, dark spot behind second radial cell in *C. azureus*, *C. felippebauerae* and *C. mojingaensis*) and by the flagellomeres without transition in length between proximal and distal series, in the others species the flagellomeres 9–12 are more elongated than 2–8 (Felippe-Bauer et al. 2013).

**Distribution:** Brazil (Amazonas and Pará).

**New record:** Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VII-2010, 0001-1 (1 female), CDC light traps, forest fragments, F.A.C. Pessoa collector.

#### *Culicoides bricenoi* Ortiz, 1915

(Figures 5, 11M)

*Culicoides bricenoi* Ortiz (1915b): 445. Venezuela.

**Identification:** Second radial cell long, twice longer than first;  $r_3$  with two large pale spots, the distal one located in distal portion of cell reaching anterodistal wing margin (Felippe-Bauer et al. 2013).

**Distribution:** Venezuela, Ecuador, Peru, Bolivia, and Brazil (Amazonas, Pará, and Roraima).

#### *Culicoides discrepans* Ortiz & Mirsa, 1951

(Figure 6)

*Culicoides discrepans* Ortiz and Mirsa (1951): 595. Venezuela.

**Identification:** A large species; wing with distal pale spot in  $r_3$  single, basal pale spot in  $m_1$  large, slightly connected with the basal pale spot in  $m_2$ , distal pale spot in

$m_2$  reaching wing margin; one distal pale spot in anal cell, pale spot on  $M_1$  restricted to extreme apex, connected with the distal pale spot in  $r_3$  and  $m_1$  (Felippe-Bauer et al. 2013).

**Distribution:** Venezuela and Brazil (Amazonas).

#### *Culicoides felippebauerae* Spinelli, 2007

(Figure 6)

*Culicoides felippebauerae* Spinelli in Spinelli et al. (2007): 660, Amazonas, Brazil.

**Identification:** The only species in the subgenus *Mataemyia* with one spermatheca; male with stout, sinuate parameres with filiform tip; posteromedial projection of aedeagus tapered to slender, narrow, blunt tip, with a pair of well developed subapical points (Spinelli et al. 2007).

**Distribution:** Brazil (Amazonas).

#### *Culicoides wallacei* Wirth and & Blanton, 1973

(Figures 6, 11N)

*Culicoides wallacei* Wirth and Blanton (1973): 449. Pará, Brazil.

**Identification:** A large species; wing as shown in figure, first and 2<sup>nd</sup> second radial cells nearly similar in length, distal pale spot in  $r_3$  oblique, broadly reaching wing margin; anal cell with two distal small, round, pale spot (Spinelli et al. 2007).

**Distribution:** Brazil (Amazonas, Pará, and Maranhão).

#### Subgenus *Oecacta* Poey, 1853

*Oecacta* Poey (1853): 238. Type species: *Oecacta furens* Poey, 1853, by monotypy.

*Diplosella* Kieffer (1921): 113, as subgenus of *Culicoides*. Type species: *Culicoides sergenti* Kieffer, 1921, by monotypy.

*Sensiculoides* Shevchenko (1977): 133, as subgenus of *Culicoides*. Type species: *Ceratopogon pictipennis* Staeger, 1839 by original designation.

#### *Culicoides alahialinus* Barbosa, 1952

(Figures 6, 11O)

*Culicoides alahialinus* Barbosa (1952): 11. Ecuador.

**Identification:** Eyes narrowly separated, bare; third palpal segment moderately swollen, with moderately large, shallow, sensory pit; wing as figured, pale area at wing base and over r-m crossvein; very faint indications of pale spots in cells on distal part of wing; halter infuscated; spermathecae two, slightly, pyriform, slightly unequal (Wirth and Blanton 1959).

**Distribution:** Costa Rica, Panama, Colombia, Ecuador, and Brazil (Amazonas).

Subgenus unplaced, *acotylus* species group


#### *Culicoides acotylus* Lutz, 1913

(Figures 6, 11P)

*Culicoides acotylus* Lutz (1913): 69. Mato Grosso, Brazil.

*Culicoides panamericanus* Fox (1947): 90. Mexico.

**Identification:** Third palpal segment moderately swollen, without sensory pit, the sensoria scattered on



**Figure 6.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. discrepans*. **B.** *C. felippebaueri*. **C.** *C. wallacei*. **D.** *C. alahialinus*. **E.** *C. acotylus*. **F.** *C. belemensis*. **G.** *C. camposi*. **H.** *C. carpenteri*. **I.** *C. dasyophrus*. **J.** *C. eublepharus* (modified after SimpleMappr, <http://www.simplemappr.net>).

surface of segment; wing as figured, second radial cell very dark, r-m crossvein blackish, in the center of a large pale spot which broadly reaches costal margin, cell  $r_3$  with five definite pale spots, apices of veins  $M_1$  and  $M_2$  pale margined a short distance, a pale spot each in apex of cell  $m_2$  and cell  $CuA_1$ , neither attaining wing margin, two pale spots in distal portion of anal cell and two in basal portion of this cell; cell  $m_2$  with a pale spot lying in front of mediocubital fork, a lying behind medial fork and another lying across cell about halfway from these spots and wing base; halter pale; spermathecae two, pyriform, slightly unequal, the bases of the ducts sclerotized a short distance (Wirth and Blanton 1959).

**Distribution:** Mexico, Honduras, Panama, Venezuela, Trinidad, Suriname, and Brazil (Amazonas, Pará, Roraima, and Mato Grosso).

Subgenus unplaced, *carpenteri* species group

### ***Culicoides belemensis* Wirth & Blanton, 1973**

(Figures 6, 11Q)

*Culicoides belemensis* Wirth and Blanton (1973): 427. Pará, Brazil.

**Identification:** Eyes narrowly separated, bare; third segment only slightly swollen with small, round, shallow, sensory pit on distal; wing as figured; halter brownish; spermathecae two, ovoid, unequal; aedeagus with disti-median process moderately slender with parallel sides and blunt, concaved tip; parameres with distinct ventral swelling ending in a distal lobe, apical portion slender, bearing lateral spines and terminating in a filiform tip (Wirth and Blanton 1973).

**Distribution:** Brazil (Amazonas, Pará) and Colombia.

### ***Culicoides camposi* Ortiz & León, 1955**

(Figure 6)

*Culicoides camposi* Ortiz and León (1955): 569. Ecuador.

*Culicoides fairchildi* Wirth and Blanton (1955): 102. Panama.

**Identification:** Third palpal segment swollen, with a broad, shallow, irregular sensory pit; second radial cell very dark; pale spot over r-m crossvein broadly reaching costal margin, distal pale spot in cell  $r_3$  rounded, almost reaching anterior wing margin, distal pale spot in cell  $m_1$  elongate and nearly reaching wing margin, two pale spots, which may be more or less fused, in distal part of anal cell; base of anal cell pale; cell  $m_2$  with a small pale spot lying in front of mediocubital fork and a pale area extending from behind medial fork to base of wing; halter yellowish (Wirth and Blanton 1959).

**Distribution:** Costa Rica, Panama, Colombia, Ecuador, and Brazil (Amazonas).

***Culicoides carpenteri* Wirth & Blanton, 1953**  
(Figure 6)

*Culicoides carpenteri* Wirth and Blanton (1953): 72. Panama

**Identification:** Third palpal segment scarcely swollen, with a broad, shallow, sensory pit; second radial cell very dark, large pale area over r-m crossvein, distal pale spot in cell  $r_3$  extensive, broadly extending across cell near apex from anterior wing margin to vein  $M_1$ ; apices of veins  $M_1$ ,  $M_2$  and  $CuA_1$  dark; subapical pale spot in cell  $m_2$ , distal pale spot in cell  $m_1$  broadly meeting wing margin; two pale spots in distal part of cell  $m_2$ , the distal one broadly meeting wing margin; the proximal one connected by a pale area extending to base of cell and including the pale spots lying in front of mediocubital fork and behind medial fork; pale area in cell  $CuA_1$  nearly filling entire cell; anal cell pale except for a large dark area centering on middle of stem of mediocubital vein; halter pale (Wirth and Blanton 1959).

**Distribution:** Panama and Brazil (Amazonas).

Subgenus unplaced, *dasyophrus* species group

***Culicoides dasyophrus* Macfie, 1940**  
(Figures 6, 11R)

*Culicoides dasyophrus* Macfie (1940): 27. Guyana.

**Identification:** Eyes narrowly separated above, with long interfacetal hairs; third palpal segments swollen, with a broad, shallow, sensory pit; wing as shown in figure, poststigmatic pale spots in cell  $r_3$  more or less fused, the posterior one located slightly proximad of the anterior one, distal pale spot in cell  $r_3$  small, only one small pale spot in distal part of anal cell and one pale spot in distal part of cell  $m_2$ ; halter whitish; spermatheca one, pyriform (Wirth and Blanton 1956).

**Distribution:** Colombia, Venezuela, Ecuador, Guyana, and Brazil (Amazonas, Pará, Roraima, and Mato Grosso).

Subgenus unplaced, *eublepharus* species group

***Culicoides eublepharus* Macfie, 1948**  
(Figures 6, 48S)

*Culicoides eublepharus* Macfie (1948): 86. Guyana.

*Culicoides transferrans* Ortíz (1953): 801. Venezuela.

**Identification:** Eyes narrowly separated, with long interfacetal hairs; third palpal segment moderately swollen, with a broad, shallow, sensory pit; wing as shown in figure, with four pale spots in a rhomboid in cell  $r_3$ , the distal pair usually fused and broadly attaining wing margin anteriorly; two pale spots each in cells  $m_1$  and apices of cells  $m_2$  and anal cell; pale spots present behind medial fork and in front of mediocubital fork; halter yellowish; spermatheca one, oval with a short portion of the duct sclerotized (Wirth and Blanton 1959).

**Distribution:** Mexico, Costa Rica to Venezuela and Ecuador, and Brazil (Amazonas, Pará, and Roraima).

***Culicoides propriipennis* Macfie, 1948**  
(Figures 7, 11T)

*Culicoides propriipennis* Macfie (1948): 84. Mexico.

**Identification:** This species is most similar to *C. tetrathyris*, but *C. propriipennis* can easily be distinguished by the presence of a third pale spot at the wing margin in cell  $m_1$ , by the second pale spot in the distal portion of cell  $m_2$ , wing as figured (Wirth and Blanton 1959).

**Distribution:** Mexico to Panama, Venezuela, Ecuador, and Brazil (Amazonas, Pará, and Bahia).

***Culicoides rangeli* Ortíz & Mirsa, 1952**  
(Figure 7)

*Culicoides rangeli* Ortíz and Mirsa (1952): 126. Venezuela.

*Culicoides donajii* Vargas (1954): 28. Mexico.

*Culicoides patulipalpis* Wirth and Blanton (1959): 421. Panama.

**Identification:** Eyes nearly contiguous, with long interfacetal hairs; third segment swollen, with a very broad, shallow, sensory pit, open and without regular pore; posterior poststigmatic pale spot in cell  $r_3$  very small or entirely absent, distal spot in cell  $r_3$  transverse and not double, not attaining wing margin; two very small pale spots each in cell  $m_1$  and in apices of cell  $m_2$  and anal cell; a pale spot present behind medial fork and another in front of mediocubital fork; halter pale; spermatheca one, pyriform, with long sclerotized neck (Wirth and Blanton 1959).


**Distribution:** Mexico to Venezuela, Trinidad, Ecuador, Bolivia, and Brazil (Amazonas).

Subgenus unplaced, *fluvialis* species group

***Culicoides fluvialis* Macfie, 1940** (Figures 7, 11U)

*Culicoides fluvialis* Macfie (1940): 25. Guyana.

**Identification:** Third palpal segment short and slightly swollen, with a small, deep sensory pit; wing as shown in figure, with second radial cell moderately dark, cell  $r_3$  with two separate round poststigmatic pale spots, the posterior one located behind second radial cell, the distal pale spot hourglass-shaped on wing margin, apices of veins  $M_1$  and  $M_2$  with a small pale spot at wing margin, cell  $CuA_1$  with a round pale spot almost meeting



**Figure 7.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. propiipennis*. **B.** *C. rangeli*. **C.** *C. fluvialis*. **D.** *C. leopoldoi*. **E.** *C. tetrathyris*. **F.** *C. benarrochi*. **G.** *C. fieldi*. **H.** *C. glabellus*. **I.** *C. leoni*. **J.** *C. galindoi* (modified after SimpleMappr, <http://www.simplemappr.net>).

wing margin, only one small round pale spot in distal part of anal cell, cell  $m_2$  with a small pale spot present lying behind medial fork; halter pale; spermathecae two, oval, slightly unequal (Wirth and Blanton 1959).

**Distribution:** Honduras to Colombia, Venezuela, Trinidad, Guyana, and Brazil (Amazonas, Pará, and Roraima).

***Culicoides leopoldoi*** Ortíz, 1951  
(Figures 7, 12A)

*Culicoides leopoldoi* Ortíz (1951a): 579. Venezuela.

**Identification:** Wing as figured, with pale spot on crossvein r-m extensive; pale spot present behind second radial cell, proximal to the poststigmatic pale spots; poststigmatic pale spots longitudinally aligned, posterior one smaller; distal pale spot in cell  $r_3$  large, with narrow proximal extension; one spermatheca (Spinelli et al. 2005).

**Distribution:** Guatemala and Belize to Bolivia, north-eastern Argentina, Trinidad, and Brazil (Amapá, Ama-

zonas, Pará, Rondônia, Roraima, Bahia, Ceará, Maranhão, Pernambuco, Espírito Santo, Minas Gerais, and Rio de Janeiro).

***Culicoides tetrathyris*** Wirth & Blanton, 1959

(Figures 7, 12B)

*Culicoides tetrathyris* Wirth and Blanton (1959): 409. Panama.

**Identification:** Resembling *C. propiipennis* in wing pattern, mesonotal and leg markings, structural characters, and male genitalia, but *C. propiipennis* differs by having the sides of the mesonotum brownish instead of bluish pruinose, a third pale spot in cell  $m_1$  at the wing margin, the pale spots in cell  $r_3$  arranged in a rhomboid instead of a trapezoid, distal sensory tufts in flagellomeres 1, 7–12, and the pit on the third palpal segment broader. The male genitalia of the two species are practically identical, but in *C. propiipennis* the stem of the paramere is stouter and not bent near the base (Wirth and Blanton 1959).

**Distribution:** Honduras, Costa Rica, Panama, Ecua-

dor, Suriname, Trinidad, and Brazil (Acre, Amazonas, Pará, Roraima, and Bahia).

Subgenus unplaced, *leoni* species group

***Culicoides benarrochi*** Ortiz & Mirsa, 1952

(Figures 7, 12C)

*Culicoides benarrochi* [sic] Ortiz and Mirsa (1952): 126. Venezuela.

**Identification:** Wing as figured; separated from *C. fieldi* by its shorter second palpal segment (9/13 as long as third segment), flagellomeres 9–11 and 12 subequal in length and by the presence of a distinct pale spot in front of the mediocubital fork (Wirth and Blanton 1959).

**Distribution:** Venezuela, Trinidad, and Brazil (Amazonas, Roraima, and Rio de Janeiro).

***Culicoides fieldi*** Wirth & Blanton, 1956

(Figure 7)

*Culicoides fieldi* Wirth and Blanton (1956): 50. Honduras.

**Identification:** Eyes contiguous, with long interfacetal hairs; third segment moderately swollen, with a small, deep, sensory pit; pale spot over r-m crossvein small, poststigmatic pale spots in cell  $r_3$  more or less separated into two distinct, distal pale spot in cell  $r_3$  small and round in center of cell, two pale spots in cell  $m_1$ , only one pale spot each in apices of cells  $m_2$ ,  $CuA_1$  and anal cell, pale spot present behind medial fork; halter infuscated; spermatheca one, pyriform (Wirth and Blanton 1959).

**Distribution:** Costa Rica, Honduras, Panama, and Brazil (Amazonas, Pará, and Rio de Janeiro).

***Culicoides glabellus*** Wirth & Blanton, 1956

(Figures 7, 12D)

*Culicoides glabellus* Wirth and Blanton (1956): 47. Panama.

**Identification:** General appearance, size, and wing markings practically identical to *C. leoni* Barbosa, 1952 and *C. gabaldoni* Ortiz, 1954. There are two spermathecae but in *C. gabaldoni* and *C. leoni* differs by having an antennal ratio of only 0.75 and sensoria present in flagellomere 1, 5, and 6–8. The male genitalia of *C. glabellus* is distinguished by the gradually curving base of the paramere, the swollen stem at the distal part of the straight portion, beyond which the paramere tapers rapidly to a simple, filamentous tip (Wirth and Blanton 1959).

**Distribution:** Honduras to Panamá, Ecuador, Trinidad, and Brazil (Amazonas, Pará, Roraima, and Bahia).

***Culicoides leoni*** Barbosa, 1952 (Figure 7)

*Culicoides leoni* Barbosa (1952): 17 Ecuador.

**Identification:** Sensilla coeloconica on flagellomeres 1, 5–8; wing length 0.63 mm, wing pattern with the caudal extension of the post-stigmatic pale spot not oblique in cell  $r_3$  and usually connected to vein  $M_1$ , smaller pale spot on r-m crossvein, one pale spot in cell  $M_1$  near base, and distal pale spot in anal cell abutting wing margin (Huerta et al. 2012).

**Distribution:** Brazil (Amazonas) and Ecuador.

Subgenus unplaced, *limai* species group

***Culicoides antunesi*** Forattini, 1954

*Culicoides antunesi* Forattini (1954b): 315. Goiás, Brazil.

**Identification:** Eyes narrowly separated, bare; third segment slightly swollen, with a small, deep, sensory pit; sensilla in flagellomeres 1, 5–8 and antennal ratio greater than 1; halter pale; species resembles *C. lobato* Felipe-Bauer 1994, *C. limai* Barretto, 1954 and *C. boliviensis* Spinelli and With, 1984 by general aspects of the wings. *Culicoides antunesi*, *C. lobato* and *C. boliviensis* have similar pale markings on all legs while *C. limai* presents an additional pale spot on subapical portion of hind femur (Forattini 1957; Felipe-Bauer and Quintelas 1994).

**Distribution:** Brazil (Amazonas, Goiás, Mato Grosso do Sul, Minas Gerais, and São Paulo).

***Culicoides galindoi*** Wirth & Blanton, 1953

(Figures 7, 12E)

*Culicoides galindoi* Wirth and Blanton (1953): 73. Panama.

**Identification:** Eyes contiguous, bare; third segment slightly swollen, with a small, shallow, sensory pit; have the pale distal spot in cell  $r_3$  rounded distally, leaving a small dark area in apex of cell; halter pale; spermathecae two, pyriform, subequal (Wirth and Blanton 1959).

**Distribution:** Costa Rica, Panama, and Brazil (Acre, Amazonas, Roraima, and Maranhão).

***Culicoides limai*** Barretto, 1944

(Figures 8, 12F)

*Culicoides limai* Barretto (1944): 99. São Paulo, Brazil.

**Identification:** Small; the much broader third palpal segment; wing as shown in figure, with distal pale spot in cell  $m_1$  broadly abutting wing margin, second radial wholly included in a dark spot, no pale spot straddling vein  $M_2$ , this vein usually dark to apex, apices of veins  $M_1$ ,  $M_2$  dark; subapical pale band on the hind femur and pale apex of the hind tibia; presence of a small ventral lobe on the male parameres (Wirth and Blanton 1959).

**Distribution:** El Salvador to Ecuador, northeastern Argentina, and Brazil (Amapá, Amazonas, Pará, Roraima, Maranhão, Mato Grosso do Sul, Minas Gerais, Rio de Janeiro, São Paulo, and Santa Catarina).


***Culicoides lopesi*** Barretto, 1944

(Figures 8, 12G)

*Culicoides lopesi* Barretto (1944): 102. São Paulo, Brazil.

**Identification:** Wing nearly identical with that of *C. almirantei* Wirth & Blanton, 1959, wing as figured, but the genitalia are different, with a tapering ninth tergum, aedeagus with slender basal arms and slender apex, and parameres with a low ventral lobe and subapical barbs (Wirth and Blanton 1959).

**Distribution:** Panama, Suriname, and Brazil (Acre, Amazonas, Rio de Janeiro, and São Paulo).



**Figure 8.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. limai*. **B.** *C. lopesi*. **C.** *C. santanderi*. **D.** *C. vernoni*. **E.** *C. caprilesi*. **F.** *C. pachymerus*. **G.** *C. aureus*. **H.** *C. castelloni*. **I.** *C. fittkaui*. **J.** *C. guamai* (modified after SimpleMappr, <http://www.simplemappr.net>).

***Culicoides santanderi* Browne, 1980**

(Figure 8)

*Culicoides santanderi* Browne (1980): 536. Colombia.

**Identification:** Eyes narrowly separated, bare; third segment cone-shaped with deep circular sensory pit; sensory pattern in flagelomeres 1, 5–8; halter pale; two spermathecae, subequal, with ducts sclerotized a considerable distance. Antennal proportion, distribution of sensoria, and overall wing pattern place *C. santanderi* within the *limai* group. This species is distinguished from others in that group by the divided poststigmatic pale spot, the posterior one larger and extending proximal to the anterior one (Browne 1980).

**Distribution:** Brazil (Amazonas) and Colombia.

***Culicoides vernoni* Wirth & Blanton, 1973**

(Figures 8, 12H)

*Culicoides vernoni* Wirth and Blanton (1973): 448. Pará, Brazil.

**Identification:** Wing as figured, distal pale spot in cell

$r_3$  large, filling apex of cell with dark area boadering tip of vein  $M_1$  very little expanded apically; fore and mid femora with suapical and all tibiae with sub-basal, narrow pale rings. This species has a nearly identical wing pattern to *C. limai* and *C. galindoi*, but in both related species the distal pale spot in cell  $r_3$  is rounded distally leaving a sizeable dark area at apex of cell. *Culicoides limai* has male genitalia nearly identical with *C. vernoni*, but the hind femur has a distinct subapical pale ring. *Culicoides galindoi* has the hind femur dark to tip, but the male parameres lack the ventral lobe (Wirt and Blanton 1973).

**Distribution:** Costa Rica, Colombia, Bolivia, and Brazil (Amazonas and Pará).

Subgenus unplaced, *pachymerus* species group

***Culicoides caprilesi* Fox, 1952**

(Figure 8)

*Culicoides caprilesi* Fox (1952): 364. Venezuela.

*Culicoides kintzi* Wirth and Blanton (1953): 72. Panama.

**Identification:** Distinguished in pinned mounts by the greater development of the dark mesonotal striae and the much hairier wing on which the dark color pattern is more extensive, dark mark extending from the base of the second radial cell almost to its apex and extending over into cell  $r_3$  as a small rounded dark spot just behind the tip of the second radial cell. In some specimens, the latter dark spot is narrowly connected to the next distal dark area (Wirth and Blanton 1959).

**Distribution:** Colombia, Panama, Venezuela, and Brazil (Amazonas, Pará, and Mato Grosso).

***Culicoides pachymerus* Lutz, 1914**

(Figure 8)

*Culicoides pachymerus* Lutz (1914): 83. –Amazonas, Brazil.

**Identification:** Smaller than *C. uniradialis* Wirth & Blanton, 1953 and *C. caprilesi*, its wing is less hairy than that of *C. caprilesi*, there is no dark spot or connection immediately posterior to the end of the second radial cell as in *C. caprilesi*. The male genitalia of *C. pachymerus* closely resemble those of *C. uniradialis*, with a long ventral lobe present on the parameres, but differing in having a median notch on the ninth tergum, aedeagal arch much broader with slender basal arms and no barbs on the tips of the parameres (Wirth and Blanton 1959).

**Distribution:** Guatemala to Colombia and Brazil (Amazonas).

Subgenus unplaced, *reticulatus* species group

***Culicoides aureus* Ortiz, 1951**

(Figures 8, 12I)

*Culicoides aureus* Ortiz (1951a): 585. Venezuela.

*C. miyamotoi* Wirth and Blanton (1953): 231. Panama.

**Identification:** Wing as figured,  $r_3$  with five pale spots in the proximal half of cell;  $m_1$  with two pale spots, the distal one far from wing margin; AR 0.72–0.76; proboscis short, P/H ratio 0.61–0.70; male with sternite 9 with slightly perceptible posteromedial excavation; aedeagus with basal arch extending to 0.3 of total length; parameres without ventral lobes, distal portion curved, apex flattened (Santarém et al. 2015).

**Distribution:** Panama, Venezuela, Bolivia, Paraguay, Argentina, and Brazil (Amazonas and Goiás).

***Culicoides castelloni* Santarém & Felipe-Bauer, 2015**

(Figure 8)

*Culicoides castelloni* Santarém and Felipe-Bauer (2015): 959. Amazonas, Brazil.

**Identification:** The only species of the *reticulatus* group with sensilla coeloconica on flagellomeres 1 and 9–13 and a tibial comb with six spines (sensilla coeloconica 1, 5–8 or 1, 6–8 and tibial comb with 4 or 5 spines in other species of the group). Third palpal segment long (PR 4.2) with a small, rudimentary sensory pit

(PR 1.6–3.2, with variable sensory pit, in other species of the group). The female of *C. castelloni* has similar wing pattern to *C. reticulatus*, but distinguished by the characters described above and by the presence of two parallel, longitudinal, admedian pale spots, anterior to the prescutellar depression, in *C. reticulatus* with four median anterior yellowish spots arranged as a leaf clover (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas).

***Culicoides fittkaui* Wirth & Blanton, 1973**

(Figures 8, 12J)

*Culicoides fittkaui* Wirth and Blanton 1973: 432. Pará, Brazil.

**Identification:** Eyes nearly contiguous; proboscis short, P/H ratio 0.60–0.63; hind femur dark; wing as shown in figure, with pale spots of the wing interconnected,  $r_3$  with two pale spots, second radial cell with pale lumen,  $m_1$  with two pale spots, the distal one reaching wing margin; aedeagus in male with basal arch extending half of total length; parameres with large ventral lobes, apical portion with lateral fringe of spicules (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Pará).

**\**Culicoides guamai* Wirth & Blanton, 1973**

(Figures 8, 12K)

*Culicoides guamai* Wirth and Blanton (1973): 438. Pará, Brazil.

**Identification:** Proboscis short, P/H ratio 0.71–0.79; hind femur dark; wing as figured, with pale spots of the wing interconnected,  $r_3$  with two pale spots,  $m_1$  with two pale spots, the distal one reaching wing margin, second radial cell short, with dark lumen; aedeagus in male with basal arch extending to 0.6 of total length; parameres with a strong ventral swelling on proximal portion of stem, distal portion with filamentous tip with fringing spines (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Pará).

**New record** — Brazil, Amazonas state, Presidente Figueiredo, Rio Pardo (01°48' S, 060°19' W), VI-2010, VIII-2010, 0001-6 (2 females), CDC light traps, forest fragments, F.A.C. Pessoa collector.


***Culicoides hildebrandoi* Farias, Pereira Junior,**

Felippe-Bauer, Pessoa, Medeiros and Santarém, 2016

(Figures 9, 12L)

*Culicoides hildebrandoi* Farias et al. (2016): 107. Amazonas and Rondônia, Brazil.

**Identification:** Second radial cell in dark spot,  $r_3$  with four sparsely distributed pale spots, r-m crossvein pale; hind femur with subapical pale band; scutum with two anterior submedian clover-leaf shaped spots; 3<sup>rd</sup> palpal segment elongate, slightly swollen, cylindrical, without a sensory pit but with capitate sensilla scattered on the surface cuticle; tergite 9 of male with a posteromedial notch, parameres with slightly sinuous stem, swollen on mid-portion and without a ventral lobe and the basal



**Figure 9.** Map of *Culicoides* species distribution in Amazonas State, Brazil. **A.** *C. hildebrandoi*. **B.** *C. irregularis*. **C.** *C. kuripako*. **D.** *C. lanei*. **E.** *C. paucienfusca*. **F.** *C. profundus*. **G.** *C. pseudoreticulatus*. **H.** *C. rhombus*. **I.** *C. fluviatilis*. **J.** *C. wokei* (modified after SimpleMappr, <http://www.simplemappr.net>).

arch of aedeagus extending two-thirds of total length.

**Distribution:** Brazil (Amazonas and Roraima).

***Culicoides irregularis*** Santarém, Felipe-Bauer & Castellón, 2014

(Figures 9, 12M)

*Culicoides irregularis* Santarém, Felipe-Bauer and Castellón in Santarém et al. (2014): 265. Roraima, Brazil.

**Identification:** Third palpus segment with a multiple, shallow, irregular sensory pit, PR 2.0–2.9; moderately long proboscis, P/H ratio 0.80–0.94; mandible with 16 teeth; scutum with four median anterior yellowish spots arranged as a leaf clover; wing as shown in figure,  $r_3$  with four pale spots,  $m_1$  with two pale spots; two unequal ovoid spermathecae (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Roraima).

***Culicoides kuripako*** Felipe-Bauer, 2010

(Figure 9)

*Culicoides kuripako* Felipe-Bauer in Felipe-Bauer et al. (2010): 863. Amazonas, Brazil.

**Identification:** Pale spots of the wing interconnected;  $r_3$  with two pale spots, poststigmatic pale spot inverted L-shaped, isolating a dark spot behind second radial cell;  $m_1$  with two pale spots, the distal one reaching wing margin;  $CuA_1$  and  $CuA_2$  pale. Male: sternite 9 with deep posteromedial excavation; aedeagus with basal arch extending to half of total length; parameres with ventral lobes, distal portion with lateral fringe of spicules (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Pará).

***Culicoides lanei*** Ortiz, 1950

(Figures 9, 12N)

*Culicoides lanei* Ortiz (1950b): 431. Panama.

**Identification:** Proboscis short, P/H ratio 0.63–0.66; AR 0.72–0.78; wing as figured,  $r_3$  with five pale spots,  $m_1$  with three pale spots, the distal one near wing margin; hind


femur dark, hind tibia dark apically; aedeagus in male with basal arch extending to half the total length; parameres with blade-like ventral lobes, distal portion with lateral fringe with fine spicules (Santarém et al. 2015).

**Distribution:** Mexico, Honduras, Costa Rica, Panama, Venezuela, Trinidad, and Brazil (Amazonas, Pará and Roraima).

***Culicoides paucienfuscatus*** Barbosa, 1947  
(Figures 9, 120)

*Culicoides paucienfuscatus* Barbosa (1947): 23. –Amazonas, Brazil.

**Identification:** Wing as figured, with pale spots of the wing interconnected;  $r_3$  with two pale spots, post-stigmatic pale spot inverted L-shaped, isolating a dark spot behind second radial cell;  $m_1$  with two pale spots,



**Figure 10.** Wing photograph of *Culicoides* species in Amazonas State, Brazil. **A.** *C. efferus*. **B.** *C. pusilloides*. **C.** *C. pusillus*. **D.** *C. freitasi*. **E.** *C. debilipalpis*. **F.** *C. glabrior*. **G.** *C. insinuatus*. **H.** *C. spurius*. **I.** *C. todatangae*. **J.** *C. paraensis*. **K.** *C. quasiparaensis*. **L.** *C. baniwa*. **M.** *C. batesi*. **N.** *C. brownei*. **O.** *C. coutinhoi*. **P.** *C. diabolicus*. **Q.** *C. filariferus*. **R.** *C. flavivenulus*. **S.** *C. foxi*. **T.** *C. franklini*. **U.** *C. fusipalpis*.


the distal one reaching wing margin; CuA<sub>1</sub> pale and CuA<sub>2</sub> dark (Santarém et al. 2015).

**Distribution:** Costa Rica to Peru, Venezuela, Trinidad, Brazil (Acre, Amazonas, Pará, Roraima, Maranhão, Goiás) and Bolivia.

***Culicoides profundus*** Santarém, Felipe-Bauer and Trindade, 2014

(Figures 9, 12P)

*Culicoides profundus* Santarém, Felipe-Bauer and Trindade in Santarém et al. (2014): 266. Pará, Brazil.



**Figure 11.** Wing photograph of *Culicoides* species in Amazonas State, Brazil. **A.** *C. hylas*. **B.** *C. ignacioi*. **C.** *C. insignis*. **D.** *C. lutzi*. **E.** *C. ocumarensis*. **F.** *C. paraignacioi*. **G.** *C. paramaruim*. **H.** *C. plaumanni*. **I.** *C. pseudodiabolicus*. **J.** *C. tidwelli*. **K.** *C. verecundus*. **L.** *C. aldomari*. **M.** *C. bricenoi*. **N.** *C. wallacei*. **O.** *C. alahialinus*. **P.** *C. acotylus*. **Q.** *C. belemensis*. **R.** *C. dasyophrus*. **S.** *C. eublepharus*. **T.** *C. propriipennis*. **U.** *C. fluvialis*.


**Identification:** Third palpal segment swollen, with large, deep sensory pit in middle portion, opening in a small, rounded pore, PR 1.8–2.4; moderately long proboscis, P/H ratio 0.86–0.95; mandible with 16 teeth; scutum with four median anterior yellowish spots arranged as a leaf clover; wing as shown in figure,  $r_3$  with four pale spots,  $m_1$  with two pale spots, the distal one far from wing margin (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Pará).

***Culicoides pseudoreticulatus*** Santarém, Felipe-Bauer & Castellón, 2014  
(Figures 9, 12Q)

*Culicoides pseudoreticulatus* Santarém, Felipe-Bauer and Castellón in Santarém et al. (2014): 268. Roraima, Brazil.

**Identification:** Scutum with four median anterior yellowish spots arranged as a leaf clover; third palpal segment with a shallow, round sensory pit in middle portion, with no well-defined contours on distal half, PR



**Figure 12.** Wing photograph of *Culicoides* species in Amazonas State, Brazil. **A.** *C. leopoldoi*. **B.** *C. tetrathyris*. **C.** *C. benarrochi*. **D.** *C. glabellus*. **E.** *C. galindoi*. **F.** *C. limai*. **G.** *C. lopesi*. **H.** *C. vernoni*. **I.** *C. aureus*. **J.** *C. fittkaui*. **K.** *C. guamai*. **L.** *C. hildebrandoi*. **M.** *C. irregularis*. **N.** *C. lanei*. **O.** *C. paucienfuscatus*. **P.** *C. profundus*. **Q.** *C. pseudoreticulatus*. **R.** *C. rhombus*.

2.0–2.7; proboscis moderately long, P/H ratio 1.0–1.06; mandible with 19 teeth; wing as figured,  $r_3$  with four pale spots;  $m_1$  with two pale spots, the distal one far from wing margin (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Roraima).

***Culicoides rhombus*** Santarém, Felipe-Bauer & Castellón, 2014

(Figures 9, 12R)

*Culicoides rhombus* Santarém, Felipe-Bauer and Castellón in Santarém et al. (2014): 270. Roraima, Brazil.

**Identification:** Third palpal segment rhomboid, with a shallow, well-developed irregular sensory pit that reaches the extreme distal portion of segment, PR 2.5–3.0; proboscis long; P/H ratio 1.08–1.28; scutum with four median anterior yellowish spots arranged as a leaf clover; wing as figured,  $r_3$  with four pale spots,  $m_1$  with two pale areas, the distal one far from wing margin (Santarém et al. 2015).

**Distribution:** Brazil (Amazonas and Roraima).

Subgenus unplaced, *stigmalis* species group

***Culicoides fluviatilis*** (Lutz, 1914)

(Figure 9)

*Johannseniella fluviatilis* Lutz (1914): 82. Amazonas, Brazil.

*Culicoides scorzai* Ortiz (1956): 93. Venezuela.

**Identification:** A large species; wing length 1.12 mm, membrane with evenly scattered, small but distinct, spinelike macrotrichia; CR 0.70; AR 0.80 (Felippe-Bauer and Wirth 1987).

**Distribution:** Colombia, Venezuela, Trinidad Ecuador, Bolivia, and Brazil (Amazonas).

Miscellaneous unplaced species

***Culicoides wokei*** Fox, 1947

(Figure 9)

*Culicoides wokei* Fox (1947): 90. Panama.

*Culicoides aethionotus* Wirth and Blanton, (1955b): 121. Panama.

**Identification:** Eyes narrowly separated, bare; third palpal segment short and swollen, with broad, shallow sensory pit; second radial cell dark, pale spot over r-m crossvein very broad, cell  $r_3$  with three pale spots, a small one at end of second radial cell, an elongate one behind second radial cell sometimes narrowly meeting the pale spot over crossvein and the anterior poststigmatic pale spot, and the third pale spot extensive, nearly filling apex of cell  $r_3$ ; spermathecae two (Wirth and Blanton 1959).

**Distribution:** Costa Rica, Panama, and Brazil (Amazonas).

## DISCUSSION

The richness of *Culicoides* in the state of Amazonas is 6.6% of all species of the genus worldwide and 77% all *Culicoides* species in Brazil (Borkent 2015; Farias et

al. 2016; Santarém and Felipe-Bauer 2016). Catalogs and lists of species are important tools in the study of biodiversity of localities, states, countries, and zoogeographic regions (Santarém and Felipe-Bauer 2016). Our list of species may be helpful in identifying the *Culicoides* fauna of Amazonas, for it gathers together data on the identification and distribution of species occurring in the state. Moreover, we extend the ranges of nine species to the state of Amazonas, including two species are recorded from Brazil for the first time.

Despite the importance *Culicoides* to human health, the *Culicoides* fauna of Amazonas state is still poorly known. Our list of *Culicoides* of Amazonas adds to the knowledge of the diversity of this genus and may be useful for medical entomology, vector epidemiology, and management of these pernicious insects.

## ACKNOWLEDGEMENTS

We thank Dr. Maria Luiza Felipe-Bauer and Maria Clara Alves Santarém for the corrections in to the manuscript. We were financially supported by PROEP ILMD/FAPEAM, Process number 062.01844/2014.

## LITERATURE CITED

- Barbosa, F.A.S. 1947. *Culicoides* (Diptera: Heleidae) da região neotropical. Anais da Sociedade de Biologia de Pernambuco 7: 3–30.
- Barbosa, F.A.S. 1952. Novos subsidios para o conhecimento dos *Culicoides* neotropicos (Diptera: Heleidae) Publicações Avulsas do Instituto Aggeu Magalhães, Recife 2: 3, 11–41.
- Barretto, M.P. 1944. Sobre o género *Culicoides* Latreille, 1809, com a descricao de três novas espécies (Diptera, Ceratopogonidae). Anais da Faculdade de Medicina (da Universidade) de Sao Paulo 20: 89–105, pls. 1–4.
- Beck, E.C. 1957. Two new species of *Culicoides* from Florida (Diptera: Heleidae). Florida Entomologist 40: 103–105.
- Borkent, A. 2015. World species of biting midges (Diptera: Ceratopogonidae). Accessed at <http://www.inhs.illinois.edu/files/1114/2384/5200/CeratopogonidaeCatalog.pdf>, 20 August 2015.
- Borkent, A. and G.R. Spinelli. 2007. Neotropical Ceratopogonidae (Diptera: Insecta). Sofia/Moscow: Pensoft Publishers. 198 pp.
- Browne, J.E. 1980. New species of biting midges of the genus *Culicoides* from Colombia and the first description of the male of *C. florenciae* (Diptera: Ceratopogonidae). Journal of Medical Entomology 17: 533–544. doi: [10.1093/jmedent/17.6.533](https://doi.org/10.1093/jmedent/17.6.533)
- Castellón, E.G. and M.L. Felipe-Bauer. 2015. Classificação e distribuição das espécies de *Culicoides* (Diptera: Ceratopogonidae) na Amazônia brasileira; pp. 30–66, in: E.G. Castellón and R.S. Veras (eds.). Maruins (Ceratopogonidae: *Culicoides*) na Amazônia brasileira, 1<sup>st</sup> edition. Manaus: Editora INPA. 146pp.
- Clastrier, J. 1968. Deux ceratopogonides nouveaux de la Guyane Française. Archives de l'Institut Pasteur de la Guyane Française et de L'Inini 21: 85–92.
- Clastrier, J. 1971. Deux nouveaux *Culicoides* (Diptera, Ceratopogonidae) de la Guyane Française. Annales de Parasitologie 46: 285–294.
- Costa Lima, A. da. 1937. Chave das especies de *Culicoides* da regio neotropical (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 32: 411–422. doi: [10.1590/S0074-02761937000300003](https://doi.org/10.1590/S0074-02761937000300003)
- Farias, E.S., A.M. Pereira-Júnior, J.A. Feijó, F.A.C. Pessoa and

- J.F. Medeiros. 2015. Hematophagous biting midges (Diptera: Ceratopogonidae) from Tefé municipality, Amazonas state, Brazil. Check List 11(4): 1676. doi: [10.15560/11.4.1676](https://doi.org/10.15560/11.4.1676)
- Farias, E.S., A.M. Pereira Júnior, M.L. Felipe-Bauer, F.A.C. Pessoa, J.F. Medeiros and M.C.A. Santarém. 2016. *Culicoides hildebrandoi*, a new species of the *reticulatus* species group from the Brazilian Amazon region (Diptera, Ceratopogonidae). ZooKeys 571: 105–111 doi: [10.3897/zookeys.571.7341](https://doi.org/10.3897/zookeys.571.7341)
- Felippe-Bauer, M.L. and A.R. Quintelas. 1994. *Culicoides lobatoui*, a new Brazilian biting midge of the limai group (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 89: 25–27. [10.1590/S0074-02761994000100005](https://doi.org/10.1590/S0074-02761994000100005)
- Felippe-Bauer, M.L., A.G. Cáceres, C. S. Silva, W. Valderrama-Bazan, A. Gonzales-Perez and J.M. Costa. 2008. Description of *Culicoides pseudoheliconiae* sp. n. from Peruvian Amazon and revalidation of *Culicoides contubernalis* Ortiz & Leon (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 103: 259–262. doi: [10.1590/S0074-02762008005000010](https://doi.org/10.1590/S0074-02762008005000010)
- Felippe-Bauer, M.L., C.P. Damasceno, V. Py-Daniel and G.R. Spinelli. 2009. *Culicoides baniwa* sp. nov. from the Brazilian Amazon region with a synopsis of the *hylas* species group (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 104: 851–857. doi: [10.1590/S0074-02762009000600005](https://doi.org/10.1590/S0074-02762009000600005)
- Felippe-Bauer, M.L., C.P. Damasceno, R. Lopes da Trindade and V. Py-Daniel. 2010. A new *Culicoides* (Diptera: Ceratopogonidae) of the *reticulatus* species group from Brazilian Amazon region. Memórias do Instituto Oswaldo Cruz 105: 863–865. doi: [10.1590/S0074-02762010000700004](https://doi.org/10.1590/S0074-02762010000700004)
- Felippe-Bauer, M.L., T.N. Silva and R.L. Trindade. 2013. New *Culicoides* Latreille of the subgenus *Mataemyia* Vargas from Pará, Brazil (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 108: 54–58. doi: [10.1590/S0074-02762013000100009](https://doi.org/10.1590/S0074-02762013000100009)
- Felippe-Bauer, M.L. and W.W. Wirth. 1987. A new *Culicoides* of the *stigmatis* group (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 82: 415–419. doi: [10.1590/S0074-02761987000300014](https://doi.org/10.1590/S0074-02761987000300014)
- Floch, H. and E. Abonnenc. 1942a. Cératopogonidés hématophages de la Guyane Française — II. Publication de l'Institut Pasteur de la Guyane et du Territoire de L'Inini 49: 1–5.
- Forattini, O.P. 1954a. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) do Brasil Central. Arquivos da Faculdade de Higiene e Saude Publica da Universidade de Sao Paulo 8: 189–192. doi: [10.11606/issn.2358-792X.v8i2p189-192](https://doi.org/10.11606/issn.2358-792X.v8i2p189-192)
- Forattini, O.P. 1954b. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) do estado de Goias, Brasil. Folia Clinica et Biologica 21: 315–320.
- Forattini, O.P. 1956a. A new *Culicoides* from Panama (Diptera, Ceratopogonidae). Proceedings of the Entomological Society of Washington 58: 35–36. <http://biodiversitylibrary.org/page/16196666>
- Forattini, O.P. 1956b. Nova espécie de *Culicoides* (Diptera, Ceratopogonidae) de Sao Vicente, estado de Sao Paulo, Brasil. Arquivos da Faculdade de Higiene e Saude Publica da Universidade de Sao Paulo 10: 81–84.
- Forattini, O.P. 1957. *Culicoides* da Regiao Neotropical (Diptera, Ceratopogonidae). Arquivos da Faculdade de Higiene e Saúde Pública da Universidade de São Paulo 11: 159–526. doi: [10.11606/issn.2358-792X.v11i2p161-526](https://doi.org/10.11606/issn.2358-792X.v11i2p161-526)
- Fox, I. 1946. A review of the species of biting midges or *Culicoides* from the Caribbean region (Diptera: Ceratopogonidae). Annals of the Entomological Society of America 39: 248–258. doi: [10.1093/aesa/39.2.248](https://doi.org/10.1093/aesa/39.2.248)
- Fox, I. 1947. Two new Central American biting midges or *Culicoides* (Diptera: Ceratopogonidae). Kuba 3: 90–91.
- Fox, I. 1948. *Hoffmania*, a new subgenus in *Culicoides* (Diptera: Ceratopogonidae). Proceedings of the Biological Society of Washington 61: 21–28. <http://www.biodiversitylibrary.org/part/43305>
- Fox, I. and W.A. Hoffman. 1944. New Neotropical biting sand flies of the genus *Culicoides* (Diptera: Ceratopogonidae). Puerto Rico Journal of Public Health and Tropical Medicine 20: 108–111.
- Fox, I. 1952. Six new Neotropical species of *Culicoides* (Diptera: Ceratopogonidae). Annals of the Entomological Society of America 45: 364–368. doi: [10.1093/aesa/45.3.364](https://doi.org/10.1093/aesa/45.3.364)
- Goeldi, E. 1905. Os mosquitos no Pará. Reunião de quatro trabalhos sobre os mosquitos indigenas, principalmente as especies que molesta o homem. Memórias do Museu Goeldi (Museu Paraense) de Historia Natural e Ethnographie 4: 1–154, 5 pls. <http://biodiversitylibrary.org/page/8677658>
- Hoffman, W.A. 1925. A review of the species of *Culicoides* of North and Central America and the West Indies. American Journal of Hygiene 5: 274–301.
- Hoffman, W.A. 1939. *Culicoides filariferus*, new species. Puerto Rico Journal of Public Health and Tropical Medicine 15: 172–176.
- Kieffer, J.J. 1917. Chironomides d'Amérique conservés au Musée National Hongrois de Budapest. Annales Historico-Naturales Musei Nationalis Hungarici 15: 292–364.
- Kieffer, J.J. 1921. Sur quelques diptères piqueurs de la tribu des Ceratopogoninae. Archives de l'Institut Pasteur de l'Afrique du Nord 1: 107–115.
- Lutz, A. 1913. Contribuição para o estudo das ceratopogoninas hematofagas do Brasil. Memórias do Instituto Oswaldo Cruz 5: 45–73, pls. 6–8. doi: [10.1590/s0074-02761913000100005](https://doi.org/10.1590/s0074-02761913000100005)
- Lutz, A. 1914. Contribuição para o conhecimento das ceratopogoninas do Brasil. Memórias do Instituto Oswaldo Cruz 6: 81–99, pls. 8–9. doi: [10.1590/S0074-02761914000200003](https://doi.org/10.1590/S0074-02761914000200003)
- Macfie, J.W.S. 1940. A report on a collection of Ceratopogonidae (Diptera) from British Guiana. The Entomologist's Monthly Magazine 76: 23–32.
- Macfie, J.W.S. 1948. Some species of *Culicoides* (Diptera, Ceratopogonidae) from the state of Chiapas, Mexico. Annals of Tropical Medicine and Parasitology. 42: 67–87.
- Mellor, P.S., J. Boorman and M. Baylis 2000. *Culicoides* biting midges: their role as arbovirus vectors. Annual Review of Entomology 45: 307–340. doi: [10.1146/annurev.ento.45.1-307](https://doi.org/10.1146/annurev.ento.45.1-307)
- Ortiz, I. 1950a. Estudios en *Culicoides* V. Informes sobre una nueva especie y lista de los machos cuyas genitalias son conocidas. Revista de Sanidad y Asistencia Social 15: 461–465.
- Ortiz, I. 1950b. Estudios en *Culicoides* IV. Revisión de las especies americanas del sub-género *Hoffmania* Fox 1948, con la descripción de dos nuevas especies. Revista de Sanidad y Asistencia Social 15: 437–460.
- Ortiz, I. 1950c. Estudios en *Culicoides* II. Diptera, Ceratopogonidae. *Culicoides lanei* n. sp. de Panama. Revista de Sanidad y Asistencia Social 15: 431–433.
- Ortiz, I. 1951a. Estudios en *Culicoides* (Diptera, Ceratopogonidae). IX. Sobre los caracteres diferenciales entre *Culicoides paraensis* (Goeldi, 1905), *C. stellifer* (Coquillett, 1901), y *C. lanei* (Ortiz, 1950). Descripción de cuatro nuevas especies con la redescipción de algunas otras poco conocidas. Revista de Sanidad y Asistencia Social 16: 573–591.
- Ortiz, I. 1951b. Estudios en *Culicoides* (Diptera, Ceratopogonidae). 6. *Culicoides bricenoi* n. sp. Boletín del Laboratorio de la Clínica "Luis Razetti" 11: 442–448.
- Ortiz, I. 1953. Nueva contribución al conocimiento de los caracteres morfológicos externos de las hembras americanas del género *Culicoides* LTR. (Diptera, Ceratopogonidae) con una espermoteca. Descripción de dos nuevas especies de Venezuela: *Culicoides transferrans* (C. *oublepharus* [sic] Ortiz, 1952) y *Culicoides mirsae*. Revista de Sanidad y Asistencia Social 18: 797–806.
- Ortiz, I. 1956. Sobre un pequeño diptero de la region del Auyantepuy. Boletín Venezolano de Laboratorio Clínico 1: 93–96.

- Ortiz, I. and L.A. León. 1955. Los *Culicoides* (Diptera: Ceratopogonidae) de la Republica del Ecuador. Boletín de Informaciones Científicas Nacionales 67: 564–594.
- Ortiz, I. and M. Mirsa. 1951. Estudios en *Culicoides*. Descripción de dos nuevas especies: *Culicoides avilaensis* y *C. discrepans*, y del macho de *C. leopoldoi* Ortiz, 1951. Redescrición de *C. limai* Barretto, 1944, *C. baueri* Hoffman 1925, *C. lichyi* Floch & Abonnenc, 1949, y *C. pusillus* Lutz, 1913. Revista de Sanidad y Asistencia Social 16: 593–605.
- Ortiz, I. and M. Mirsa. 1952. Sobre las especies Americanas del género *Culicoides* Latr., (Diptera, Ceratopogonidae) con una espermateca. Acta Científica Venezolana 2: 125–128.
- Poe, F. 1853. Memórias sobre la historia natural de la Isla de Cuba, acompañadas de sumarios latinos y extractos en francés. Habana: Impr. de Barcina. 1: 463 pp, 34 pls. doi: [10.5962/bhl.title.2485](https://doi.org/10.5962/bhl.title.2485)
- Ronderos, M.M. and G.R. Spinelli. 1995. Redescrición de *Culicoides lahillei* y descripción de *Culicoides ichesi*, n. sp. de la Argentina, Paraguay y Uruguay (Diptera: Ceratopogonidae). Neotropica 41: 77–81.
- Santarém, M.C.A., R.L. Trindade, T.N. Silva, E.G. Castellón, C.A. Mello-Patiu and M.L. Felipe-Bauer. 2014. New Neotropical *Culicoides* and redescription of *Culicoides reticulatus* Lutz (Diptera: Ceratopogonidae). Zootaxa 3795: 255–274. doi: [10.11646/zootaxa.3795.3.2](https://doi.org/10.11646/zootaxa.3795.3.2)
- Santarém M.C., E.S. Farias and M.L. Felipe-Bauer. 2015. *Culicoides castelloni* sp. nov. from the Brazilian Amazon region with a revision of the *reticulatus* species group (Diptera, Ceratopogonidae). Annals of the Brazilian Academy of Sciences 87: 955–72. doi: [10.1590/0001-3765201520140517](https://doi.org/10.1590/0001-3765201520140517)
- Santarém M.C. and M.L. Felipe-Bauer. 2016. Brazilian species of biting midges — espécies de maruins do Brasil (Diptera: Ceratopogonidae). Accessed at [https://portal.fiocruz.br/sites/portal.fiocruz.br/files/documentos/2016\\_brazilian\\_species\\_of\\_biting\\_midges\\_final.pdf](https://portal.fiocruz.br/sites/portal.fiocruz.br/files/documentos/2016_brazilian_species_of_biting_midges_final.pdf), 19 April 2016.
- Shevchenko, A.K. 1977. Bloodsucking midges. Fauna Ukraini 13: 1–251. [In Russian]
- Spinelli, G.R., E.C. Greiner and W.W. Wirth. 1993. The Neotropical bloodsucking midges of the *Culicoides guttatus* group of the subgenus *Hoffmania* (Diptera: Ceratopogonidae). Contributions of the American Entomological Institute 27: 1–91.
- Spinelli, G.R., M.M. Ronderos, P.I. Marino, D.S. Carrasco and R.L. Menezes Ferreira. 2007. Description of *Culicoides (Mataemyia) felippebauerae* sp. n., *Forcipomyia musae* immatures in Brazilian Amazonia (Diptera: Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 102: 659–669. doi: [10.1590/S0074-02762007005000083](https://doi.org/10.1590/S0074-02762007005000083)
- Tavares, O. and R.A. Ruiz. 1980. Sobre duas espécies novas do genero *Culicoides* Latreille, 1809, do Brasil (Diptera, Ceratopogonidae). Memórias do Instituto Oswaldo Cruz 75: 27–32. doi: [10.1590/S0074-02761980000100002](https://doi.org/10.1590/S0074-02761980000100002)
- Tavares, O. and M.A. de Souza. 1978. Sobre tres espécies novas do genero *Culicoides* Latreille, 1809 do Brasil (Diptera, Ceratopogonidae). Revista Brasileira de Biologia 38: 619–624.
- Trindade, R.L. and I.S. Gorayeb. 2005. Maruins (Ceratopogonidae: Diptera) do estuário do rio Pará e do litoral do estado do Pará, Brasil. Entomologia y Vectores 12: 61–74. doi: [10.1590/S0328-03812005000100005](https://doi.org/10.1590/S0328-03812005000100005)
- Trindade, R.L. and I.S. Gorayeb. 2010. Maruins (Diptera: Ceratopogonidae: *Culicoides*), após a estação chuvosa, na Reserva de Desenvolvimento Sustentável Itatupã-Baquiá, Gurupá, Pará, Brasil. Revista Pan-Amazônica de Saúde 1: 121–130. doi: [10.5123/S2176-62232010000200015](https://doi.org/10.5123/S2176-62232010000200015)
- Vargas, L. 1954. Dos nuevas especies de *Culicoides* Mexicanos (Diptera: Heleidae). Revista del Instituto de Salubridad y Enfermedades Tropicales 14: 25–32.
- Vargas, L. 1960. The subgenera of *Culicoides* of the Americas (Diptera: Ceratopogonidae). Revista de Biología Tropical 8: 35–47.
- Waller, J. R., Lancelot, P.C. Lefevre and M. Kremer. 1990. Note faunistique sur *Culicoides* de Guyane (Diptera: Ceratopogonidae). Revue d'Élevage et de Médecine Vétérinaire des Pays Tropicaux 43(3): 349–364.
- Wirth, W.W. and F.S. Blanton. 1959. Biting midges of the Genus *Culicoides* from Panama (Diptera: Heleidae). Proceedings of the United States National Museum 109(1959): 237–482. doi: [10.5479/si.00963801.109-3415.237](https://doi.org/10.5479/si.00963801.109-3415.237)
- Wirth, W.W. and P. Barreto. 1978. New species of *Culicoides* biting midges (Diptera: Ceratopogonidae) from Colombia. Journal of Medical Entomology 14: 553–564. doi: [10.1093/jmedent/14.5.553](https://doi.org/10.1093/jmedent/14.5.553)
- Wirth, W.W. and F.S. Blanton. 1953. Studies in Panama *Culicoides* (Diptera: Heleidae): I, Descriptions of six new species. Journal of the Washington Academy of Sciences 43: 69–77. doi: [10.2307/3273942](https://doi.org/10.2307/3273942)
- Wirth, W.W. and F.S. Blanton. 1955. Studies in Panama *Culicoides* (Diptera, Heleidae) IV. Descriptions of three new species. Bulletin of the Brooklyn Entomological Society 50: 100–106.
- Wirth, W.W. and F.S. Blanton. 1956. Studies in Panama *Culicoides* (Diptera, Heleidae). IX. Two new species related to *leoni* Barbosa and *reevesi* Wirth. Bulletin of the Brooklyn Entomological Society 51: 45–52.
- Wirth, W.W. and F.S. Blanton. 1968. A revision of the Neotropical biting midges of the *hylas* group of *Culicoides* (Diptera: Ceratopogonidae). Florida Entomologist 51: 201–215. doi: [10.2307/3493420](https://doi.org/10.2307/3493420)
- Wirth, W.W. and F.S. Blanton. 1973. A review of the maruins or biting midges of the genus *Culicoides* (Diptera: Ceratopogonidae) in the Amazon Basin. Amazoniana 4: 405–470.
- Wirth, W.W. and N. Marston. 1968. A method for mounting small insects on microscope slides in Canada balsam. Annals of the Entomological Society of America 61: 783–784. doi: [10.1093/aesa/61.3.783](https://doi.org/10.1093/aesa/61.3.783)

**Author contributions:** ESF participated in sample design, identified the insects, and co-wrote the text; JFA mounted the insects and co-wrote the article; FACP participated in sample design, collected insects, and co-wrote the text.

**Received:** 17 November 2015

**Accepted:** 25 April 2016

**Academic editor:** Fabio Silva