



Birds of the Parque Ecológico Lagoa da Fazenda, Sobral, Ceará state, northeastern Brazil

Ana Carolina Brasileiro^{1,2,6}, Frede Lima-Araujo¹, Jamile Aragão Alcântara¹, Alano Sousa Martins Pontes¹, José André Neto³, Ricardo de Oliveira Tavares¹, Amanda Souza Silvino⁴ & Jeferson Vizentin-Bugoni⁵

¹Universidade Estadual Vale do Acaraú, Avenida da Universidade, 850 CEP 62040-370, Campus da Betânia, Sobral, CE, Brazil

²Current address: Universidade Federal do Ceará, Programa de Pós-graduação em Ecologia e Recursos Naturais, CEP 60455-760, Fortaleza, CE, Brazil

³Rua Castro Alves, 316, Alto da Brasília, CEP 62041320, Sobral, CE, Brazil

⁴Universidade Estadual de Campinas, Programa de Pós-graduação em Ambiente e Sociedade, CEP 13083-970, Campinas, SP, Brazil

⁵Universidade Estadual de Campinas, Programa de Pós-graduação em Ecologia, CEP 13083-970, Campinas, SP, Brazil

⁶Corresponding author. E-mail: carolbrmelo@hotmail.com

Abstract: Natural areas within cities are important as they contribute to maintain biodiversity and the functioning of ecosystems. In Ceará state, inventories of birds in human-impacted areas are still scarce. Here, we inventory bird species and estimate the species richness at Parque Ecológico Lagoa da Fazenda, an urban park in the municipality of Sobral, in northeastern Brazil. We found 82 bird species, of which 16 breed in the area, three are endemics, and three others are introduced. Despite the moderate species richness detected, richness estimators revealed that sampling was sufficient to detect most species. To maintain or even increase local species richness, we suggest the cessation of urban expansion within the park, sewage dumping, and filling of the wetland, as well as further planting of native vegetation.

Key words: anthropic avifauna; Brazilian semiarid ecosystems; breeding areas; bird diversity; Caatinga; inventory; species richness

INTRODUCTION

Among the main threats to the bird diversity in Brazil are overhunting, introduced species, pollution and, remarkably, habitats loss and fragmentation (MARINI & GARCIA 2005), which are especially important in densely populated areas around cities. Along with decreases in biodiversity due to urban expansion, several ecosystem services can also be affected, such as pollination and diseases control, which ultimately decreases quality of life for humans (LOBADA & DE ANGELIS 2005). An alternative scenario is the maintenance of natural areas within urban matrices in order to harbor biodiversity and provide several direct benefits, such as water drainage, control of air and noise pollution, and recreational areas (MATSUMOTO et al. 2012). These benefits are even more remarkable when wetlands are associated to urban natural areas

in arid regions as such aquatic ecosystems present high primary productivity and often high vegetation heterogeneity (KEDDY 2000) and therefore increase bird diversity, for instance (BRANCO 2007).

In South America, one of the most arid biogeographic provinces is the Caatinga, which covers 844,453 km² in northeastern Brazil (IBGE 2004). In this region, annual rainfall is typically low and seasonal, with evapotranspiration rates usually overcoming precipitation and, hence, promoting water deficit. Vegetation is xerophytic, with short, spiny, and leafless trees or treelets dominant (Caatinga *strictu sensu*), although scarce forest patches can be found in some humid areas of hills (PRADO 2003). Caatinga also presents endemic fauna adapted to the scarcity of water. Among the birds, 510 species are present if all phytophysiognomies are considered (SILVA et al. 2003), and of these species, 348 can be found within Caatinga *strictu sensu* (PACHECO 2004); 23 of them are restricted or nearly exclusive to Caatinga (OLMOS et al. 2005).

Despite of the high bird diversity in the Caatinga and their role to the functioning of the ecosystem by acting, for example, on pollination, seed dispersal, and pest control, at least 20 species are under risk of extinction (MMA 2002). The fact that only less than 2% of Caatinga is under integral protection worsens this scenario (MACIEL 2010). In this sense, providing species inventories is an important step towards a better understanding of the regional distributions of species, and these data may be valuable to drive conservation efforts. Even though the state of Ceará has been explored by ornithologists for many years (see PACHECO 2004), systematic bird inventories are still scarce and focused mainly on highland areas (NASCIMENTO 2000; NASCIMENTO et al. 2000; FARIA et al. 2005; NASCIMENTO et al. 2005; ALBANO AND GIRÃO 2008) and arboreal Caatinga (i.e., 'Caatinga arbórea e




Figure 1. Location of Sobral (black dot) in Ceará state, northeastern Brazil (A) and satellite image showing the seven sampling points in the Parque Ecológico Lagoa da Fazenda (B). B was adapted from Google Earth™.

arbustiva'; OLMOS et al. 2005). Notably, avifaunas from anthropic areas have been poorly studied in Ceará state to date, which limits the understanding on how urbanization influences the regional bird diversity.

In this context, we provide an inventory of the avifauna of Parque Ecológico Lagoa da Fazenda, a suburban park adjacent to Sobral city, and estimate the species richness. In addition, we report evidences of breeding, and identify the endemic Caatinga species occurring in the park. Finally, we discuss the potential effects of urbanization on the local avifauna and suggest management strategies to maintain, or even increase, the local bird diversity.

MATERIAL AND METHODS

Study area

Data were collected at Parque Ecológico Lagoa da Fazenda (PELF; 03°40'52" S, 40°20'39" W; 70 m above sea level), in the municipality of Sobral, Ceará state (Figure 1). This park was implemented in 1991 (decree 21.303) and covers around of 19 ha (Ceará 1991). Regional climate presents mean annual temperature around 28°C and annual precipitation lower than 750 mm, which is restricted to few months per year. The vegetation of the region is composed by open and bushy caatingas ('Caatinga Arbustiva Aberta'), forest with

Carnauba Wax Palm (*'Floresta Mista Dicotilo-Palmácea'*), deciduous forest with spiny species (*'Floresta Caducifólia Espinhosa'*) and, at higher altitudes, tropical semideciduous forest (*'Floresta Subcaducifólia Tropical Pluvial'*) (IPECE 2013; NIMER 1972). The hydrographic system encompasses three rivers (Acaraú, Aracatiáçu and Jaibaras) that, along with '*Serra da Meruoca*' range, influence the local climate (LIMA & ASSIS 2005).

PELF encompasses a central lake surrounded by emergent or floating macrophytic vegetation. Drier terrains are covered by sparse, mostly exotic trees and treelets species, such as *Pithecellobium dulce* (Roxb.) Benth., but native species are also present such as *Copernicia prunifera* (Miller) H. E. Moore and *Mimosa tenuiflora* (Willd.) Poir. Historically, the PELF has been widely used for recreational purposes but these activities have decreased due to sewage disposal into the lake (AGUIAR JÚNIOR 2005).

Data collection and analysis

Data collection was from May 2014 to April 2015 within seven circular plots each with a radius of 35 m and placed at least 70 m apart (Figure 2). Each plot was sampled twice a month for 15 minutes, starting around 6:30 h, and totaling 42 h of observations. During each 15-minute sampling period, all bird species were identified and the number of individuals was counted. Binocular and photographic cameras (30 × zoom) were used to record birds and specialized field guides were used for species identification (Gwynne et al. 2003; Corbo et al. 2013; Sigrist 2013). Species were documented by voucher photos or voice recordings which were deposited at Wikiaves (<http://www.wikiaves.com.br>), an open-access on-line repository. Species names follow the Brazilian Ornithological Records Committee (CBRO 2014).

Point counts were distributed in order to cover all the representative environments within the PELF (Figures 1 and 2): Point 1 is covered mostly by tall aquatic grasses which are occasionally removed to feed cattle; Point 2 borders a little lake surrounded by tall aquatic grasses; Point 3 covers the interface between a flooded area covered by grasses and a drier terrain with scattered trees; Point 4 borders the main lake which is often covered by dense patches of the floating macrophyte *Eichhornia crassipes* (Mart.) Solms, which changes its location within the lake according to the wind direction. The water level is seasonally variable in both lakes (Points 2 and 4) but neither lake was observed totally dry; Point 5 is a drier terrain covered by scarce short grasses and planted exotic trees such as *P. dulce*, *Azadirachta indica* A. Juss., and *Libidibia ferrea* (Mart. ExTul.) L.P.Queiroz; Point 6 presents a patch of *Cocos nucifera* L. which were planted along the shore of the larger lake where *E. crassipes* and *Ludwigia octovalvis* (Jacq.) P.H.Raven treelets occur; Point 7 is seasonally flooded and presents scattered individuals of *Mimosa tenuiflora*, *C. prunifera*, and *P. dulce*.

Species richness was estimated by Chao 1 and Jackknife 1 (MAGURRAN 2004) using the EstimateS 9.1.0 software




Figure 2. Environmental features and vegetation at each of the seven sampling points in Parque Ecológico Lagoa da Fazenda. Photos by A. C. Brasileiro. See Figure 1 for location within the park.

(COLWELL 2011). We defined sampling as the list of species and their abundances at all seven points summed per sampling day, so that a month had two samples. Species recorded outside the sampling intervals were added to the inventory, but were not considered for the richness estimation. To check sampling sufficiency, we plotted accumulated species richness and mean estimated richness by both metrics along the gradient of sampling accumulation (Figure 3).

Evidences of breeding were collected from January 2014 to April 2015. As evidence, we considered active nests (or those under construction), nestlings, or fledglings (i.e., individuals with low dispersal ability), and/or adults carrying feeding items. Endangered species were classified according BIRDLIFE INTERNATIONAL (2015) and Caatinga endemics, according OLMOs et al. (2005).

RESULTS

We recorded 82 species at the Parque Ecológico Lagoa da Fazenda, of which 69 (84.0% of the total) species were documented by photographs or voice recordings (Table 1). *Dendrocygna viduata*, *Nomonyx dominica*, *Tyto furcata*,

Aramides cajaneus, and *Sturnella superciliaris* were only observed outside the sampling period. The most species-rich families were Tyrannidae (10 species) and Thraupidae, Ardeidae and Rallidae (seven species each). Species richness at PELF was estimated between 88.14 ± 8.23 (Chao1) and 94.25 ± 3.97 species (Jackknife 1). None of the species is currently classified as globally endangered. Three species are endemic to the Caatinga: *Pseudoseisura cristata*, *Agelaioides fringillarius*, and *Paroaria dominicana*. Three species are introduced: *Passer domesticus*, *Estrilda astrild*, and *Columba livia*.

Evidence for breeding was found for 16 species (19.5% of the total; Table 2). Curiously, *Certhiaxis cinnamomeus* and *Pseudoseisura cristata* used plastic materials to build their nests (Figure 4A, B). Remarkably, a nest of *Porzana flavigaster* was found in January 2014 containing five cream-colored eggs (Figure 4C, D).

Below, we list characteristics used to identify species that are either rare to Ceará or potentially confused with similar species:

- *Porzana flavigaster*: black bill; brown iris; pale yellow legs; yellowish throat and breast; body sides black and white stripes; black crown; marked upper ocular.
- *Pardirallus maculatus*: long yellow-greenish bill with a red spot in the basis; reddish legs; red iris; black and white spots or stripes in the neck, breast and belly; black and brownish spots in the back with greenish bright.
- *Laterallus melanophaius*: dark bill with greenish in the inner basis; dark olive legs; brownish upperparts; ferruginous face and neck-sides; white throat, breast and abdomen; back and white barring on the flanks.
- *Laterallus exilis*: black bill with greenish in the inner basis; light grey face sides, neck and breast; white-barrred belly; notable ferrugineous nape and upper neck.
- *Ixobrychus exilis*: straight long yellow bill; blackish cape (male); pale tawny-chestnut face and neck-sides; white throat; light brownish neck with white vertical stripes;




Figure 3. Observed species accumulation curve (square) and estimated by Chao 1 (triangles) and Jackknife 1 (dots) for 28 samples.

light brownish underparts and legs; black back.

- *Botaurus pinnatus*: straight long yellowish bill; white throat; pale yellow and black barred head and neck, turning into thicker irregular bars on the wings and back.
- *Polytmus guainumbi*: long black bill, slightly decurved (reddish upper mandible in males); golden brighten green body; white upper ocular stripe; black ocular stripe; whitish belly.
- *Nomonyx dominica*: male: blue bill with black tip; black face; white periocular ring; dark brown body with black spots; female: plumbeous bill; greyish body with irregular small black dots in the breast; two black stripes in the face with light brown background.
- *Ardea alba*: long straight yellow bill; yellow iris; black legs and feet; entirely white body; bigger than any other white heron.
- *Aramides cajaneus*: yellow bill with a greenish tip; red iris; yellow legs; entirely grey head and neck; rufous breast and belly; brown back and short darker tail.
- *Tringa solitaria*: long straight yellow bills with black tip;

Table 1. Bird species recorded in Parque Ecológico Lagoa da Fazenda, Ceará state, northeastern Brazil. Asterisk indicates species recorded only outside of the sampling periods. Species with several records within the park were demarcated with a general geographical coordinate ($03^{\circ}40'52.98''$ S, $040^{\circ}20'39.51''$ W).

| Order/family/species | Photo or voice vouchers | Latitude | Longitude |
|---|-------------------------|--------------------------|---------------------------|
| Anseriformes Linnaeus, 1758 | | | |
| Anatidae Leach, 1820 | | | |
| * <i>Dendrocygna viduata</i> (Linnaeus, 1766) | wikiaves.com/1663415 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| * <i>Nomonyx dominica</i> (Linnaeus, 1766) | | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| Podicipediformes Fürbringer, 1888 | | | |
| Podicipedidae Bonaparte, 1831 | | | |
| <i>Tachybaptus dominicus</i> (Linnaeus, 1766) | wikiaves.com/1668651 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| <i>Podilymbus podiceps</i> (Linnaeus, 1758) | wikiaves.com/1219037 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| Pelecaniformes Sharpe, 1891 | | | |
| Ardeidae Leach, 1820 | | | |
| <i>Tigrisoma lineatum</i> (Boddaert, 1783) | wikiaves.com/1666016 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| <i>Botaurus pinnatus</i> (Wagler, 1829) | wikiaves.com/1663414 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| <i>Ixobrychus exilis</i> (Gmelin, 1789) | wikiaves.com/1304464 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |
| <i>Butorides striata</i> (Linnaeus, 1758) | wikiaves.com/1689417 | $03^{\circ}40'52.98''$ S | $040^{\circ}20'39.51''$ W |

Continued

Table 1. Continued.

| Order/family/species | Photo or voice vouchers | Latitude | Longitude |
|---|--------------------------------|-----------------|------------------|
| <i>Bubulcus ibis</i> (Linnaeus, 1758) | wikiaves.com/1662738 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Ardea alba</i> Linnaeus, 1758 | | 03°40'52.98" S | 040°20'39.51" W |
| <i>Egretta thula</i> (Molina, 1782) | wikiaves.com/1666015 | 03°40'52.98" S | 040°20'39.51" W |
| Cathartiformes Seebold, 1890 | | | |
| Cathartidae Lafresnaye, 1839 | | | |
| <i>Coragyps atratus</i> (Bechstein, 1793) | wikiaves.com/1689356 | 03°40'38.66" S | 040°20'31.74" W |
| <i>Cathartes aura</i> (Linnaeus, 1758) | wikiaves.com/1677964 | 03°40'38.96" S | 040°20'30.95" W |
| <i>Cathartes burrovianus</i> Cassin, 1845 | wikiaves.com/1662724 | 03°40'52.98" S | 040°20'39.51" W |
| Accipitriformes Bonaparte, 1831 | | | |
| Accipitridae Vigors, 1824 | | | |
| <i>Geranospiza caerulescens</i> (Vieillot, 1817) | wikiaves.com/1663487 | 03°40'41.78" S | 040°20'30.34" W |
| <i>Rupornis magnirostris</i> (Gmelin, 1788) | | 03°40'52.98" S | 040°20'39.51" W |
| Gruiformes Bonaparte, 1854 | | | |
| Aramidae Bonaparte, 1852 | | | |
| <i>Aramus guarauna</i> (Linnaeus, 1766) | | 03°40'51.31" S | 040°20'37.64" W |
| Rallidae Rafinesque, 1815 | | | |
| * <i>Aramides cajaneus</i> (Statius Muller, 1776) | | 03°40'52.98" S | 040°20'39.51" W |
| <i>Laterallus melanophaius</i> (Vieillot, 1819) | wikiaves.com/1636037 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Laterallus exilis</i> (Temminck, 1831) | wikiaves.com/1677983 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Porzana flavigaster</i> (Boddaert, 1783) | wikiaves.com/1696003 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Pardirallus maculatus</i> (Boddaert, 1783) | wikiaves.com/1175917 | 03°40'45.23" S | 040°20'40.70" W |
| <i>Gallinula galeata</i> (Lichtenstein, 1818) | wikiaves.com/1450751 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Porphyrio martinicus</i> (Linnaeus, 1766) | wikiaves.com/1662726 | 03°40'52.98" S | 040°20'39.51" W |
| Charadriiformes Huxley, 1867 | | | |
| Charadriidae Leach, 1820 | | | |
| <i>Vanellus chilensis</i> (Molina, 1782) | wikiaves.com/1175925 | 03°40'52.98" S | 040°20'39.51" W |
| Scolopacidae Rafinesque, 1815 | | | |
| <i>Tringa solitaria</i> Wilson, 1813 | | 03°40'42.24" S | 040°20'36.88" W |
| <i>Tringa flavipes</i> (Gmelin, 1789) | wikiaves.com/1666022 | 03°40'52.98" S | 040°20'39.51" W |
| Jacanidae Chenu & Des Murs, 1854 | | | |
| <i>Jacana jacana</i> (Linnaeus, 1766) | wikiaves.com/1663417 | 03°40'52.98" S | 040°20'39.51" W |
| Columbiformes Latham, 1790 | | | |
| Columbidae Leach, 1820 | | | |
| <i>Columbina talpacoti</i> (Temminck, 1811) | wikiaves.com/1663486 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Columbina picui</i> (Temminck, 1813) | wikiaves.com/1677985 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Columba livia</i> Gmelin, 1789 | wikiaves.com/1677972 | 03°40'52.98" S | 040°20'39.51" W |
| Cuculiformes Wagler, 1830 | | | |
| Cuculidae Leach, 1820 | | | |
| <i>Coccyzus</i> sp. | | 03°40'53.84" S | 040°20'45.93" W |
| <i>Crotophaga ani</i> Linnaeus, 1758 | wikiaves.com/1662722 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Guira guira</i> (Gmelin, 1788) | | 03°40'52.98" S | 040°20'39.51" W |
| Strigiformes Wagler, 1830 | | | |
| Tytonidae Mathews, 1912 | | | |
| * <i>Tyto furcata</i> (Temminck, 1827) | | 03°40'52.98" S | 040°20'39.51" W |
| Caprimulgiformes Ridgway, 1881 | | | |
| Caprimulgidae Vigors, 1825 | | | |
| <i>Hydropsalis parvula</i> (Gould, 1837) | wikiaves.com/1634479 | 03°40'52.98" S | 040°20'39.51" W |
| Apodiformes Peters, 1940 | | | |
| Apodidae Olphe-Galliard, 1887 | | | |
| <i>Tachornis squamata</i> (Cassin, 1853) | wikiaves.com/1666014 | 03°40'52.98" S | 040°20'39.51" W |
| Trochilidae Vigors, 1825 | | | |
| <i>Eupetomena macroura</i> (Gmelin, 1788) | wikiaves.com/1666041 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Chlorostilbon lucidus</i> (Shaw, 1812) | | 03°40'52.98" S | 040°20'39.51" W |
| <i>Polytmus guainumbi</i> (Pallas, 1764) | wikiaves.com/1666023 | 03°40'53.95" S | 040°20'45.82" W |
| <i>Amazilia fimbriata</i> (Gmelin, 1788) | wikiaves.com/1666017 | 03°40'44.44" S | 040°20'38.37" W |
| Coraciiformes Forbes, 1844 | | | |
| Alcedinidae Rafinesque, 1815 | | | |
| <i>Chloroceryle amazona</i> (Latham, 1790) | wikiaves.com/1696010 | 03°40'40.89" S | 40°20'30.15" W |
| Piciformes Meyer & Wolf, 1810 | | | |
| Picidae Leach, 1820 | | | |
| <i>Melanerpes candidus</i> (Otto, 1796) | wikiaves.com/1662736 | 03°40'52.79" S | 040°20'38.20" W |

Continued

Table 1. Continued.

| Order/family/species | Photo or voice vouchers | Latitude | Longitude |
|--|-------------------------|----------------|-----------------|
| Falconiformes Bonaparte, 1831 | | | |
| Falconidae Leach, 1820 | | | |
| <i>Caracara plancus</i> (Miller, 1777) | wikiaves.com/1689391 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Falco femoralis</i> Temminck, 1822 | | 03°40'52.98" S | 040°20'39.51" W |
| Psittaciformes Wagler, 1830 | | | |
| Psittacidae Rafinesque, 1815 | | | |
| <i>Forpus xanthopterygius</i> (Spix, 1824) | wikiaves.com/1662725 | 03°40'52.98" S | 040°20'39.51" W |
| Passeriformes Linnaeus, 1758 | | | |
| Furnariidae Gray, 1840 | | | |
| <i>Furnarius figulus</i> (Lichtenstein, 1823) | wikiaves.com/1689357 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Pseudoseisura cristata</i> (Spix, 1824) | wikiaves.com/1670934 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Certhiaxis cinnamomeus</i> (Gmelin, 1788) | wikiaves.com/1662737 | 03°40'52.98" S | 040°20'39.51" W |
| Tityridae Gray, 1840 | | | |
| <i>Xenopsaris albinucha</i> (Burmeister, 1869) | wikiaves.com/1636101 | 03°40'54.12" S | 040°20'45.22" W |
| Rhynchocyclidae Berlepsch, 1907 | | | |
| <i>Tolmomyias flaviventris</i> (Wied, 1831) | wikiaves.com/1696011 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Todirostrum cinereum</i> (Linnaeus, 1766) | wikiaves.com/1663485 | 03°40'52.98" S | 040°20'39.51" W |
| Tyrannidae Vigors, 1825 | | | |
| <i>Machetornis rixosa</i> (Vieillot, 1819) | wikiaves.com/1666037 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Phaeomyias murina</i> (Spix, 1825) | wikiaves.com/1450750 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Pitangus sulphuratus</i> (Linnaeus, 1766) | wikiaves.com/1677965 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Myiozetetes similis</i> (Spix, 1825) | wikiaves.com/1666036 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tyrannus melancholicus</i> Vieillot, 1819 | wikiaves.com/1696012 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tyrannus savanna</i> Vieillot, 1808 | wikiaves.com/1689392 | 03°40'54.38" S | 040°20'45.22" W |
| <i>Empidonax varius</i> (Vieillot, 1818) | | 03°40'54.32" S | 040°20'45.88" W |
| <i>Fluvicola albiventer</i> (Spix, 1825) | wikiaves.com/1696009 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Fluvicola nengeta</i> (Linnaeus, 1766) | wikiaves.com/1662723 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Arundinicola leucocephala</i> (Linnaeus, 1764) | wikiaves.com/1662735 | 03°40'52.98" S | 040°20'39.51" W |
| Hirundinidae Rafinesque, 1815 | | | |
| <i>Progne chalybea</i> (Gmelin, 1789) | wikiaves.com/1689408 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tachycineta albiventer</i> (Boddaert, 1783) | wikiaves.com/1663416 | 03°40'52.98" S | 040°20'39.51" W |
| Troglodytidae Swainson, 1831 | | | |
| <i>Troglodytes musculus</i> Naumann, 1823 | wikiaves.com/1696004 | 03°40'52.98" S | 040°20'39.51" W |
| Polioptilidae Baird, 1858 | | | |
| <i>Polioptila plumbea</i> (Gmelin, 1788) | wikiaves.com/1661592 | 03°40'42.21" S | 040°20'35.34" W |
| Turdidae Rafinesque, 1815 | | | |
| <i>Turdus rufiventris</i> Vieillot, 1818 | wikiaves.com/1689398 | 03°40'52.98" S | 040°20'39.51" W |
| Mimidae Bonaparte, 1853 | | | |
| <i>Mimus saturninus</i> (Lichtenstein, 1823) | wikiaves.com/1689397 | 03°40'52.98" S | 040°20'39.51" W |
| Parulidae Wetmore Friedmann, Lincoln, Miller, Peters, van Rossem, Van Tyne & Zimmer, 1947 | | | |
| <i>Myiothlypis flaveola</i> Baird, 1865 | wikiaves.com/1634480 | 03°40'53.36" S | 040°20'44.68" W |
| Icteridae Vigors, 1825 | | | |
| <i>Chrysomus ruficapillus</i> (Vieillot, 1819) | wikiaves.com/1677989 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Agelaioides fringillarius</i> (Spix, 1824) | wikiaves.com/1661595 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Molothrus bonariensis</i> (Gmelin, 1789) | wikiaves.com/1677976 | 03°40'52.98" S | 040°20'39.51" W |
| <i>*Sturnella superciliaris</i> (Bonaparte, 1850) | | 03°40'44.93" S | 040°20'40.20" W |
| Thraupidae Cabanis, 1847 | | | |
| <i>Thlypopsis sordida</i> (d'Orbigny & Lafresnaye, 1837) | wikiaves.com/1414514 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tangara sayaca</i> (Linnaeus, 1766) | wikiaves.com/1670955 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tangara palmarum</i> (Wied, 1823) | wikiaves.com/1670933 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Tangara cayana</i> (Linnaeus, 1766) | wikiaves.com/1668652 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Paroaria dominicana</i> (Linnaeus, 1758) | wikiaves.com/1668675 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Sicalis flaveola</i> (Linnaeus, 1766) | wikiaves.com/1689358 | 03°40'52.98" S | 040°20'39.51" W |
| <i>Volatinia jacarina</i> (Linnaeus, 1766) | wikiaves.com/1677977 | 03°40'52.98" S | 040°20'39.51" W |
| Estrildidae Bonaparte, 1850 | | | |
| <i>Estrilda astrild</i> (Linnaeus, 1758) | wikiaves.com/1661593 | 03°40'52.98" S | 040°20'39.51" W |
| Passeridae Rafinesque, 1815 | | | |
| <i>Passer domesticus</i> (Linnaeus, 1758) | wikiaves.com/1670958 | 03°40'52.98" S | 040°20'39.51" W |



Figure 4. Breeding evidence in Parque Ecológico Lagoa da Fazenda, Sobral, Ceará state, northeastern Brazil. Nest of *Certhiaxis cinnamomeus* (**A**) and *Pseudoseisura cristata* (**B**), nest and eggs of *Porzana flavigaster* (**C**), and an adult incubating eggs (**D**). Nests A and B utilized blue plastic stick and pieces of plastic bags, respectively. Photos by A.C. Brasileiro (A and B) and J.A. Neto (C and D).

Table 2. Bird species breeding at Parque Ecológico Lagoa da Fazenda, Ceará state, northeastern Brazil.

| Family/Species | Type of breeding evidence | Month(s) | NBE* |
|----------------------------------|----------------------------|--------------|------|
| Podicipedidae | | | |
| <i>Tachybaptus dominicus</i> | Fledglings | May | 1 |
| Ardeidae | | | |
| <i>Butorides striata</i> | Nest under construction | Feb | 1 |
| Rallidae | | | |
| <i>Porzana flavigaster</i> | Active nest | Jan | 1 |
| <i>Gallinula galeata</i> | Fledglings | Feb–Apr, Jun | 6 |
| <i>Porphyrio martinicus</i> | Fledglings | Mar, Jul | 2 |
| Charadriidae | | | |
| <i>Vanellus chilensis</i> | Active nest | Apr | 1 |
| Jacanidae | | | |
| <i>Jacana jacana</i> | Fledglings | Jan–Apr | 4 |
| Columbidae | | | |
| <i>Columbina talpacoti</i> | Nest under construction | Apr | 1 |
| <i>Columbina picui</i> | Active nest | Sep | 1 |
| Furnariidae | | | |
| <i>Furnarius figulus</i> | Nest under construction | Sep | 2 |
| <i>Pseudoseisura cristata</i> | Nest under construction | Jan | 1 |
| | Adults carrying food items | Mar | 1 |
| <i>Certhiaxis cinnamomeus</i> | Nest under construction | Mar–Apr, Jun | |
| | Adults carrying food items | Dec–Jan | 2 |
| Rhynchocyclidae | | | |
| <i>Todirostrum cinereum</i> | Nest under construction | Mar | 1 |
| Tyrannidae | | | |
| <i>Fluvicola albiventer</i> | Nest under construction | Mar | 2 |
| <i>Arundinicola leucocephala</i> | Nest under construction | Mar | 1 |
| Troglodytidae | | | |
| <i>Troglodytes musculus</i> | Adults carrying food items | Mar | 1 |

* Number of independent observations of breeding evidence

yellowish legs; white ocular ring; grey upperparts with pale spots; head and breast finely streaked dark brown; white breast and belly.

DISCUSSION

Our findings reveal that PELF harbors moderate species richness. Even though some additional species are expected, both richness estimators indicate that our sampling was sufficient to detect most species occurring in the area (Figure 3). The assemblage encompasses mostly aquatic birds, such as those belonging to families Rallidae, Ardeidae, and Scolopacidae. Several species breed in the area and endemic species are also present, but endangered species were absent.

Species richness at PELF can be considered intermediate to other bird inventories from northeastern Brazil, which reported about 130 species for other areas of Caatinga *strictu sensu* (NASCIMENTO 2000; ROOS et al. 2006; FARIA 2007, 2009; FARIA et al. 2010; PEREIRA & AZEVEDO JÚNIOR 2011). However, direct comparisons among studies should be carefully considered due to environmental and sampling differences, such as the size of the sampled area, time span, methods and effort (VIZENTIN-BUGONI et al. 2015). When similar but non-urbanized areas such as other wetlands in Caatinga region are considered, similar richness are reported: 90 species for three vernal pools in Rio Grande do Norte state (PEREIRA 2010) and 125 species in vernal pools and surroundings in Pernambuco

state ('*Lagoas de Petrolina*'; OLMO et al. 2005). Moreover, PELF presents species richness similar to other areas more strongly influenced by urbanization, such as a patch of Atlantic Forest in the city of Aracaju (72 species; MOREIRA & MACHADO 2014) and the Botanical Garden in the city of Recife (90 species; PEREIRA et al. 2011), which suggests that the PELF avifauna is likely reduced due to the effects of the urbanization process.

Several species reported for PELF are known from few previous records in Ceará, such as *Pardirallus maculatus*, *Laterallus melanophaius*, *L. exilis*, *Ixobrychus exilis*, *Botaurus pinnatus*, and *Polytmus guainumbi* (NASCIMENTO 2000; PACHECO 2004; GIRÃO et al. 2007; ALBANO & GIRÃO 2008; GIRÃO et al. 2009; GIRÃO et al. 2013). *Porzana flaviventer* had not yet been documented in inventory birds for Ceará state, so the record extends its distribution in northeastern Brazil (LAMM, 1948; SOUZA 2008; PEREIRA et al. 2014).

The effects of anthropogenic disturbances are apparent in several features of the PELF bird assemblage. For example, the lack of globally endangered species at PELF is likely a consequence of local extinction, which could be indicative of the increasing urban expansion, sewage dumping, and vegetation removal on the avifauna of the park. Anthropogenic influences may also explain the presence of three exotic species which are typically associated with urban areas. Interestingly, the ability of some species to tolerate anthropogenic disturbances reflects also in their behavior; for instance, *Certhiaxis cinnamomeus* and *Pseudoseisura cristata* incorporated artificial material on their nest structures (Figure 4A, B, respectively).

Despite of the depauperate bird assemblage, the PELF harbors several aquatic and endemic species, and also includes *Sicalis flaveola* and *Paroaria dominicana* (endemic), which are widely caught for the illegal cage-bird market in northeastern Brazil (FERNANDES-FERREIRA et al. 2012; PEREIRA & BRITO 2005). Additionally, the PELF harbors long-distance migrant species, such as *Tringa solitaria* and *Tringa flavipes*, which are dependent on wetlands for stopovers along their migratory routes (SILVA et al. 2003). The park is also a breeding site for *Porzana flaviventer*, a widely distributed but rare species, with a poorly understood life history (TAYLOR & DE JUANA 2013; SOUZA 2008). Its eggs were found in a nest on flooded ground, built with the leaves of surrounding vegetation (Figure 4C, D) and similar to an earlier description (VILELLA et al. 2011). Altogether, we suggest that the PELF is regionally important for these birds.

Our findings are relevant to planning management strategies to maintain the local bird diversity. Among the measures that could maintain or even increase the bird species richness at PELF, the most urgent is the cessation of filling of the natural areas and the dumping of sewage in the lakes. Because waste water is usually nutrient-rich, it promotes eutrophication, leading to wetland loss and, therefore, affecting drastically the associated biodiversity (KEDDY 2000). To increase species richness, we recommend the planting of native vegetation, such

as ornithocoric or ornithophilous species, which provide food resources for birds, i.e. fruits and nectar, respectively. Some recommended species are *Ziziphus joazeiro* Mart., *Commiphora leptophloeos* (Mart.) J. B. Gillett, *Tabebuia aurea* (Silva Manso) Benth. & Hook. f. ex S. Moore, and *Erythrina velutina* Willd. that in Portuguese are named *joazeiro*, *imburana*, *ipê-amarelo* and *mulungu*, respectively. In addition, the removal of the exotic *Azadirachta indica* should be considered, as this species is invasive and outcompetes the native flora (MORO et al. 2013, 2014) and indirectly may affect the avifauna.

In summary, we found that the Parque Ecológico Lagoa da Fazenda harbours considerable avifauna encompassing endemic, long-distance migratory, and several aquatic species. Despite the anthropogenic disturbances, which arguably have affected composition of the avifauna, the park is still used as breeding site for several species. Finally, we hope that our recommendations can guide further management plans aimed at maintaining or even increasing the avifauna of this park, which should contribute to bird conservation locally.

ACKNOWLEDGEMENTS

We thank to Universidade Estadual Vale do Acaraú (UVA) for financial support by means of the Programa de Iniciação Científica (PIC-PBU); CAPES for the PhD scholarship to JVB; Diego Santos and Álvaro Nepomuceno for plant identification; Mara Silva for access to the official decree for implantation of the PELF; Wellington Galvão for providing the park shapefile; Marcelo Moro and Weber Girão for access to literature; Daniel Passos for help with statistics; and to the two anonymous reviewers for their contributions to the manuscript.

LITERATURE CITED

- AGUIAR JÚNIOR, P.R. 2005. A cidade e o rio: produção do espaço urbano em Sobral-Ceará. Mercator—Revista de Geografia da UFC 4(8).
- ALBANO, C.G. & W. GIRÃO. 2008. Aves das matas úmidas das Serras de Aratanga, Baturité e Maranguape, Ceará. Revista Brasileira de Ornitologia 16(2): 142–154. <http://www4.museu-goeldi.br/revistabronito/revista/index.php/BJO/article/view/0532>
- BIRD LIFE INTERNATIONAL. 2015. IUCN Red List for birds. Accessed at <http://www.birdlife.org>, 23 April 2015.
- BRANCO, J.O. 2007. Avifauna aquática do Saco da Fazenda (Itajaí, Santa Catarina, Brasil): uma década de monitoramento. Revista Brasileira de Zoológia 24(4):873–882. doi: [10.1590/S0101-81752007000400003](https://doi.org/10.1590/S0101-81752007000400003)
- CBRO (COMITÉ BRASILEIRO DE REGISTROS ORNITOLÓGICOS). 2014.11ª edição. 1 Jan 2014. Lista de Aves do Brasil. Florianópolis. Accessed at <http://www.cbro.org.br/CBRO/listabr.html>, 20 January 2014.
- CEARÁ (ESTADO). Decreto nº 21.303 de 11 de março de 1991. Palácio do Governo do estado do Ceará, Ceará, 1991.
- CORBO, M.A.M., G.B. D'ANGELO, C.H. ALMEIDA & I. SAZIMA. 2013. Aves do Campus da Unicamp e arredores. Vinhedo: Avis Brasilis. 320 pp.
- COLWELL, R.K. 2011. Estimates: statistical estimation of species richness and shared species from samples. Version 9. User's guide

- and application. Accessed at <http://purl.oclc.org/estimates>, 5 April 2016.
- FARIAS, G.B. 2007. Avifauna em quatro áreas de caatinga *strictu sensu* no centro-oeste de Pernambuco, Brasil. Revista Brasileira de Ornitologia 15(1): 53–60. http://www4.museu-goeldi.br/revista_brornito/revista/index.php/BJO/article/viewFile/2803/pdf_447
- FARIAS, G.B. 2009. Aves do Parque Nacional de Pernambuco, Brasil do Catimbau, Buíque, Pernambuco, Brasil. Atualidades Ornitológicas On-line 147: 36–39. http://www.ao.com.br/download/ao147_36.pdf
- FARIAS, G.B., W.A.G. SILVA & C.G. ALBANO. 2005. Diversidade de aves em áreas prioritárias para conservação da Caatinga; pp. 203–226, in: F.S. ARAÚJO, M.J.N. RODAL & M.R.V. BARBOSA (eds.). Análise das variações da biodiversidade do bioma Caatinga: Suporte a estratégias regionais de conservação. Brasília: Ministério do Meio Ambiente.
- FARIAS, G.B., G.A. PEREIRA & K.Q. BURGOS. 2010. Aves da Floresta Nacional de Negreiros (Serrita, Pernambuco). Atualidades Ornitológicas On-line 157: 41–46. http://www.ao.com.br/download/AO157_41.pdf
- FERNANDES-FERREIRA, H., S.V. MENDONÇA, C. ALBANO, F.S. FERREIRA & R.R.N. ALVES. 2012. Hunting, use and conservation of birds in northeast Brazil. Biodiversity and Conservation 21: 221–244. doi: [10.1007/s10531-011-0179-9](https://doi.org/10.1007/s10531-011-0179-9)
- GIRÃO W., C. ALBANO & H.T. ABREU. 2009. Registros do socoí-vermelho no estado do Ceará, Brasil *Ixobrychus exilis* (Gmelin, 1789). Atualidades Ornitológicas 149: 14–15.
- GIRÃO, W., A. NETTO, R. LEITE, T.T. SILVA, T.T. & A.C. SAMPAIO. 2013. Primeiros registros documentados do socó-boi-baio *Botaurus pinnatus* no Ceará. Atualidades Ornitológicas 173: 16–17.
- GIRÃO, W., C. ALBANO, T. PINTO & L.F. SILVEIRA. 2007. Avifauna da Serra de Baturité: dos naturalistas à atualidade; pp. 187–224, in: T.S. OLIVEIRA & E.F.S. ARAÚJO (eds.). Biodiversidade e conservação da biota na serra de Baturité, Ceará. Fortaleza: Edições UFC, Coelce.
- Gwynne, J.A., R.S. RIDGELY, G. TUDOR & M. ARGEL. 2003. Aves do Brasil: Pantanal e Cerrado. São Paulo: Editora Horizonte. 322 pp.
- IBGE (INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA). 2004. Mapa de biomas do Brasil: primeira aproximação. Rio de Janeiro. Accessed at <http://www.ibge.gov.br/>, 20 July 2014.
- IEPECE (INSTITUTO DE PESQUISA E ESTRATÉGIA ECONÔMICA DO CEARÁ). 2013. Perfil básico municipal 2013: Sobral. Accessed at http://www.ipece.ce.gov.br/publicacoes/perfil_basico/pbm-2013/Sobral.pdf, 10 June 2014.
- KEDDY, P. 2000. Wetland ecology: principles and conservation. Cambridge: Cambridge University Press. 614 pp.
- LAMM, D. W. 1948. Notes on the birds of the states of Pernambuco and Paraíba, Brazil. Auk 65: 261–283. <https://sora.unm.edu/sites/default/files/journals/auk/v065n02/p0261-p0283.pdf>
- LIMA, V.B. & L.F. ASSIS. 2005. Mapeando alguns roteiros de trabalho de campo em Sobral (CE): uma contribuição ao ensino de geografia. Revista da Casa da Geografia 6/7(1): 109–121. <http://www.uvanet.br/rcgs/index.php/RCGS/article/view/125>
- LOBADA, C.R., & B.L.D. DE ANGELIS. 2005. Áreas verdes públicas urbanas: conceitos, usos e funções. Revista Ambiência 1(1): 125–139. doi: [10.1007/s10531-011-0179-9](https://doi.org/10.1007/s10531-011-0179-9)
- MACIEL, B.A. 2010. Unidades de conservação no Bioma Caatinga; pp. 76–81, in: M.A. GARIGLIO, E.V. S.B. SAMPAIO, L.A. CESTARO & P.Y. KAGEYAMA (eds.). Uso Sustentável e Conservação dos Recursos Florestais da Caatinga. Brasília: Serviço Florestal Brasileiro.
- MAGURRAN, A.E. 2004. Measuring biological diversity. Oxford: Blackwell Publishing. 256 pp.
- MARINI, M.A. & F.I. GARCIA. 2005. Conservação de Aves no Brasil. Mega diversidade, Belo Horizonte 1(1): 95–102. <https://simonprojetos.files.wordpress.com/2012/12/marini-e-garcia-2005-conservacao-de-aves-no-brasil.pdf>
- MATSUMOTO, M.L., Z.F. PEREIRA, J.P. GONÇALVES, A. SANGALLI & S.S.F. FERNANDES. 2012. Avaliação ambiental do parque urbano Arnulpho Fioravante para adoção de estratégias de restauração. Boletim Paranaense de Geociências 66-67: 51–60. <http://ojs.c3sl.ufpr.br/ojs/index.php/geociencias/article/viewFile/19400/20449>
- MMA (MINISTÉRIO DO MEIO AMBIENTE). 2002. Biodiversidade Brasileira — Avaliação e identificação de áreas e ações prioritárias para conservação, utilização sustentável e repartição dos benefícios da biodiversidade nos biomas brasileiros. Brasília. Série Biodiversidade. 404 pp.
- MOREIRA, A.L.O. & C.G. MACHADO. 2014. Descaracterização da avifauna no último remanescente de Mata Atlântica de Aracaju, capital de Sergipe, Brasil. Atualidades Ornitológicas 178: 33–38. http://www.ao.com.br/download/AO178_33.pdf
- MORO, M.F., C. WESTERKAMP & F.S. ARAÚJO. 2014. How much importance is given to native plants in cities' treescape? A case study in Fortaleza, Brazil. Urban Forestry and Urban Greening 13(2): 365–374. doi: [10.1016/j.ufug.2014.01.005](https://doi.org/10.1016/j.ufug.2014.01.005)
- MORO, M.F.C. WESTERKAMP & F.R. MARTINS. 2013. Naturalization and potential impact of the exotic tree *Azadirachta indica* A.Juss. in northeastern Brazil. Check List 9(1): 53–156. doi: [10.15560/9.1.153](https://doi.org/10.15560/9.1.153)
- NASCIMENTO, J.L.X. 2000. Estudo comparativo da avifauna em duas Estações Ecológicas da caatinga: Aiuba e Seridó. *Melopsittacus* 3(1): 12–35.
- NASCIMENTO, J.L.X., I.L.S. NASCIMENTO & S.M. AZEVEDO JÚNIOR. 2000. Aves da Chapada do Araripe (Brasil): biologia e conservação. Ararajuba 8(2): 115–125. http://www4.museu-goeldi.br/revista_brornito/revista/index.php/BJO/article/viewFile/1307/pdf_407
- NASCIMENTO, J.L.X., L.Z. SALES JÚNIOR, A.E.B.A. SOUSA, & J. MINNS. 2005. Avaliação rápida das potencialidades ecológicas e econômicas do Parque Nacional de Ubajara, Ceará, usando aves como indicadores. Ornithologia 1(1): 33–42. http://observatorio.wwf.org.br/site_media/upload/gestao/documentos/Nascimento_2005.pdf
- NIMER, E. 1972. Climatologia da região Nordeste do Brasil. Introdução à climatologia dinâmica. Revista Brasileira de Geografia 34(2): 3–51.
- OLMOS, F., W.A.G. SILVA & C.G. ALBANO. 2005. Aves em oito áreas de caatinga no Sul do Ceará e Oeste de Pernambuco, Nordeste do Brasil: composição, riqueza e similaridade. Papéis Avulsos de Zoologia. São Paulo 45(14): 179–199. doi: [10.1590/S0031-10492005001400001](https://doi.org/10.1590/S0031-10492005001400001)
- PACHECO, J.F. 2004. As aves da Caatinga: uma análise histórica do conhecimento; pp.190–250, in: J.M.C. SILVA, M. TABARELLI, M.T. FONSECA & L.V. LINS (eds.). Biodiversidade da caatinga: áreas e ações prioritárias para conservação. Brasília: Ministério do Meio Ambiente.
- PEREIRA, G.A. Avifauna associada a três lagoas temporárias no estado do Rio Grande do Norte, Brasil. 2010. Atualidades Ornitológicas 156: 53 – 60. http://www.ao.com.br/download/AO156_53.pdf
- PEREIRA, G.A., P.P. RODRIGUES, S. LEAL, M.C. PERIQUITO, G.B.M. DA SILVA et al. 2014. Important bird records from north-east Brazil. Cotinga 36: 45 – 50. <http://wordpress.neotropicalbirdclub.org/wp-content/uploads/2015/03/C36-Pereira2.pdf>
- PEREIRA, G.A., M.C. PERIQUITO, M.T. BRITO & M. MENEZES. 2011. Estrutura trófica da avifauna no Jardim Botânico do Recife, Pernambuco, Brasil. Atualidades Ornitológicas 164: 57–63. http://www.ao.com.br/download/AO164_57.pdf
- PEREIRA, G.A. & S.M. AZEVEDO JÚNIOR. 2011. Estudo comparativo entre as comunidades de aves de dois fragmentos florestais de Caatinga em Pernambuco, Brasil. Revista Brasileira de Ornitologia 19(1): 22–31. http://www4.museu-goeldi.br/revista_brornito/revista/index.php/BJO/article/view/4203
- PEREIRA, G.A. & M.T. BRITO. 2005. Diversidade de aves silvestres brasileiras comercializadas nas feiras livres da Região Metropolitana

- do Recife, Pernambuco. Atualidades Ornitológicas (126):14–19. <http://www.ao.com.br/download/glauco.pdf>
- PRADO, D.E. 2003. As Caatingas da América do Sul; pp. 3–74; in: I.R. LEAL, M. TABARELLI & J.M.C. SILVA (eds.). Ecologia e conservação da Caatinga. Recife: Editora Universitária da UFPE.
- ROOS, A.L., M.F.C. NUNES, E.A. SOUSA, A.E.B.A. SOUSA, J.L.X. NASCIMENTO & R.C.A. LACERDA. 2006. Avifauna da região do Lago de Sobradinho: composição, riqueza e biologia. Ornithologia 1(2): 135–160. <http://cemave.net/publicacoes/index.php/ornithologia/article/view/16>
- SIGRIST, T. 2013. Guia de campo avis brasiliis: avifauna brasileira. São Paulo: Avis Brasilis. 592 pp.
- SILVA, J.M.C., M.A. SOUZA, A.G.D. BIEBER & C.J. CARLOS. 2003. Aves da Caatinga: status, uso do habitat e sensitividade; pp. 237–273, in: I.R. LEAL, M. TABARELLI & J.M.C. SILVA (eds.). Ecologia e conservação da Caatinga. Recife: Editora Universitária da UFPE.
- SOUZA, D. 2008. Primeiro registro documentado da Sanã-amarela *Porzana flavigaster* (Boddaert 1783), Aves, Rallidae, para a Bahia e revisão da distribuição brasileira da espécie. Atualidades Ornitológicas 144: 4–6.
- TAYLOR, B. & E. DE JUANA. 2013. Yellow-breasted Crake (*Hapalocrex flavigaster*). In: J. DEL HOYO, A. ELLIOTT, J. SARGATAL, D.A. CHRISTIE & E. DE JUANA (eds.). Handbook of the birds of the world alive. Lynx Edicions: Barcelona. Accessed at <http://www.hbw.com/node/53668>, 15 July 2015.
- VILELLA, F.J., J.A. CRUZ & M. LÓPEZ. 2011. Nesting ecology of the Yellow-Breasted Crake (*Porzana flavigaster*) in Puerto Rico. Waterbirds 34(3): 366–368. doi: [10.1675/063.034.0311](https://doi.org/10.1675/063.034.0311)
- VIZENTIN-BUGONI, J., F.P. JACOBS, M.A.A. COIMBRA & R.A. DIAS. 2015. Birds of the Reserva Biológica do Mato Grande and surroundings, Rio Grande do Sul, Brazil. Check List 11(3): 1641. doi: [10.15560/11.3.1641](https://doi.org/10.15560/11.3.1641)

Authors' contributions: ACB collected and analyzed the data, and wrote the text. FLA collected the data and made maps; JAA, AMP and JAN collected the data, ROT and ASS coordinated the research and JVB analyzed the data, wrote the text and guided the research.

Received: 27 April 2016

Accepted: 27 December 2016

Academic editor: Nárgila Gomes Moura