



First record of the Sungrebe, *Heliornis fulica* (Boddaert, 1783) (Gruiformes, Heliornithidae), from Piauí, northeastern Brazil

Leonardo Moura dos Santos Soares,¹ André Bastos da Silva,² Anderson Guzzi²

1 Universidade Federal do Piauí, Campus Professora Cinobelina Elvas, Departamento de Biologia, Br 135, km 3, Planalto Horizonte, CEP 64900-000, Bom Jesus, PI, Brazil. **2** Universidade Federal do Piauí, Campus Ministro Petrônio Portella, Programa de Pós-Graduação em Desenvolvimento e Meio Ambiente, Núcleo de Referências em Ciências Ambientais do Trópico Ecotonal do Nordeste, Av. Universitária 1310, Ininga, CEP 64049-550, Teresina, PI, Brazil.

Corresponding author: Anderson Guzzi, guzzi@ufpi.edu.br

Abstract

This study presents the first record of the Sungrebe *Heliornis fulica* for the state of Piauí in northeastern Brazil, a first sighting also in a Cerrado/Caatinga ecotone. On 19 December 2014, a female Sungrebe was collected by a local fisherman in the Corrente River located between the towns of Capitão de Campos and Piripiri in the northcentral part of the state of Piauí.

Key words

Ecotone; Parnaíba River watershed; Corrente River; aquatic bird; geographic distribution; conservation.

Academic editor: Johan Ingels | Received 31 May 2017 | Accepted 8 July 2017 | Published 11 September 2017

Citation: Soares LMS, Silva AB, Guzzi A (2017) First record of the Sungrebe, *Heliornis fulica* (Boddaert, 1783) (Gruiformes, Heliornithidae), from Piauí, northeastern Brazil. Check List 13 (5): 411–415. <https://doi.org/10.15560/13.5.411>

Introduction

The family Heliornithidae includes only 1 species, *Heliornis fulica* (Boddaert, 1783), which is widely distributed throughout the Neotropical region but apparently with low population densities (Bertram and Kirwan 2017). Its preferred habitats are quiet forest streams and rivers, and freshwater ponds and lakes, with overhanging riparian vegetation. These aquatic birds tend to swim near the vegetation hanging from the river banks, taking refuge under it at the first sign of danger. They feed on aquatic insects and their larvae, as well as beetles, ants, spiders, crustaceans, amphibians, and small fish. Sungrebes are difficult to observe in their natural habitat, due to their inconspicuous color pattern and secretive behavior (Ingels and Tauleigne 2011). Sometimes when disturbed from their roosts at night, they are caught in fishing nets (Sick 1997).

The state of Piauí has an extensive ecotone zone at the intersection of the Amazon Forest, Cerrado (Neotropical savanna), and semiarid Caatinga (deciduous, thorny, dryland vegetation), and it represents the largest phytoecological domain in the Parnaíba River watershed (Sousa et al. 2009). Due to the climatic complexity of the region and its habitat heterogeneity, this ecotone zone harbors many bird species (Gonçalves et al. 2017) and has been surveyed since the beginning of the 20th century (Hellmayr 1929, Pacheco 2004). Surveys are now more intense within conservation areas (Novaes 1992, Olmos 1993, Zaher 2001a, 2001b, Santos 2001, Olmos and Albano 2012, Santos et al. 2012, Silveira and Santos 2012, Guzzi 2012), as well as in less-protected sites (Santos 2004, Olmos and Brito 2007).

Although numerous studies in recent years have focused on the birds of Piauí, there are no records of




Figure 1. Map showing new record of *Heliornis fulica* from a section of the Corrente River located between the towns of Capitão de Campos and Piripiri, north-central Piauí in a Cerrado/Caatinga ecotone and other records from northeastern Brazil.



Figures 2, 3. The female specimen of Sungrebe *Heliornis fulica* collected by a local fisherman in the Corrente River located between the towns of Capitão de Campos and Piripiri in the northcentral part of the state of Piauí, Brazil. **2.** Side view. **3.** Dorsal view. Photographs by A.B. Silva.

the occurrence of *H. fulica*. As such, the present study reports the first record of the Sungrebe for that state, a first sighting also in an ecotone area (Cerrado/Caatinga). Additionally, this record extends the known range of that species towards the interior of northeastern Brazil.

Methods

The region where the specimen of *H. fulica* was collected is situated in an ecotone zone (Rivas 1996) at the confluence of Cerrado and dryland Caatinga vegetation (Jacomine et al. 1986) (Fig. 1). The climate is hot and semiarid, presenting 6 months of drought and has an average annual rainfall of 800–1000 mm, with the heaviest rains occurring between January and March (CODEVASF 2006). The average annual minimum and maximum temperatures of are 22 °C and 33 °C respectively (Lima 2002).

Results

On 19 December 2014, a Sungrebe *H. fulica* was collected by a local fisherman in a section of Corrente River located between the towns of Capitão de Campos and Piripiri in northern-central Piauí, Brazil. According to the

fisherman, the Sungrebe was captured about 200 m downstream of the Oiteiro dam (04°28'37" S, 041°50'11" W) (Fig. 4), in the Longá watershed. The specimen was brought to the Center for the Reception of Wild Animals (CETAS) of the Brazilian Institute of Environment and Renewable Resources (IBAMA) of the state of Piauí, for release back into nature later.

The characters that aided in the identification of this Sungrebe, a female, were: head and neck strikingly



Figure 4. Corrente River between the towns of Capitão de Campos and Piripiri in north-central Piauí, Brazil, where the female specimen of *Heliornis fulica* was collected about 200 m from the Oiteiro dam by a local fisherman. Photograph by A.B. Silva.

marked black and white, cinnamon cheeks typical for a female, red eye-ring, and reddish maxilla (Mata et al. 2006) (Figs 2, 3).

Discussion

There are few records of *H. fulica* for northeastern Brazil (e.g., Oren and Roma 2011). One specimen deposited in the Paraense Emílio Goeldi Museum (voucher MPEG 43389) was collected in Balsas, state of Maranhão, Brazil. Unofficial records (not found in the scientific literature) of *H. fulica* can be found on WikiAves (2017): the town of Passo de Camaragibe (Alagoas state); the towns of Ilhéus and São Desidério (Bahia state); and the towns of Riachão and Penalva (Maranhão state) (Fig. 1). The absence of records of *H. fulica* in other parts of the state of Piauí and northeastern Brazil may be due to a lack of detailed observations in areas with appropriate habitat.

Although *H. fulica* has high habitat specificity (Stotz et al. 1996), its conservation status was categorized as Least Concern due to its extremely wide range and large numbers, as well as populations trends that do not appear to be rapidly decreasing (BirdLife International 2016). However, defaunation caused by habitat destruction (Hoffmann et al. 2010, Dirzo et al. 2014) has motivated actions directed toward its conservation. Some populations of *H. fulica* in Brazil's Atlantic Forest and Amazon regions are considered Vulnerable (Silva et al. 2008, Pacheco et al. 2010, Oren and Roma 2011) or Critically Endangered (CONSEMA 2011) and therefore are conservation priorities.

The region where the new record of *H. fulica* was found is in an advanced state of environmental degradation, principally due to ecosystem fragmentation resulting from farming and cattle raising. Additional anthropogenic threats to forest remnants are burning and bushmeat hunting. Thus, actions to protect towards those natural habitats are extremely important to the continued presence of this species in Piauí.

This first record of *H. fulica* from Piauí helps to fill a gap in the geographic distribution of this species in northeastern Brazil and suggests that the distribution of *H. fulica* is considerably wider than believed. The new data increases is a useful contribution to the conservation of this aquatic bird species.

Acknowledgements

We thank Ferdinand Costa and Maria de Lourdes Silva Costa for their help during fieldwork, Fabiano Barbosa Pessoa for taking care of the specimen at CETAS (IBAMA), Karoline Veloso Ribeiro for preparing the map, and Nathalie Reis Monteiro for helping with the brief review.

Authors' Contributions

LMSS and ABS collected the data, LMSS identified the specimen; LMSS, ABS and AG wrote the text.

References

- Bertram BCR, Kirwan GM (2017) Sungrebe (*Heliornis fulica*). In: del Hoyo J, Elliott A, Sargatal J, Christie DA, Juana E (Eds) Handbook of the Birds of the World Alive. Lynx Edicions, Barcelona. <http://www.hbw.com/node/53707>. Accessed on: 2017-4-24.
- BirdLife International (2016) *Heliornis fulica*. The IUCN Red List of Threatened Species 2016. <https://dx.doi.org/10.2305/IUCN.UK.2016-3.RLTS.T22692192A93340756.en>
- CODEVASF (2006) Plano de Ação para o Desenvolvimento Integrado da Bacia do Parnaíba, PLANAP: Síntese Executiva: Território dos Carnaubais. TDA Desenhos & Arte Ltda, Brasília, 76 pp. <http://files.assuncaoliveira.com/200000890-a0ca6a2c2e/SINTESE%20-%20TERRIT%C3%93RIO%20DOS%20CARNAUBAIS.pdf>. Accessed on: 2017-4-15. [Technical report].
- CONSEMA (2011) Resolução CONSEMA Nº 002. http://www.fatma.sc.gov.br/upload/Fauna/resolucao_fauna__002_11_fauna.pdf. Accessed on: 2017-4-15.
- Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJB, Collen B (2014) Defaunation in the Anthropocene. *Science* 345 (401): 401–406. <https://doi.org/10.1126/science.1251817>
- Gonçalves GR, Santos MPD, Cerqueira PV, Juen L, Bispo AA (2017) The relationship between bird distribution patterns and environmental factors in an ecotone area of northeast Brazil. *Journal of Arid Environments* 140: 6–13. <https://dx.doi.org/10.1016/j.jaridenv.2017.01.004>
- Guzzi A, Tavares AA, Santos AGS, Cardoso CO (2012) Aves do Delta do Parnaíba. In: Guzzi A (Ed.). Biodiversidade da APA Delta do Parnaíba, porção piauiense, 1ª ed. EDUFPI, Teresina, 291–338.
- Hellmayr CE (1929) A contribution to the ornithology of northeastern Brazil. *Fieldiana, Zoological Series* 12 (18): 1–284.
- Hoffmann M, Hilton-Taylor C, Angulo A, Böhm M, Brooks TM, et al. (2010) The impact of conservation on the status of the world's vertebrates. *Science* 330: 1503–1509.
- Ingels J, Tauleigne D (2011) First record of a Sungrebe (*Heliornis fulica*) nest in French Guiana. *Boletín SAO* 20 (2): 52–55.
- Jacomine PKT, Cavalcanti AC, Pessoa SCP, Burgos N, Mélo Filho HFR, Lopes OF, Medeiros LAR (1986) Levantamento Exploratório de Solos do Estado do Piauí. EMBRAPA-SNLCs, Boletim de pesquisa, Série recursos de solos 36 (18): 1–782.
- Lima MG (2002) Estimativa da temperatura do ar no Piauí. UFPI, Teresina, 48 pp.
- Mata JRR, Erize F, Rumboll M (2006) Collins Field Guide: Birds of South America. Princeton University Press, Princeton/Oxford, 384 pp.
- Novaes FC (1992) Bird observations in the state of Piauí, Brazil. *Goeldiana Zoologia* 16–18: 1–5.
- Olmos F (1993) Birds of Serra da Capivara National Park, in the “caatinga” of north-eastern Brazil. *Bird Conservation International* 3 (1): 21–36. <https://doi.org/10.1017/S0959270900000769>
- Olmos F, Albano CG (2012) As aves da região do Parque Nacional Serra da Capivara (Piauí, Brasil). *Revista Brasileira de Ornitologia* 20 (3): 173–187.
- Olmos F, Brito GRR (2007) Aves da região da Barragem de Boa Esperança, médio rio Parnaíba, Brasil. *Revista Brasileira de Ornitologia* 15 (3): 37–52.
- Oren DC, Roma JC (2011) Composição e vulnerabilidade da avifauna da Amazônia maranhense, Brasil. In: Martins MB, Oliveira TG (Eds) Amazônia Maranhense: Diversidade e Conservação. MPEG, Belém, 221–250.
- Pacheco JF (2004) As aves da Caatinga: uma análise histórica do conhecimento. In: Silva JMC, Tabarelli M, Fonseca MT, Lins LV (Eds) Biodiversidade da Caatinga: Áreas e Ações Prioritárias para Conservação. MMA, Brasília, 189–250.
- Pacheco JF, Astor INC, Cesar CB (2010) Avifauna da Reserva Biológica de Poço das Antas, Silva Jardim, RJ. *Atualidades Ornitológicas* 157: 55–74
- Rivas MP (1996) Macrozoneamento Geoambiental da Bacia Hidrográfica do Rio Parnaíba. IBGE, Rio de Janeiro, 111 pp.

- Santos MPD (2001) Composição da avifauna nas Áreas de Proteção Ambiental Serra da Tabatinga e Chapada das Mangabeiras, Brasil. Boletim do Museu Paraense Emílio Goeldi, Série Zoologia 17 (1): 43–67.
- Santos MPD (2004) As comunidades de aves em duas fisionomias da vegetação de Caatinga no estado do Piauí, Brasil. Revista Brasileira de Ornitologia 12 (2): 113–123.
- Santos MPD, Santana A, Soares LMS, Sousa SA (2012) Avifauna of Serra Vermelha, south of Piauí, Brazil. Revista Brasileira de Ornitologia 20 (3): 199–214. http://www4.museu-goeldi.br/revistabrornito/revista/index.php/BJO/article/download/4904/pdf_632
- Sick H (1997) Ornitologia brasileira. Editora Nova Fronteira, Rio de Janeiro, 912 pp.
- Silva WR, Silveira LF, Uezu A, Antunes AZ, Sugieda AM, Hasui E, Figueiredo LF, Develey PF (2008) Aves. In: Rodrigues RR, Bononi VLR (Eds) Diretrizes para a Conservação e Restauração da Biodiversidade no Estado de São Paulo. Instituto de Botânica, São Paulo, 77–171.
- Silveira LF, Santos MPD (2012) Bird richness in Serra das Confusões National Park, Brazil: How many species may be found in an undisturbed caatinga? Revista Brasileira de Ornitologia 20 (3): 188–198.
- Sousa SR, Castro AAJF, Farias RRS, Sousa GM, Castro NMCF (2009) Fitoecologia do complexo de Campo Maior, Piauí, Brasil. Publicações Avulsas em Conservação de Ecossistemas 22: 1–25. <https://dx.doi.org/10.18029/1809-0109/pacc.n22p1-22>
- Stotz DF, Fitzpatrick JW, Parker TA, Moskovitz DK (1996) Neotropical Birds: Ecology and Conservation. University of Chicago Press, Chicago, 502 pp.
- WikiAves (2017) Mapa de Registros da Espécie Picaparra (*Heliornis fulica*). http://www.wikiaves.com/mapaRegistros_picaparra. Accessed on: 2017-4-15.
- Zaher HED (2001a) Projeto Diversidade de Vertebrados Terrestres da Estação Ecológica de Uruçuí-Una, Piauí (PI): Subsídios para o Plano de Manejo. Fundação O Boticário, Curitiba, 110 pp. [Technical report].
- Zaher HED (2001b) Relatório Sobre o Levantamento Preliminar da Fauna de Vertebrados Terrestres do Parque Nacional da Serra das Confusões, Piauí. IBAMA, Teresina, 41 pp. [Unpublished report].