



First records of water scavenger beetle species (Coleoptera, Hydrophilidae) from Quindío Department, Colombia

Liza M. González-Rodríguez,¹ Andrea L. García-Hernández,¹ Bruno Clarkson²

1 Universidad del Quindío, Centro de Estudio e Investigaciones en Biodiversidad y Biotecnología de la Universidad del Quindío, Grupo de Estudios en Artrópodos del Quindío, Carrera 15 Calle 12 Norte, Armenia, Quindío, Colombia. **2** Departamento de Entomología, Museu Nacional, Universidade Federal do Rio de Janeiro, Quinta da Boa Vista, São Cristóvão, CEP 20940-040, Rio de Janeiro, RJ, Brazil.

Corresponding author: Liza M. González-Rodríguez, lizmgr287@gmail.com

Abstract

A list of species of Hydrophilidae from Quindío Department, Colombia is presented. This list is based on recently collected specimens, a survey of data from the literature, and specimens deposited in the Colección de Insectos de la Universidad del Quindío (CIUQ). Twenty-three species, including 16 new records from Colombia, are listed. Information about biology of each species and a checklist of the overall Colombian species are given.

Key words

Aquatic beetles; checklist; new records; range extension; Neotropics; South America.

Academic editor: Juan Pablo Botero | Received 8 October 2016 | Accepted 2 August 2017 | Published 13 October 2017

Citation: González-Rodríguez LM, García-Hernández AL, Clarkson B (2017). First records of water scavenger beetle species (Coleoptera, Hydrophilidae) from Quindío Department, Colombia. Check List 13 (5): 605–620. <https://doi.org/10.15560/13.5.605>

Introduction

Currently, there are few taxonomic and ecologic published studies about Colombian aquatic beetles. Studies on specific groups or families are even fewer. Data on biodiversity of the Hydrophilidae of Colombia is extremely limited, as well as information about habitat preference, feeding habits, life cycle, and geographic distribution of species (Arias-Díaz et al. 2007). According to the major species catalogues (Hansen 1999, Short and Hebauer 2006, Short and Fikáček 2011), there are 34 species recorded from Colombia, including representatives of the subfamilies Acidocerinae (2 spp.), Enochrinae (2 spp.), Hydrophilinae (20 spp.), and Sphaeridiinae (10 spp.). The present list contributes to the knowledge of diversity and biology of Colombian hydrophilid beetles and adds new localities in Quindío Department.

Methods

Specimens were examined, identified, measured and illustrated using a Stemi 2000-C ZEISS microscope with attached AxiomCam ERc 5s camera. Taxonomic identification of the specimens was done with the support of the keys and information by Leech (1943), Mouchamps (1963), Young (1967), Bachmann (1969), Gundersen (1977), Smetana (1978), Oliva (1981, 1987, 1989, 1994), Fernández (1988, 1989, 1994, 1997), Oliva et al. (2002), Short (2004), Komarek and Beutel (2007), Clarkson and Ferreira-Jr (2009), Arce-Pérez and Morón (2013). Morphological terminology follows mainly Hansen (1991), with some terminology adapted from Komarek (2004).

The geographic distribution of each species was taken from the publications already cited and from major species catalogues (Zaitzev 1908, Blackwelder 1944, Hansen 1999, Short and Hebauer 2006, Short and

Fikáček 2011), as well as from Bedel (1891), Régimbart (1901), Orchymont (1921a, 1921b, 1922), Mouchamps (1959), Wooldridge (1973), Smetana (1975, 1978, 1984), Gundersen (1977), Fernández (1982, 1988, 1989, 1997), Oliva (1983, 1994, 2010, 2014), Short (2004, 2005), Komarek (2005), Fikáček and Short (2006), Peck (2006), Short and Torres (2006), García-Hernández (2009), Oliva and Short (2012), Arce-Pérez and Morón (2013), and Deler-Hernández et al. (2013). All the locality data was provided as detailed as available in the above papers. Records of species available in old catalogues (e.g., Zaitzev 1908, Blackwelder 1944) were confirmed with the examination of the literature above cited. Information on species synonymy was obtained from Fernández (1982, 1988, 1989), Oliva (1989), Hansen (1991, 1999), Komarek (2005), Oliva and Short (2012), and Fikáček and Vondráček (2014). The species are presented in alphabetical order but keeping the current classification of the Hydrophilidae as proposed by Short and Fikáček (2013). Lists of hydrophilid species from Colombia and the complete distribution of species from Quindío

Department are shown in Tables 1 and 2, respectively, and a distributional map is presented (Figure 48).

Results

Subfamily Acidocerinae

Helochaeres abbreviatus (Fabricius, 1801) (Figs 1, 24)

Hydrophilus abbreviatus Fabricius 1801: 251. *Helochaeres* (s. str.) *abbreviatus* (Fabricius)—Orchymont 1936: 10.

Philydrus pallidus Castelnau 1840: 53, secondary homonym of

Hydrophilus pallidus Rossi, 1972. Syn.: Orchymont 1936: 10

Philydrus pallidus Castelnau—Gemminger and Harold 1868: 482.

Helochaeres pallidus (Castelnau)—Fleutiaux and Sallé 1889: 376.

Enochrus (*Lumetus*) *pallidus* (Castelnau)—Zaitzev 1908: 388.

Helochaeres (*Helochaeres*) *pallidus*—Orchymont 1926: 232–233.

Helochaeres (*Hydrobaticus*) *rufobrunneus* Balfour-Browne 1939: 293.

Syn.: Spangler 1981: 158, Fernández 1982: 34.

Helochaeres (*Helochaeres*) *abbreviatus* Orchymont 1943: 55.

Diagnosis. Body 6.60–6.70 mm long. Dorsal punctation on head, pronotum and elytra evenly fine and sparse. Mesoventral process slightly elevated. Last abdominal ventrite deeply notched. Parameres longer than median

Table 1. List of species of Hydrophilidae from Colombia. * Without locality specification. ** Cited from Colombia by Blackwelder (1944), but not by Zaitzev (1908), Hansen (1999), Short and Hebauer (2006), and Short and Fikáček (2011). *** Recently recorded without nominal identification (Short et al. 2017).

Species	Geographic distribution
Subfamily Acidocerinae	
<i>Globulosis</i> sp.	Colombia [Meta]***
<i>Helochaeres abbreviatus</i> (Fabricius, 1801)	Argentina, Bolivia, Brazil, Colombia* [Quindío], Cuba, French Guiana, Granada, Panama, Paraguay, Suriname, Venezuela
<i>Helochaeres atratus</i> Bruch, 1915	Argentina, Brazil, Colombia [Quindío], Ecuador (?), Paragau
<i>Helochaeres femoralis</i> (Brullé, 1841)	Antigua and Barbuda, Argentina, Brazil, Colombia (?) [Nueva Granada], French Guiana (?)
Subfamily Chaetarthriinae	
Tribe Anacaenini	
<i>Crenitulus solstitialis</i> (Kirsch, 1873)	Bolivia, Brazil, Colombia [Quindío], Costa Rica, Guatemala, Mexico, Panama, Peru
<i>Crenitulus suturalis</i> (LeConte, 1866)	Canada to Argentina, Colombia [Quindío]
Subfamily Enochrinae	
<i>Enochrus circumcinctus</i> (Bruch, 1915)	Argentina, Colombia [Quindío], Paraguay, Uruguay
<i>Enochrus lampros</i> Kirsch, 1924b	Argentina, Bolivia, Brazil, Colombia [Quindío], Peru, Venezuela
<i>Enochrus paraensis</i> Fernández, 1997	Brazil, Colombia [Quindío]
<i>Enochrus pigrans</i> Knisch, 1925	Colombia [Valle del Cauca (Darién)]
<i>Enochrus pygmaeus</i> (Fabricius, 1792)	Bahamas, Colombia [Quindío], Cuba, Mexico, USA
<i>Enochrus sharpi</i> Gundersen, 1977	Colombia [Quindío], Mexico
<i>Enochrus sublongus</i> (Fall, 1924)	Argentina, Colombia [Quindío], Guatemala, Paraguay, USA
<i>Enochrus tropicus</i> (Kirsch, 1871)	Colombia [Cundinamarca (Bogotá)]
Subfamily Hydrophilinae	
Tribe Berosini	
<i>Berosus festai</i> Knisch, 1925	Colombia [Caldas? (S. José)], Ecuador
<i>Berosus maculinus</i> Knisch, 1921	Argentina, Bolivia, Brazil, Colombia**, Paraguay
<i>Berosus nigrinus</i> Knisch, 1921	Brazil, Colombia**
<i>Berosus singularis</i> Knisch, 1921	Brazil, Colombia**
<i>Berosus truncatipennis</i> Castelnau, 1840	Argentina, Bolivia, Brazil, Colombia [Quindío], Guatemala, Mexico, Nicaragua, Panama, Paraguay, Venezuela
<i>Berosus wintersteineri</i> Knisch, 1921	Brazil, Colombia**
<i>Berosus zimmermanni</i> Knisch, 1921	Argentina, Brazil, Colombia**, Paraguay
<i>Derallus terranova</i> Oliva, 1983	Brazil, Colombia [Quindío]
<i>Hemiosus aequatorialis</i> Oliva, 1994	Colombia [Quindío], Peru, Venezuela
Tribe Hydrophilini	
<i>Hydrobiomorpha colombica</i> Mouchamps, 1959	Colombia*
<i>Hydrophilus ensifer</i> Brullé, 1837	Argentina, Bolivia, Brazil, Colombia*, Ecuador, French Guyana, Paraguay, Suriname, Uruguay
<i>Tropisternus apicalpalpis</i> Chevrolat, 1834	Argentina, Bolivia, Brazil, Colombia* [Quindío], Costa Rica, Cuba, Lesser Antilles, Mexico, Nicaragua, Paraguay, Venezuela
<i>Tropisternus collaris</i> Fabricius, 1775	Argentina, Bolivia, Brazil, Colombia*, Cuba, Haiti, Lesser Antilles, Mexico, Paraguay, Peru, Puerto Rico, USA, Venezuela


Continued

Table 1. Continued.

Species	Geographic distribution
<i>Tropisternus knischi</i> Orchymont, 1943	Colombia [Cauca, Quindío]
<i>Tropisternus lancifer</i> Sharp, 1883	Colombia [Cundinamarca (Bogotá), Nueva Granada?]
<i>Tropisternus mergus</i> Say, 1835	Argentina, Bolivia, Colombia*, Cuba, Guatemala, Honduras, Lesser Antilles, Mexico, Nicaragua, Panama, Paraguay Venezuela
<i>Tropisternus oculatus</i> Sharp, 1882	Colombia*, Mexico, Venezuela
<i>Tropisternus ovalis</i> Castelnau, 1840	Argentina, Brazil, Colombia*, Cuba, Guatemala, Mexico, Paraguay
<i>Tropisternus sharpi</i> Orchymont, 1922	Argentina, Brazil, Colombia* [Quindío], Paraguay, Uruguay, Venezuela
Tribe Laccobiini	
<i>Oocyclus luteus</i> García-Hernández, 2009	Colombia [Quindío (Pijao)]
<i>Oocyclus morgani</i> García-Hernández, 2009	Colombia [Quindío (Calarcá)]
<i>Paracymus giganticus</i> Wooldridge, 1973	Colombia [Cundinamarca (Bogotá, Tocancipá)]
<i>Paracymus indigenus</i> Wooldridge, 1969	Colombia*, Costa Rica, Cuba, El Salvador, Guatemala, Honduras, Mexico
<i>Paracymus insularis</i> Wooldridge, 1973	Colombia [Magdalena (Barranquilla)]
<i>Paracymus limbatus</i> Wooldridge, 1973	Bolivia, Brazil, Colombia [Meta (Villavicencio)], Paraguay, Peru, Trinidad and Tobago, Venezuela
Subfamily Sphaeridiinae	
Tribe Coelostomatini	
<i>Dactylosternum subdpressum</i> (Castelnau, 1840)	Bolivia, Brazil, Colombia*, Cuba Guatemala, Guayana, Mexico, Nicaragua, Panama, Peru, Venezuela
<i>Dactylosternum subrotundum</i> (Fabricius, 1792)	Colombia (?), Peru
<i>Phaenonotum exstriatum</i> (Say, 1835)	Argentina?, Brazil?, Colombia [Quindío], Costa Rica, Cuba, Dominican Republic, Guatemala, Jamaica, Lesser Antilles, Mexico, Panama, Puerto Rico, USA
<i>Phaenonotum globulosum</i> (Mulsant, 1844)	Argentina, Brazil, Colombia [Bolívar (Cartagena)], Guatemala, Mexico, Nicaragua
<i>Phaenonotum</i> sp. 1	Colombia [Quindío]
Tribe Megasternini	
<i>Cercyon variegatus</i> Sharp, 1882	Argentina, Brazil, Colombia*, Guatemala, Jamaica, Lesser Antilles, Mexico, Nicaragua, Panama, Paraguay, Peru, Puerto Rico, Venezuela
<i>Motonerus apterus</i> Fikáček and Short, 2006	Colombia [Norte de Santander (Chinácota)], Venezuela
<i>Motonerus depressus</i> Fikáček and Short, 2006	Colombia [Caldas (La Aquadita)]
<i>Pelosoma cercyonoides</i> Sharp, 1882	Colombia [Quindío], Guatemala, and Mexico
<i>Pelosoma lafertei</i> Mulsant, 1844	Argentina, Brazil, Colombia [Quindío], France, Guatemala, Italia, Mexico, Nicaragua, Panama, Venezuela
<i>Pelosoma sordidum</i> Sharp, 1882	Colombia*, Costa Rica, Ecuador, Guatemala, Mexico, Trinidad and Tobago
Tribe Omicrini	
<i>Aculomicrus testudo</i> Smetana, 1975	Colombia [Vale del Cauca (Anchicaya)]
<i>Omicrus intermedius</i> Smetana, 1975	Colombia [Amazonas (Leticia)], Costa Rica, Mexico, Panama, USA
<i>Omicrus laevipennis</i> Smetana, 1975	Colombia [Vale del Cauca (Anchicaya, San Andres)], Panama

Table 2. Distribution records of Hydrophilidae species in Quindío Department municipalities (*New records from Colombia, † New records from Quindío Department).

Species	Filandia	Quimbaya	Circasia	Salento	Montenegro	Armenia	Calarcá	La Tebaida	Córdoba
<i>Helochares abbreviatus</i> †								x	
<i>H. atratus</i> *†								x	
<i>Crenitulus solstitialis</i> *†	x	x	x	x	x		x	x	x
<i>C. suturalis</i> *†	x	x	x	x	x		x	x	x
<i>Enochrus circumcinctus</i> *†		x						x	
<i>E. lampros</i> *†								x	
<i>E. paraensis</i> *†		x							
<i>E. pygmaeus</i> *†	x	x					x	x	x
<i>E. sharpi</i> *†				x	x			x	
<i>E. sublongus</i> *†		x			x			x	
<i>Berosus truncatipennis</i> *†								x	
<i>Derallus terranova</i> *†								x	
<i>Hemiosus aequatorialis</i> *†		x						x	x
<i>Hydrophilus ensifer</i> †					x			x	
<i>Tropisternus apicalpis</i> †					x		x	x	
<i>T. collaris</i> †				x				x	x
<i>T. sharpi</i> †								x	
<i>Paracymus insularis</i> †		x						x	x
<i>P. limbatus</i> †		x						x	
<i>Phaenonotum exstriatum</i> *†								x	
<i>P. sp. 1</i> *†								x	
<i>Pelosoma cercyonoides</i> *†		x	x			x			
<i>P. lafertei</i> *†		x				x	x		



Figures 1–12. Dorsal habitus of some species of Hydrophilidae found in Quindío Department, Colombia. **1, 2.** *Helachares* spp. **1.** *H. abbreviatus*, **2.** *H. atratus*. **3, 4.** *Crenitulus* spp. **3.** *C. solstitialis*. **4.** *C. suturalis*. **5–10.** *Enochrus* spp. **5.** *E. circumcinctus*. **6.** *E. lampros*. **7.** *E. paraensis*. **8.** *E. pygmaeus*. **9.** *E. sharpi*. **10.** *E. sublongus*. **11.** *Berosus truncatipennis*. **12.** *Hydrophilus ensifer*. Scale bars = 1mm.

lobe; apices slightly rounded with inner margins produced and tooth-like projection on outer margins; dorsal lobule narrow, swollen apically with two protuberances; ventral lobule shorter and wider than dorsal lobule, acuminate at apex.

Geographic distribution. Argentina (Misiones), Bolivia, Brazil, Colombia, Costa Rica, Cuba, Guyana, Lesser Antilles, Panama, Paraguay, Surinam, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°41'30.1" N, 075°83'98.1" W, 1057 m above mean sea level [a.m.s.l.], 2014-X-10. L.M. González coll., CIUQ 10636–10642 (8 males, 10 females)

Remarks. Specimens were collected from ponds with muddy/sandy and muddy/leaf litter substrates and in marshes with predominance of muddy substrate and saturated litter.

Helochares atratus Bruch, 1915 (Figs 2, 25)

Helochares atratus Bruch 1915: 451. (Not a syn. of *gibbus* Brullé (= *ventricosus* Bruch), as in Orchymont 1926: 236).

Helochares (Helochares) atratus Knisch 1925: 4.

Helochares (Helochares) pallipes Orchymont 1926: 236–237.

Helochares (s. str.) *parhedrus* Orchymont 1939: 259. Syn.: Fernández 1982: 35.

Diagnosis. Body 6.10–6.80 mm long. Dorsal punctation on elytra finer and sparser than head and pronotum. Serial punctures on rows 8, 9 and 10 of elytra deeply impressed. Mesoventral process slightly elevated, rounded in lateral view. Last ventrite weakly notched. Parameres longer than median lobe, apices slightly rounded; dorsal lobule wide, bifid at apex, each production of apex slightly acute, turned inward.

Geographic distribution. Argentina, Brazil, Ecuador, and Paraguay. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04. L.M. González coll., CIUQ 10643 (2 males).

Remarks. Specimens were collected from marshes with muddy substrate and saturated litter.

Subfamily Chaetarthiinae

Tribe Anacaenini

Crenitulus solstitialis (Kirsch, 1873) (Figs 3, 26)

Hydrobius solstitialis Kirsch 1873: 133.

Paracymus solstitialis (Kirsch)—Zaitzev 1908: 380; generic assignment.

Anacaena solstitialis (Kirsch)—Orchymont 1933: 304; generic assignment.

Anacaena perplexa Orchymont 1942: 39.

Anacaena perspicua Orchymont 1942: 40.

Crenitulus solstitialis (Kirsch)—Fikáček and Vondráček 2014: 502.

Diagnosis. Body 1.40–1.90 mm long. Body oval in shape.

Clypeus, front, pronotum and elytra black or dark to light brown. Dorsal punctation on head, pronotum and elytra fine and dense. Elytra with longitudinal rows of punctures black, sometimes seeming to form striae. Antennae with 9 antennomeres. Mesoventral process slightly elevated. Procoxae pubescent, with five spine-like projections. Parameres longer than phallobase; median lobe narrow and crown-shaped at apex.

Geographic distribution. Widely distributed in North, Central, and South America. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'50.4" N, 075°51'09.3" W, 998 m a.m.s.l., 2015-I-30. L.M. González coll., CIUQ 10443–10468 (1083 males, 1767 females). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03. L.M. González coll., CIUQ 10447–10482 (136 males, 229 females). Córdoba Municipality, Río Verde neighborhood, 04°24'15.0" N, 075°43'32.6" W, 1100 m a.m.s.l., 2015-I-27. L.M. González coll., CIUQ 10473–10480 (520 males, 871 females). Circasia Municipality, Membrillal neighborhood, 04°37'35.0" N, 075°38'30.0" W, 1700 m a.m.s.l., 2007-IX-22, A.L. García coll., CIUQ 12725 (6 males, 11 females). Calarcá Municipality, La Virginia neighborhood, 04°29'30.0" N, 075°38'27.0" W, 1650 m a.m.s.l., 2007-IX-23, A.L. García coll., CIUQ 12726 (21 males, 36 females). Salento Municipality, Boquía neighborhood, 04°38'46.1" N, 075°34'52.56" W, 2500 m a.m.s.l., 2007-IX-25, A.L. García coll., CIUQ 12727 (8 males, 13 females). Filandia Municipality, Cruces neighborhood, 04°41'17.0" N, 075°36'29.2" W, 1800 m a.m.s.l., 2008-II-14, A.L. García coll., CIUQ 12728 (5 males, 8 females). Montenegro Municipality, Silencio neighborhood, 04°29'50.0" N, 075°47'16.2" W, 1180 m a.m.s.l., 2007-II-02, A.L. García coll., CIUQ 12729 (4 males, 6 females).

Remarks. Specimens were found on the banks of rivers, ponds and marshes, on different substrates: submerged vegetation, stony/sandy, and muddy.

Crenitulus suturalis (LeConte, 1866) (Figs 4, 27)

Limnebius suturalis Leconte 1866: 366–367.

Hydrobius suturalis (Leconte)—Horn 1873: 136, generic assignment.

Creniphilus suturalis (Leconte)—Horn 1890: 272, generic assignment.

Paracymus suturalis (LeConte)—Zaitzev 1908: 380, generic assignment.

Crenitulus suturalis (LeConte)—Winters, 1926: 54, generic assignment.

Anacaena suturalis (Leconte)—Orchymont, 1933: 302, generic assignment.

Hydrobius debilis Sharp 1882: 65.

Paracymus debilis (Sharp)—Zaitzev 1908: 379, generic assignment.

Anacaena debilis (Sharp)—Orchymont 1933: 305, generic assignment.

Anacaena moreirai Orchymont 1921a: 245.

Anacaena pescheti Orchymont 1921a: 247.

Paracymus attenuatus Orchymont 1921a: 248.

Anacaena attenuata (Orchymont)—Orchymont 1933: 305, generic assignment.

Anacaena cordobana Knisch 1924b: 122.

Anacaena morosa Orchymont 1942: 41.

Anacaena peta Orchymont 1942: 42.

Anacaena morula Orchymont 1942: 46.

Anacaena sternalis Leech 1948: 447.

Crenitulus suturalis (LeConte)—Fikáček and Vondráček, 2014: 502.

Diagnosis. Body 1.50–2.10 mm long. Body elongate oval in shape, attenuated posteriorly. Dorsal punctation on head and elytra fine. Clypeus, front, pronotum and elytra black; pronotum and elytra with brown to light brown area at margins. Antennae with 9 antennomeres, apical antennomere elongate. Mesoventral process weakly elevated or absent. Pubescence on metafemur sparse. Parameres almost as long as phallobase, slightly acuminate at apex. Median lobe shorter than parameres, crown-shaped at apex.

Geographic distribution. Widely distributed from Canada to Argentina. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10487–10494 (20 males, 14 females). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10483–10486, CIUQ 10495–10497, CIUQ 10506 and CIUQ 10507 (169 males, 114 females). Córdoba Municipality, Santo Domingo neighborhood, 04°24'19.8" N, 075°43'35.8" W, 1112 m a.m.s.l., 2015-I-27, L.M. González coll., CIUQ 10498–10505 (30 males, 20 females). Calarcá Municipality, La Virginia neighborhood, 04°29'30.0" N, 075°38'27.0" W, 1650 m a.m.s.l., 2007-IX-23, A.L. García coll., CIUQ 12730 (15 males, 10 females). Salento Municipality, San Juan de Carolina neighborhood, 04°35'46.4" N, 075°43'37.5" W, 1510 m a.m.s.l., 2007-IX-25, A.L. García coll., CIUQ 12731 (19 males, 12 females). Circasia Municipality, Membrillal neighborhood, 04°37'35.0" N, 075°38'30.0" W, 1700 m a.m.s.l., 2007-IX-22, A.L. García coll., CIUQ 12732 (35 males, 23 females). Filandia Municipality, Cruces neighbourhood, 04°41'17.0" N, 075°36'29.2" W, 1800 m a.m.s.l., 2008-II-14, A.L. García coll., CIUQ 12733 (3 males, 2 females). Montenegro Municipality, Silencio neighborhood, 04°29'50.0" N, 075°47'16.2" W, 1180 m a.m.s.l., 2007-II-02, A.L. García coll., CIUQ 12734 (16 males, 11 females).

Remarks. Specimens were collected in ponds, riverbanks and marshes, on submerged vegetation, as well as muddy and stony/sandy substrates.

Subfamily Enochrinae

***Enochrus (Methydrus) circumcinctus* (Bruch, 1915)**
(Figs 5, 28)

Philhydrus circumcinctus Bruch 1915: 454.

Enochrus (Lumetus) circumcinctus (Bruch)—Knisch 1924a: 208.

Enochrus circumcinctus (Bruch)—Blackwelder 1944: 172.

Enochrus (Methydrus) circumcinctus (Bruch)—Fernández 1989: 126

Diagnosis. Body 3.20–4.25 mm in length. Head dark

brown to black with triangular reddish-brown area in front of eyes; elytra dark brown, pale brown at margins. Prosternum not carinate. Mesoventral process with laminar medial elevation, triangular, slightly concave at anterior portion, rounded anteriorly with a small tooth-like projection pointed posteriad. Last abdominal ventrite deeply notched. Parameres wide at base, acute at apex, longer than median lobe; median lobe triangular, strongly acute at apex.

Geographic distribution. Argentina, Paraguay and Uruguay. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10585–10593 (42 males, 99 females). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10594 (1 male)

Remarks. This species was found in marshes on submerged vegetation, in temporary ponds on muddy/stony substrates, and in streams on stony/sandy substrates.

***Enochrus (Methydrus) lampros* Kirsch, 1924**
(Figs 6, 29)

Enochrus (Lumetus) lampros Kirsch, 1924b: 124.

Enochrus (Methydrus) lampros Kirsch; Fernández, 1997: 28.

Diagnosis. Body 3.15–3.85 mm long. Head black with triangular yellow area in front of eyes; pronotum and elytra light brown. Dorsal punctation on elytra weaker than those on head and pronotum. Prosternum strongly carinate. Mesoventral process elevated. Last abdominal ventrite deeply notched, as deep as a quarter of the ventrite length. Parameres as long as phallobase, longer than median lobe, rounded at apex, outer margin concave at medial portion.

Geographic distribution. Argentina, Bolivia, Brazil, Perú and Venezuela. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Murillo neighborhood, 04°29'01.5" N, 075°45'44.2" W, 1050 m a.m.s.l., 2015-I-24, A.L. García coll., CIUQ 12735 (4 males, 2 females)

Remarks. Specimens were found in ponds on stony substrate and submerged vegetation.

***Enochrus (Methydrus) paraensis* Fernández, 1997**
(Figs 7, 30)

Diagnosis. Body 2.80–3.60 mm long. Head, pronotum and elytra light brown. Prosternum slightly elevated with a small tooth-like projection on anterior portion. Mesoventral process strongly carinate, posterior margin of carina angulate in lateral view. Parameres longer than phallobase, rounded at apex; median lobe shorter than parameres, convex at apex.

Geographic distribution. Brazil. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, Quimbaya Municipality, Manabí neighborhood, 04°36'46.7" N, 075°51'59.6" W, 995 m a.m.s.l., 2015-II-03-02, L.M. González coll., CIUQ 10602 and 10603 (9 males, 4 females)

Remarks. Specimens were collected in streams on stony/sandy substrates.

Enochrus (Methydus) pygmaeus (Fabricius, 1792) (Figs 8, 31)

Hydrophilus pygmaeus Fabricius, 1792: 186.

Hydrobius pygmaeus (Fabricius)—Gemminger and Harold 1868: 480.

Enochrus (Lumetus) pygmaeus (Fabricius)—Zaitzev 1908: 388 (as genus dubium) (generic placement confirmed by Orchymont 1933: 307).

Enochrus (Methydus) pygmaeus (Fabricius)—Young 1954: 176.

Enochrus (Methydus) rossi Leech 1948: 451. Syn.: Gundersen 1977: 257.

Diagnosis. Body 2.70–4.80 mm long. Head, pronotum and elytra light brown, part of clypeus black. Prosternum slightly carinate. Mesoventral process with laminar medial elevation, triangular, anterior margin with a small tooth-like projection pointed posteriad. Parameres as long as phallobase, narrow and rounded at apex with apical outer margins convex; median lobe shorter than parameres, narrow and strongly acute at apex.

Geographic distribution. Bahamas Islands, Belize, Cuba, Costa Rica, Dominican Republic, Haiti, Jamaica, Mexico, Nicaragua, Puerto Rico, USA, and Virgin Islands. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'47.6" N, 075°50'29.2" W, 1064 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10514–10517, and CIUQ 10519–10536 (581 males, 421 females). Quimbaya Municipality, La Buenavista neighborhood, 04°37'30.5" N, 075°44'40.3" W, 1291 m a.m.s.l., 2015-I-24, L.M. González coll., CIUQ 10537 (1 male). Quimbaya Municipality, Manabí, neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10518, CIUQ 10545 and CIUQ 10546 (38 males, 28 females). Córdoba Municipality, Río Verde neighborhood, 04°24'11.3" N, 075°43'20.9" W, 1139 m a.m.s.l., 2015-I-31, L.M. González coll., CIUQ 10538, CIUQ10539, CIUQ10543, and CIUQ 10544 (12 males, 8 females). Córdoba Municipality, Santo Domingo neighborhood, 04°24'26.8" N, 075°43'30.5" W, 1158 m a.m.s.l., 2015-I-27, L.M. González coll., CIUQ 10540–CIUQ 10542 (11 males, 9 females). Calarcá Municipality, Santo Domingo neighborhood Bajo, 04°24'32.4" N, 075°43'27.1" W, 1450 m a.m.s.l., 2007-IX-23, A.L. García coll., CIUQ 12736 (4 males, 2 females). Filandia Municipality, La Julia neighborhood, 04°24'35.0" N, 075°40'12.4" W, 1020 m a.m.s.l., 2008-I-

20, A.L. García coll., CIUQ 12737 (1 male).

Remarks. Specimens were collected in ponds on muddy, muddy/sandy, muddy/leaf litter substrates, as well as in marshes on submerged vegetation and in streams on stony/sandy substrate.

Enochrus (Methydus) sharpi Gundersen, 1977 (Figs 9, 32)

Enochrus (Methydus) sharpi Gundersen 1977: 262.

Diagnosis. Body 3.00–4.20 mm long. Head black, with triangular light brown area in front of eyes; pronotum and elytra light brown. Prosternum not carinate, completely pubescent. Mesoventral process rectangular, with a small tooth-like projection pointed posteriad. Last abdominal ventrite shallowly notched. Parameres longer than phallobase, laterally convex; median lobe acute, triangular, with corona subapical (on apical third); phallobase as long as parameres, narrowing at distal half.

Geographic distribution. Mexico and Puerto Rico. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, Montenegro Municipality, El Gigante neighborhood, 04°33'55.2" N, 075°48'54.4" W, 1060 m a.m.s.l., 2007-VIII-30, A.L. García coll., CIUQ 13275 (1 male, 2 females). La Tebaida Municipality, Pizamal neighborhood, 04°25'24.0" N, 075°51'36.0" W, 1140 m a.m.s.l., 2007-X-03, A.L. García coll., CIUQ 13276 (1 male, 1 female). Salento Municipality, Boquia neighborhood, 04°38'46.1" N, 075°34'52.5" W, 2200 m a.m.s.l., 2007-IX-28, A.L. García coll., CIUQ 13277 (2 males, 3 females).

Remarks. Specimens were collected in ponds on grass with muddy substrates.

Enochrus (Methydus) sublongus (Fall, 1926) (Figs 10, 33)

Philhydus elongatulus Fall 1924: 85, primary homonym of *Philhydus elongatulus* MacLeay, 1871.

Philhydus sublongus Fall, 1926: 125, replacement name for *P. elongatulus* Fall.

Enochrus (Methydus) sublongus (Fall)—Winters 1927: 20.

? *Enochrus (Methydus) curialis* Knisch 1924c: 55. Syn. *dubium*: Gundersen 1978: 24.

Diagnosis. Body 2.25–2.70 mm long. Coloration yellow, dark brown to black on medial portion of clypeus and on front. Dorsal punctation on head, pronotum and elytra weakly marked. Prosternum not carinate. Mesoventral process with a slightly longitudinal elevation, not laminar. Parameres slender on distal third, acuminate at apex, strongly convex laterally; median lobe acute, subtriangular.

Geographic distribution. Argentina, Guatemala, Paraguay, and USA. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'

46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10568–10582 (221 males, 196 females). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10567, CIUQ 10583 and CIUQ 10584 (2 males, 1 female). Montenegro Municipality, El Gigante neighborhood, 04°34'22.3" N, 075°51'11.6" W, 1060 m a.m.s.l., 2007-VIII-30, A.L. García coll., CIUQ 13278 (4 males, 2 females).

Remarks. Specimens were found in streams on stony/sandy substrate, in marshes on muddy substrate and submerged vegetation, and in ponds on muddy, muddy/sandy, muddy/leaf litter and muddy/stony substrates.

Subfamily Hydrophilinae

Tribe Berosini

Berosus truncatipennis Castelnau, 1840

(Figs 11, 34, 35)

Berosus truncatipennis Castelnau 1840: 56.

Berosus (*Enoplurus*) *truncatipennis* Castelnau—Zaitzev 1908: 357.

Berosus truncatipennis Castelnau—Oliva 1989: 184.

Diagnosis. Body 5.60–6.50 mm long. Body elongate in shape. Dorsal coloration pale brown; pronotum darker on medial portion, with a pale brown medial line, suboval in shape. Prosternum not carinate. Mesoventral process laminar, slender, convex anteriorly, with acute tooth-like projection on anterior portion and concave on posterior margin in lateral view. Last abdominal ventrite deeply notched. Parameres wide, acuminate at apex, with long ventral setae on apical portion, membranous inner appendices present; median lobe slender, cylindrical, shorter than half of the parameres length, apices slightly bifid.

Geographic distribution. Argentina, Bolivia, Brazil, Peru, Panama, Paraguay, and Venezuela. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'57.8" N, 075°50'49.9" W, 1064 m a.m.s.l., 2015-I-28, L.M. González coll., CIUQ 10430–CIUQ10442 (119 males, 110 females).

Remarks. Specimens were collected in marshes on muddy substrate and on submerged vegetation; in ponds on muddy/sandy, muddy/stony and muddy/leaf substrates.

Derallus terranova Oliva, 1983 (Figs 13, 37)

Derallus terranova Oliva 1983: 348.

Diagnosis. Body 2.20–2.30 mm long. Dorsal coloration evenly black. Punctuation on head fine and dense, on pronotum fine and sparse, on elytra strongly impressed forming longitudinal striae. Mesoventral process elevated with a tooth-like projection on anterior margin and a posterior elevation. Parameres shorter than median lobe, widened and rounded at apex; median lobe cylindrical, apex rounded; phallobase somewhat as long as parameres.

Geographic distribution. Brazil (Amazonia), Panama, and Venezuela. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°41'30.1" N, 075°83'98.1" W, 1057 m a.m.s.l., 2014-X-10, L.M. González coll., CIUQ 10508–10511 (5 males, 3 females).

Remarks. Specimens were collected in marshes on muddy substrate and on submerged vegetation.

Hemiosus aequatorialis Oliva, 1994 (Figs 14, 38)

Hemiosus aequatorialis Oliva 1994: 283.

Diagnosis. Body 2.30–2.50 mm long. Head and pronotum black with greenish to reddish metallic sheen; elytra pale brown with black maculae. Punctuation on head and pronotum dense and strongly impressed; elytral striae strongly impressed. Mesoventral process strongly elevated, deeply hollowed longitudinally, in lateral view anterior margin strongly projected as a tooth, followed by a smaller elevation almost as high as anterior tooth, posterior portion also salient. Parameres longer than median lobe, apices slightly acuminate, concave on outer margin, straight on inner margin; median lobe wide basally, slightly slender and rounded at apex; phallobase somewhat shorter than parameres.

Geographic distribution. Peru and Venezuela. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'47.5" N, 075°51'11.1" W, 1030 m a.m.s.l., 2015-I-29, L.M. González coll., CIUQ 10628–10632 (22 males, 16 females). Córdoba Municipality, Santo Domingo neighborhood, 04°24'20.3" N, 075°43'29.1" W, 1166 m a.m.s.l., L.M. González coll., CIUQ 10634 (2 males, 1 female). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10635 (6 males, 5 females).

Remarks. Specimens were collected along riverbanks and in temporary ponds on muddy leaf litter as well as on muddy/stony and stony/sandy substrates.

Tribe Hydrophilini

Hydrophilus ensifer Brullé, 1837 (Figs 12, 36)

Hydrophilus ater Olivier 1792: 125, primary homonym of *Hydrophilus ater* Gmelin, 1790.

Hydrous ater (Olivier)—Cristofori and Jan 1832: 31.

Stethoxus ater (Olivier)—Solier 1834: 307.

Hydrous (*Pagipherus*) *ater* (Olivier)—Kuwert 1893: 90.

Hydrophilus ater Fabricius 1792: 183, primary homonym of *Hydrophilus ater* Gmelin, 1790 and *Hydrophilus ater* Olivier, 1792. Syn.: Castelnau 1840: 49.

Hydrophilus (*Hydrocus*) *ensifer* Brullé 1837: 52. Syn.: Bedel 1881a: xciv (sub nomine *ater* Olivier).

Stethoxus (s. str.) *ensifer* (Brullé)—Orchymont 1943: 69 (reinstated as valid name).

Hydrophilus (Hydrocus) ovalis Brullé 1837: 53. Syn.: Bedel 1881a: xciv.

Hydrous (Pagipherus) brasiliensis (Castelnau)—Kuwert, 1893: 90.

Diagnosis. Body 30–37 mm long. Body entirely black, except for the light brown labial and maxillary palps and antennae. Meso and metaventral process forming a long keel, posteriorly projected as a long spine which almost reaches posterior margin of the abdominal ventrite II. Glabrous portion of the ventrites III to V triangular in shape. Parameres apically projected outward with lateral tooth-like projection, slender and rounded at apex; median lobe subtriangular, shorter than parameres; phallobase longer than parameres.

Geographic distribution. Argentina, Bolivia, Brazil, Colombia, Costa Rica, Cuba, Dominican Republic, Ecuador, French Guyana, Guatemala, Mexico, Nicaragua, Panama, Paraguay, Surinam, Uruguay, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°41'30.1" N, 075°83'98.1" W, 1057 m a.m.s.l., 2014-X-10, L.M. González coll., CIUQ 10646–CIUQ 10650 (4 males, 5 females). Montenegro Municipality, El Gigante neighborhood, 04°34'22.3" N, 075°51'11.6" W, 1060 m a.m.s.l., 2007-VIII-30, A.L. García coll., CIUQ 13279 (1 male, 1 female).

Remarks. Specimens were collected in marshes on muddy substrate and submerged vegetation.

Tropisternus apicipalpis Chevrolat, 1834 (Figs 15, 39)

Hydrophilus apicipalpis Chevrolat 1834: 44.

Tropisternus apicipalpis (Chevrolat)—Castelnau 1840: 53.

Tropisternus (Cyphostethus) apicipalpis (Chevrolat)—Orchymont 1921b: 364.

Tropisternus (Pristoternus) apicipalpis (Chevrolat)—Orchymont 1943: 54.

Tropisternus agilis Castelnau, 1840: 53. Syn.: Orchymont 1921b: 364.

Diagnosis. Body 14–16 mm in length. Color black with dark green sheen. Punctuation on head fine and strongly impressed. Meso-metaventral process forming a long keel posteriorly projected as a long spine that reaches the anterior third of abdominal ventrite II. Last ventrite with strong spine-like elevation projected posteriad with a tuft of short golden setae at apex. Parameres truncate apically, slightly longer than median lobe, weakly narrower at apex than at base; median lobe acuminate and wide at apex, slightly wider sub-basally; corona U-shaped, apically bifid; phallobase shorter than parameres.

Geographic distribution Argentina, Bolivia, Brazil, Costa Rica, Cuba, Lesser Antilles, Mexico, Nicaragua, Paraguay, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10685–10690 (57 males, 42

females). Montenegro Municipality, El Gigante neighborhood, 04°34'22.3" N, 075°51'11.6" W, 1060 m a.m.s.l., 2007-VIII-30, A.L. García coll., CIUQ 13280 (3 males, 1 female). Calarcá Municipality, Planadas Altas neighborhood, 04°27'05.4" N, 075°39'22.3" W, 2900 m a.m.s.l., 2007-IX-15, A.L. García coll., CIUQ 13281 (1 male).

Remarks. Specimens were found in marshes and stream margins on stony and muddy substrates and on submerged vegetation.

Tropisternus collaris (Fabricius, 1775) (Figs 16, 40)

Hydrophilus collaris Fabricius 1775: 229.

Dytiscus collaris (Fabricius)—Goeze 1777: 615.

Tropisternus collaris (Fabricius)—Castelnau 1840: 54.

Hydrophilus lineatus Dejean 1821: 50, nomum nudum. Syn.: Orchymont 1919: 159.

Tropisternus lineatus (Dejean)—Solier 1834: 310, nomum nudum.

Diagnosis. Body 6–10 mm long. Dorsal coloration pale-brown to yellow, pronotum dark brown to black on medial portion with metallic green suboval spot. Punctuation on head, pronotum and elytra fine and strongly impressed. Each elytron with four longitudinal metallic green lines. Prosternum elevated, medially opened, which receive anterior margin of mesoventrite process. Last abdominal ventrite with wide subtriangular protuberance, which overlaps $\frac{3}{5}$ of ventrite length. Parameres sinuate on inner margin, projected on outer margin, wide at base, longer than median lobe; median lobe slender at apex, abruptly widened on basal half; phallobase somewhat shorter than parameres.

Geographic distribution. Argentina, Bolivia, Brazil, Colombia, Cuba, Haiti, Lesser Antilles, Mexico, Paraguay, Peru, Puerto Rico, USA, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24'47.6" N, 075°50'29.2" W, 1064 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10694–10713 (665 males, 328 females). Salento Municipality, San Juan de Carolina neighborhood, 04°35'46.4" N, 075°43'37.5" W, 2007-IX-25, 1510 m a.m.s.l., A.L. García coll., CIUQ 13282 (3 males, 1 female). Córdoba Municipality, Corosal neighborhood, 04°24'19.5" N, 075°43'37.0" W, 1300 m a.m.s.l., 2007-X-05, A.L. García coll., CIUQ 13283 (3 males, 2 females).


Remarks. Specimens were collected in ponds, in marshes, and along riverbanks, on muddy/sandy, muddy/leaf litter, muddy/stony, and muddy substrates, as well as on submerged vegetation.

Tropisternus sharpi Orchymont, 1922 (Figs 17, 41)


Tropisternus nigrinus Sharp 1883: 107.

Tropisternus (Cyphostethus) sharpi Orchymont 1922: 21 (replacement name for *nigrinus* sensu Sharp). *Tropisternus (Pristoternus) sharpi* Orchymont—Bachmann 1981: 8.

Diagnosis. Body 11–13 mm long. Head, pronotum and elytra dark brown. Meso-metaventral process forming a



Figures 13–23. Dorsal habitus of some species of Hydrophilidae found in Quindío Department, Colombia. **13.** *Derallus terranova*. **14.** *Hemiosus aequatorialis*. **15–17.** *Tropisternus* spp. **15.** *T. apicipalpis*. **16.** *T. collaris*. **17.** *T. sharpi*. **18–19.** *Paracymus* spp. **18.** *P. insularis*. **19.** *P. limbatus*. **20, 21.** *Phaenonotum* spp. **20.** *P. exstriatum*. **21.** *P. sp.1*. **22, 23.** *Pelosoma* spp. **22.** *P. cercyonoides*. **23.** *P. lafertei*. Scale bars = 1mm.



Figures 24–35. Aedeagus of some species of Hydrophilidae found in Quindío Department, Colombia. **24, 25.** *Helachares* spp. **24.** *H. abbreviatus*. **25.** *H. atratus*. **26, 27.** *Crenitulus* spp. **26.** *C. solstitialis*. **27.** *C. suturalis*. **28–33.** *Enochrus* spp. **28.** *E. circumcinctus*. **29.** *E. lampros*. **30.** *E. paraensis*. **31.** *E. pygmaeus*. **32.** *E. sharpi*. **33.** *E. sublongus*. **34, 35.** *Berosus truncatipennis*.

Figures 36–47. Aedeagus of some species of Hydrophilidae found in Quindío Department, Colombia. **36.** *Hydrophilus ensifer*. **37.** *Derallus terranova*. **38.** *Hemiosus aequatorialis*. **39–41.** *Tropisternus* spp. **39.** *T. apicalpis*. **40.** *T. collaris*. **41.** *T. sharpi*. **42, 43.** *Paracymus* spp. **42.** *P. insularis*. **43.** *P. limbatus*. **44, 45.** *Phaenonotum* spp. **44.** *P. exstriatum*. **45.** *P. sp.1*. **46, 47.** *Pelosoma* spp. **46.** *P. cercyonoides*. **47.** *P. lafertei*.

long keel posteriorly projected as a long spine that reaches medially of abdominal ventrite II length. Last ventrite with a short spine-like posteriad, little evident, with a tuft of golden setae at apex. Parameres longer than phallobase, laterally convex, median lobe acute, triangular, phallobase as long as parameres, narrowing at distal half.

Geographic distribution. Argentina, Brazil, Colombia, Paraguay, Uruguay, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°41' 30.1"N, 075°83'98.1"W, 1057 m a.s.l., 201-X-10, L.M. González coll., CIUQ 10691–10693 (2 males, 1 female).

Remarks. Specimens were collected in marshes on mud and submerged vegetation.

Tribe Laccobiini

Paracymus insularis Wooldridge, 1973 (Figs 18, 42)

Paracymus limbatus Wooldridge 1973: 119.

Diagnosis. Body 1.80–2.00 mm long. Dorsal coloration metallic green, pale brown on pronotal margins and on posterior third of elytra. Prosternum carinate, with a rounded tooth-like projection on anterior margin. Parameres slightly longer than median lobe, outer margin sinuate at apex and slightly convex on inner margin; median lobe with apex sharp, narrowed on the medial

portion; phallobase shorter than parameres.

Geographic distribution. Colombia. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°25' 13.4" N, 075°49'57.2" W, 1034 m a.m.s.l., 2015-II-28, L.M. González coll., CIUQ 10673 and CIUQ 10674 (3 males, 1 female). Quimbaya Municipality, Manabí neighborhood, 04°36'46.7" N, 075°51'59.6" W, 995 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10670–10672, CIUQ 10675, and CIUQ 10676 (10 males, 7 females). Córdoba Municipality, Corosal neighborhood, 04°24'19.5" N, 075°43'37.0" W, 1200 m a.m.s.l., 2008-I-13, A.L. García coll., CIUQ 13284 (8 males, 5 females).

Remarks. Specimens were found in ponds and along riverbanks, on muddy/leaf litter, muddy/sandy, stony/sandy and muddy substrates.

Paracymus limbatus Wooldridge, 1973 (Figs 19, 43)

Paracymus limbatus Wooldridge 1973: 120.

Diagnosis. Body 1.70–1.90 mm in length. Dorsal coloration metallic green, pale brown on pronotal margins and on posterior third of elytra. Prosternum with slightly elevation, rounded on anterior margin. Mesoventral process elevated on anterior margin with a small tooth-like projection directed posteriad. Parameres convex on outer margin; acute and slightly curve at apex. Median lobe triangular, slightly convex laterally, wider at base; phallobase strongly shorter than parameres; rounded apically.

Geographic distribution. Bolivia, Brazil, Colombia, Paraguay, Peru, Trinidad and Tobago, and Venezuela. First record from Quindío Department.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24' 46.9" N, 075°50'34.2" W, 1031 m a.m.s.l., 2015-II-04, L.M. González coll., CIUQ 10651, CIUQ 10652, and CIUQ 10655–CIUQ 10666 (72 males, 62 females). Quimbaya Municipality, Manabí neighborhood, 04°36'48.4" N, 075°51'58.9" W, 965 m a.m.s.l., 2015-II-03, L.M. González coll., CIUQ 10653, CIUQ 10654, and CIUQ 10667–10669 (14 males, 12 females).

Remarks. Specimens were collected in ponds and marshes and along riverbanks, on muddy/sandy, muddy/leaf litter, muddy/stony, and stony/sandy substrates.

Subfamily Sphaeridiinae

Tribe Coelostomatini

Phaenonotum exstriatum (Say, 1835) (Figs 20, 44)

Hydrophilus exstriatus Say 1835: 171.

Cyclonotum estriatum [sic] (Say)—Melsheimer 1853: 36, unjustified emendation.

Phaenonotum estriatum [sic] (Say)—Fleutiaux and Sallé 1889: 377, unjustified emendation.

Coelostoma minutum Castelanu 1840: 58. Syn.: Zaitzev 1908: 405.

Cyclonotum minutum (Castelanu)—Gemminger and Harold 1868: 495.

Cyclonotum globulosum Mulsant 1844b: 167.

Cyclonotum semiglobosum Zimmermann 1869: 250. Syn.: Orchymont 1941: 7.

Phaenonotum semiglobosum (Zimmermann)—Zaitzev 1908: 405.

Phaenonotum dubium Sharp 1882: 98. Syn.: Smetana 1978: 14.

Phaenonotum exstriatum (Say)—Orchymont 1941: 7.

Diagnosis. Body 2.30–4.00 mm long. Body oval in shape. Dorsal coloration entirely black. Dorsal punctation on head and pronotum fine and dense, on elytra coarser. Meso-metaventral process strongly elevated, acuminate on anterior margin, widened posteriorly, with dense punctation on surface. Parameres as long as median lobe, slightly acuminate at apex, outer margin sinuate; median lobe wider than parameres, slightly acuminate at apex; phallobase shorter than parameres.

Geographic distribution. Argentina?, Brazil? Costa Rica, Cuba, Dominican Republic, Guatemala, Jamaica, Lesser Antilles, Mexico, Panama, Puerto Rico, and USA. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24' 48.9" N, 075°51'10.5" W, 979 m a.m.s.l., 2015-I-22, L.M. González coll., CIUQ 10683 and CIUQ 10684 (3 males, 2 females).

Remarks. Specimens were collected in a marsh on muddy substrate and on submerged vegetation.

Phaenonotum sp. 1 (Figs 21, 45)

Diagnosis. Body 1.60–2.40 mm long. Body oval in shape. Dorsal coloration entirely black. Dorsal punctation on head and pronotum fine and dense, on elytra sparser. Meso-metaventral process slender and acuminate on anterior margin, widened posteriorly. Parameres as long as median lobe, apex slightly acuminate, convex on outer margin, wide medially; median lobe wider than parameres, slightly acuminate at apex; phallobase shorter than parameres.

Geographic distribution. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, La Tebaida Municipality, Pizamal neighborhood, 04°24' 50.2" N, 075°51'06.3" W, 1012 m a.m.s.l., 2015-I-30, L.M. González coll., CIUQ 10678–10682 (11 males, 19 females).

Remarks. Specimens were found in marshes on muddy substrates and on submerged vegetation. *Phaenonotum* sp. 1 was very difficult to identify given that in many cases the genitalia have slight differences. The external morphology of this species needs to be carefully studied.

Tribe Megasternini

Pelosoma cercyonoides Sharp, 1882 (Figs 22, 46)

Pelosoma cercyonoides Sharp 1882: 111.

Diagnosis. Body 2.00–2.30 mm long. Dorsal coloration dark brown to black. Each elytron with ten striae with



Figure 48. Distribution map of species of Hydrophilidae in the Quindío Department, Colombia.

strongly impressed punctures. Prosternum slightly carinate medially with a small rounded tooth-like projection on anterior margin. Mesoventral process flat, elongate, pentagonal, acuminate on anterior margin and truncate on posterior margin, with punctures coarse and strongly impressed. Abdominal ventrite I carinate medially, slightly overlapping anterior margin of ventrite II. Parameres longer than median lobe, widened at apex and base, concave on inner margin; median lobe wide at base, narrow and rounded at apex; phallobase strongly asymmetrical at base.

Geographic distribution. Guatemala and Mexico. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, Armenia Municipality, Caimo-Arcoris neighborhood, 04°27'55.0" N, 075°44'40.2" W, 1350 m a.m.s.l., 2010-XI-25, A.L García coll., CIUQ 13285 (2 males, 3 females). Circasia Municipality, Finca el Calamar neighborhood, 04°37'15.0" N, 075°39'16.3" W, 1480 m a.m.s.l., 2010-IX-22, A.L García coll., CIUQ 13286 (6 males, 4 females). Quimbaya Municipality, El Laurel neighborhood, Reserva Natural La Montaña del Ocaso, 04°34'59.9" N, 075°49'16.3" W, 1100 m a.m.s.l., 2010-VI-22, A.L García coll., CIUQ 13287 (3 males, 2 females). Quimbaya Municipality, Pizamal neighborhood, 04°25'24.0" N, 075°51'36.0" W, 950 m a.m.s.l., 2010-XII-21, A.L García coll., CIUQ 13288 (5 males, 3 females).

Remarks. Specimens were found in phytotelmata, including the inflorescence of *Calathea inocephala*, *C. lutea*, *Heliconia stricta*, and *H. latisphata*. These plants

have characteristic small water tanks and show different stages of decomposition of their inflorescences.

Pelosoma lafertei (Mulsant, 1844) (Figs 23, 47)

Cercyon bicolor Dejean 1833: 134, nomen nudum. Syn.: Mulsant 1844a: 186.

Cercyon littorale var. *minutum* Dejean 1833: 134, nomen nudum. Syn.: Mulsant 1844a: 186.

Pelosoma lafertei Mulsant 1844a: 185.

Cercyon lafertei (Mulsant)—Fairmaire and Laboulbène 1854: 256.

Cercyon (Pelosoma) lafertei (Mulsant)—Bedel 1881b: lxxxvii.

Pelosoma meridionale Bruch 1915: 468. Syn.: Orchymont 1941: 14.

Diagnosis. Body 2.10–2.50 mm long. Head and pronotum reddish-brown, elytra slightly darker. Punctures on elytra strongly impressed forming ten longitudinal striae on each elytron. Prosternum slightly carinate on anterior quarter, anteriorly elevated as a rounded small tooth. Mesoventrite flat, pentagonal, elongate. Parameres as longer as or slightly longer than median lobe, widened and rounded at apex; median lobe wide basally, narrow and rounded at apex; phallobase asymmetrical.

Geographic distribution. Argentina, Brazil, Guatemala, Mexico, Nicaragua, Panama, and Venezuela; introduced to France and Italy. First record from Quindío Department and from Colombia.

Material examined. Colombia, Quindío Department, Quimbaya Municipality, El Jasmin neighborhood, Reserva la Guajira, 04°39'17.2" N, 075°39'16.4" W, 980 m a.m.s.l., 2010-VII-24, A.L García coll., CIUQ 13289 (1 male, 4 females). Calarcá Municipality, La Bella neighborhood, 04°29'15.3" N, 075°49'31.6" W, 1300 m

a.m.s.l., 2010-XII-29, A.L. García coll., CIUQ 13290 (5 males, 4 females). Armenia Municipality, Neighborhood: Caimo-Arcóiris, 04°27'55.0" N, 075°44'40.2" W, 1350 m a.m.s.l., 2010-XI-25, A.L. García coll., CIUQ 13291 (3 males, 2 females). Quimbaya Municipality, El Laurel neighborhood, Reserva Natural La Montaña del Ocaso, 04°35'19.0" N, 075°49'56.7" W, 1100 m a.m.s.l., 2010-VI-22, A.L. García coll., CIUQ 13292 (2 males, 5 females).

Remarks. Specimens were found in phytotelmata in the inflorescence of *Calathea inocephala*, *C. lutea*, *Heliconia latispatha* and *H. stricta*. These plants have characteristic small water tanks and show different stages of decomposition of their inflorescences.

Discussion

Most of the knowledge about Colombian Hydrophilidae available is from late 19th to 20th centuries (e.g., Régimbart 1901, Orchymont 1921a, 1922, Zaitzev 1908, Leech 1943, Blackwelder 1944). Some of these works did not provide precise locality and almost never biological information. There is a clear lack of taxonomic data and knowledge on the distribution of this family, especially in Colombia, where the high diversity and the range of environments suitable to the occurrence of these beetles is little-studied.

Thirty-seven species of Hydrophilidae were previously known from Colombia. Even with the limited sampling made in Quindío Department, our study significantly improves our knowledge of this family with 16 new species records. Our list, as well as summarizing environmental and habitat information, gives an overview of the biodiversity of the Hydrophilidae species in the field. We hope that this study will be valuable as a foundation to guide students and researchers in future studies on the family. These data may enable the use of hydrophilid species in research on basic and applied ecology, sustainable use of water (Archangelsky and Durand 1992), biological control of agricultural pests (Koppenhofer and Schmutterer 1993), vectors of diseases (Shalan and Canyon 2009), and even forensic sciences (Almeida and Mise 2009).

Acknowledgements

We thank the staff of arthropod studies from Centro de Estudios y Investigaciones en Biodiversidad of the Universidad del Quindío (CIBUQ) and from Colección de Insectos of the Laboratorio de Entomología of Universidad del Quindío-Colombia (CIUQ) for encouragement and aid. We are indebted to Martin Fikáček and Albert Deler-Hernández who provided assistance and valuable information that made our work possible.

Authors' Contributions

LMGR and ALGH collected the specimens and data,

LMGR, ALGH and BC wrote the text.

References

- Almeida LM, Mise KM (2009) Diagnosis and key of the main families and species of South American Coleoptera of forensic importance. *Revista Brasileira de Entomologia* 53 (2): 227–244. <https://doi.org/10.1590/S0085-56262009000200006>
- Arce-Pérez R, Morón MA (2013) El género *Hydrophilus* (Coleoptera: Hydrophilidae: Hydrophilinae) en México y Centroamérica. *Revista Mexicana de Biodiversidad* 84: 140–152. <https://doi.org/10.7550/rmb.32113>
- Archangelsky M, Durand ME (1992) Description of the preimaginal stages of *Dibolocelus ovatus* (Gemminger and Harold, 1868) (Coleoptera, Hydrophilidae: Hydrophilinae). *Aquatic Insects* 14 (2): 107–116. <https://doi.org/10.1080/01650429209361471>
- Arias-Díaz DM, Reinoso-Flórez G, Guevara-Cardona G, Villa-Navarro FA (2007) Distribución espacial y temporal de los coleópteros acuáticos en la cuenca del río Coello (Tolima, Colombia). *Caldasia* 29 (1): 177–194.
- Bachmann A (1969) *Tropisternus (pristoternus) gaeae* una nueva especie de La República Argentina (Coleoptera-Hydrophilidae). *Revista de la Sociedad Entomológica Argentina* 31 (1–4): 93–96.
- Bachmann AO (1981) Claves para determinar las familias, las subfamilias y los géneros de Hydrophiloidea acuáticos, y las especies de Hydrophilinae, de la República de Argentina (Coleoptera). *Revista de la Sociedad Entomológica Argentina* 40 (1–4): 1–9.
- Balfour-Browne J (1939) Contribution to the study of the Palpicornia. Part III. *Annals and Magazine of Natural History* (2nd series) 4: 289–310.
- Bedel L (1881a) Synonymie de quelques Hydrophilidae et Sphaeridiidae exotiques décrits par Brullé. *Annales de la Société entomologique de France* (6) 1: xciv–xcv.
- Bedel L (1881b) Diverses synonymies nouvelles d'Hydrophilidae et de Sphaeridiidae. *Annales de la Société entomologique de France* (6) 1: lxxxvi–lxxxvii.
- Bedel L (1891) Synopsis des grands hydrophiles (genre *Stethoxus* Solier). *Revue d'Entomologie* 10: 306–323.
- Blackwelder RE (1944) Checklist of the coleopterous insects of Mexico, Central America, the West Indies, and South America. Part I. *Bulletin of the United States National Museum*, 185: 1–188.
- Bruch C (1915) Nuevas especies de coleópteros hidrofilidos. *Revista del Museo de la Plata* 19 (2): 447–470.
- Brullé A (1837) Famille des hydrophiliens [part; continued in 1841]. In: d'Orbigny A (Ed.) *Voyage dans l'Amerique méridionale*. Vol. 6, part 2. Insectes. P. Bertrand, Paris/Strasbourg, 52–56.
- Castelnau FLN (1840) Histoire naturelle des insectes coléoptères, avec une introduction renfermant l'anatomie et la physiologie des animaux articulés, par M. Brullé. *Histoire naturelle des Animaux articulés: annélidés, crustacés, arachnides, Myriapodes et Insectes*. Tome 2: 1–563. P. Duménil, Paris. <https://doi.org/10.5962/bhl.title.35290>
- Chevrolat A (1834 [1833–35]) *Coleopteres du Mexique*. 3e Fascicule. Impr. de G. Silbermann, Strasbourg, 200 pp.
- Clarkson B, Ferreira-Jr N (2009) Three new species of *Hemiosus* Sharp (Coleoptera: Hydrophilidae) and new state records of *Hemiosus fittkai* Oliva and *H. moreirai* d'Orchymont from Brazil. *Zootaxa* 2139: 61–68.
- Cristofori J de, Jan G (1832) *Catalogus in IV sectiones divisus rerum naturalium in musaeo exstantium Josephi de Cristofori et Georgii Jan. Sectio III, Enlomologia. Pars 1. Conspectus Melhodicus Insectorum*. Fasc. I, Coleoptera. Carmignani, Milano, 4 + 16 + iii pp.
- Dejean PFMA (1821) *Catalogue de la collection de coléoptères de M. le Baron Dejean*. Crevot, Paris, viii + 136 + 2 pp.
- Dejean PFMA (1833 [1833–37]). *Catalogue des Coléoptères de la collection de M. le Comte Dejean* (2 ed.). Méquignon-Marvis Père et Fils, Paris, 443 pp. [Pp. 1–176, 1833; pp. 177–256, 1834; pp. 257–360, 1835; 361–443, 1837.]

- Deler-Hernández A, Fikáček M, Cala-Riquelme F (2013) A review of the genus *Berosus* Leach of Cuba (Coleoptera, Hydrophilidae). *ZooKeys* 273: 73–106. <https://doi.org/10.3897/zookeys.273.4591>
- Fabricius JC (1775) *Systema entomologiae, sistens insectorum classes, ordines, genera, species, adjectis synonymis, locis, descriptionibus, observationibus*. Libraria Korte, Flensburgi et Lipsiae, 32 + 832 pp. <https://doi.org/10.5962/bhl.title.36510>
- Fabricius JC (1792) *Entomologia systematica emendata et aucta. Secundum classes, ordines, genera, species adjectis synonymis, locis, observationibus, descriptionibus*. vol. 1, pars 1. C.G. Proft, Hafniae, 20 + 330 pp.
- Fabricius JC (1801) *Systema eleutheratorum secundum ordines, genera, species: adiectis synonymis, locis, observationibus, descriptionibus*. vol. 1. Bibliopoli Academici Novi, Kiliae, 24 + 506 pp.
- Fairmaire L, Laboulbène A (1854 [1854–56]). *Faune entomologique française ou description des insectes qui se trouvent en France*. Vol. 1. Coléoptères. Deyrolle, Paris, 35 + 665 pp.
- Fall HC (1924) New species of North American Hydrobiini. *Journal of the New York Entomological Society* 32: 85–89.
- Fall HC (1926) Two new names and a correction in synonymy. *Bulletin of the Brooklyn Entomological Society* 21: 125.
- Fernández LA (1982) Notas sobre el género *Helochares* (Insecta, Coleoptera, Hydrophilidae). *Neotropica* 28 (79): 31–40.
- Fernández LA (1988) Contribución al conocimiento del género *Enochrus* Thomson. I. (Coleoptera: Hydrophilidae). *Physis* (Buenos Aires) B 46 (111): 85–89.
- Fernández LA (1989) Las especies de *Enochrus* descriptas por Bruch. Redescripciones basadas sobre material típico. *Revista de la Sociedad Entomológica Argentina* 46 (1–4): 21–128.
- Fernández LA (1994) Contribución al conocimiento del género *Enochrus* Thomson. (Coleoptera: Hydrophilidae) II. El subgénero *Hugoscottia* en la República Argentina. *Physis* (Buenos Aires) B 49 (116–117): 37–45.
- Fernández LA (1997) Nuevos aportes para el conocimiento del género *Enochrus* Thomson (Coleoptera: Hydrophilidae). *Physis* (Buenos Aires) B 53 (124–125): 21–29.
- Fikáček M, Short AEZ (2006) A revision of the Neotropical genus *Motonerus* Hansen (Coleoptera: Hydrophilidae: Sphaeriini). *Zootaxa* 1268: 1–38.
- Fikáček M, Vondráček D (2014) A review of *Pseudorygmodus* (Coleoptera: Hydrophilidae), with notes on the classification of the Anacaenini and on distribution of genera endemic to southern South America. *Acta Entomologica Musei Nationalis Pragae* 54 (2): 479–514.
- Fleutiaux E, Sallé A (1889 [1889–90]). *Liste des coléoptères de la Guadeloupe et descriptions d'espèces nouvelles*. *Annales de la Société entomologique de France* (6) 9: 351–484. [Only pp. 351–424 issued in 1889].
- García-Hernández AL (2009) Two new species of *Oocyclus* (Coleoptera: Hydrophilidae) from Colombia. *Revista Colombiana de Entomología* 35 (2): 250–252.
- Gemminger M, Harold B de (1868) *Catalogus Coleopteromm hucusque descriptorum synonymicus et systematicus*. Vol. 2. E.H. Gummi, Monachii, 425–752 + [6] pp [unnumbered index].
- Goeze JAE (1777) *Entomologische Beyträge zu des Ritter Linné zwölften Ausgabe des Natursystems*. Vol. 1. Weidmanns Erben und Reich, Leipzig, 16 + 736 pp.
- Gundersen RW (1977) New species and taxonomic changes in the genus *Enochrus* (Coleoptera: Hydrophilidae). *The Coleopterist's Bulletin* 31 (3): 251–272.
- Gundersen RW (1978) Nearctic *Enochrus*: Biology, Keys, Descriptions and Distribution (Coleoptera: Hydrophilidae). St Cloud State University, St Cloud, Minnesota, 54 pp.
- Hansen M (1991) The hydrophiloid beetles. Phylogeny, classification and a revision of the genera (Coleoptera, Hydrophiloidea). *Biologiske Skrifter* 40: 1–368.
- Hansen M (1999) *World Catalogue of Insects 2: Hydrophiloidea* (Coleoptera). Apollo Books, Stenstrup, 416 pp.
- Horn GH (1873) Revision of the genera and species of the tribe Hydrobiini. *Proceedings of the American Philosophical Society* 13: 118–137.
- Horn GH (1890) A revision of the Sphaeriini inhabiting boreal America. *Transactions of the American Entomological Society* 17: 279–314, pl. 9.
- Kirsch TFW (1873) Beiträge zur Kenntniss der Peruanischen Käferfauna auf Dr Abendroth's Sammlungen basirt. *Berliner entomologische Zeitschrift* 17: 121–152.
- Knisch A (1924a) Hydrophilidae. In: Junk W, Schenkling S (Eds) *Coleopterorum Catalogus*. Vol. 14, Part 79. W. Junk, Berlin, 306 pp.
- Knisch A (1924b) Neue neotropische Palpicornier. (Col. Hydrophilidae.—Op. 16.). *Wiener entomologische Zeitung* 41: 114–140.
- Knisch A (1924c) Zwei neue nearktische Palpicornier. (Col. Hydrophilidae Op. 13.). *Entomologischer Anzeiger* 4: 55–56.
- Knisch A (1925) Coleoptera aus dem tropischen Amerika. Hydrophilidae (Palpicornia). *Bollettino dei Musei di Zoologia e di Anatomia comparata della R. Università di Torino* (n. s.) 30 (39) (1924): 1–12.
- Komarek A (2004) Taxonomic revision of *Anacaena* Thomson, 1859. I. Afrotropical species (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 74: 303–349.
- Komarek A (2005) Taxonomic revision of *Anacaena* Thomson, 1859. II. Neotropical species (Coleoptera: Hydrophilidae). *Koleopterologische Rundschau* 75: 253–301.
- Komarek A, Beutel RG (2007) Phylogenetic analysis of Anacaenini (Coleoptera: Hydrophilidae: Hydrophilinae) based on morphological characters of adults. *Systematic Entomology* 32: 205–226. <https://doi.org/10.1111/j.1365-3113.2006.00359.x>
- Koppenhofer AM, Schmutterer H (1993) *Dactylosternum abdominale* (F.) (Coleoptera: Hydrophilidae): a predator of the banana weevil. *Biocontrol Science and Technology* 3: 141–147.
- Kuwert A (1893) Die grossen Hydrophiliden des Erdballs des Genus *Hydrous* Leach. *Deutsche entomologische Zeitschrift* 1893: 81–93.
- LeConte JL (1866) Additions to the coleopterous fauna of the United States. No. I. *Proceedings of the Academy of natural Sciences of Philadelphia*: 361–394.
- Leech HB (1943) *Tropisternus paredesi*, nuevo coleopteron acuatico palpicornio de Nayarit, Mexico (Coleoptera, Hydrophilidae). *Revista de la Sociedad Mexicana de Historia Natural* 4 (1–2): 17–20.
- Leech HB (1948) Contributions toward a knowledge of the insect fauna of Lower California. No. II. Coleoptera: Haliplidae, Dytiscidae, Gyrinidae, Hydrophilidae, Limnebiidae. *Proceedings of the California Academy of Sciences* (4) 24: 375–483
- Melsheimer FE (1853) *Catalogue of the described Coleoptera of the United States*. Revised by Haldeman SS, LeConte JL. Smithsonian Institution, Washington DC, xvi + 174 pp. -
- Mouchamps R (1959) Remarques concernant les genres *Hydriobiomorpha* Blackburn et *Neohydrophilus* Orchymont (Coléoptères Hydrophilides). *Bulletin et annales de la Société royale d'entomologie de Belgique* 95: 295–335.
- Mouchamps R (1963) Beiträge zur Kenntnis der Insektenfauna Boliviens XVIII. Coleoptera XIV. Sur quelques Berosini (Coléoptères-Hydrophilidae) de l'Amérique du Sud. *Mitteilungender Münchner Entomologischen Gesellschaft* 53: 118–149.
- Mulsant E (1844a) *Histoire naturelle des coléoptères de France*. Palpicornies. L. Maisson, Paris/Ch. Savy Jeune, Lyon, 7 + 196 pp., 1 pl., [errata et addenda: 197 pp.]
- Mulsant E (1844b) *Essai monographique du genre cyclonotum*. *Annales de la Société d'Agriculture de Lyon* 7: 167–180.
- Oliva A (1981) El género *Derallus* Sharp en la Argentina (Coleoptera: Hydrophilidae) *Revista de la Sociedad Entomológica Argentina* 40: 285–296.
- Oliva A (1983) *Derallus* de la Cuenca del Amazonas (Coleoptera: Hydrophilidae). *Revista de la Sociedad Entomológica Argentina* 42 (1–4): 343–351.
- Oliva A (1987) El género *Hemiosus* en los andes meridionales de la Patagónica (Coleoptera: Hydrophilidae). *Revista de la Sociedad*

- Entomológica Argentina 44 (3–4): 377–381.
- Oliva A (1989) El género *Berosus* (Coleoptera: Hydrophilidae) en América del Sur. Revista del Museo Argentino de Ciencias Naturales (Buenos Aires, Argentina) (Entomología) 6 (4): 57–235.
- Oliva A (1994) A revision of the genus *Hemiosus* Sharp, 1882 in South America (Coleoptera: Hydrophilidae). Bulletin and Annales de la Société, Royal Belge, Entomologie 130: 267–303.
- Oliva A (2010) Redescription of *Derallus strigipennis* Orchymont, 1940 (Coleoptera: Hydrophilidae), new records and a key to species of *Derallus* Sharp, 1882. Revista del Museo Argentino de Ciencias Naturales, Nueva Serie 12 (1): 91–97.
- Oliva A (2014) Distribution of the species of Berosini (Coleoptera: Hydrophilidae) in South America. Revista Colombiana de Entomología 40 (2): 272–280.
- Oliva A, Short AEZ (2012) Review of the *Berosus* Leach of Venezuela (Coleoptera, Hydrophilidae, Berosini) with description of fourteen new species. ZooKeys 206: 1–69. <https://doi.org/10.3897/zookeys.206.2587>
- Oliva A, Fernández LA, Bachmann AO (2002) Sinopsis de los Hydrophiloidea acuáticos de la Argentina (Insecta, Coleoptera). Monografías del Museo Nacional de Ciencias Naturales 2: 1–67.
- Oliva AG (1792) Encyclopédie méthodique. Dictionnaire des Insectes. Vol. 7 (Han-Mou). Pankouke, Paris, 827 pp. [Pl. 183, with illustrations of *Hydrophilus*, issued in 1797].
- Orchymont A d' (1919) Contribution à l'étude des sous-familles des Sphaeriidiinae et des Hydrophilinae (Col. Hydrophilidae). Annales de la Société entomologique de France 88: 105–168.
- Orchymont A d' (1921a) Palpicornia de l'Amérique du Sud. Bulletin et Annales de la Société entomologique de Belgique, 61: 244–255.
- Orchymont A d' (1921b) Le genre *Tropisternus* I (Col. Hydrophilidae). Annales de la Société entomologique de Belgique 61: 349–374.
- Orchymont A d' (1922) Le genre *Tropisternus* (Col. Hydrophilidae). II. Annales de la Société entomologique de Belgique 62: 11–47.
- Orchymont A d' (1926) Contribution à l'étude des Hydrophilides VI. Bulletin et Annales de la Société entomologique de Belgique 66: 201–248.
- Orchymont A d' (1933) Contribution à l'étude des Palpicornia VIII. Bulletin et Annales de la Société entomologique de Belgique 73: 271–273.
- Orchymont Ad' (1936) Quelques synonymies nouvelles d'Hydrophilidae (Col.). Bulletin du Musée royal d'Histoire naturelle de Belgique 12 (23): 1–29.
- Orchymont A d' (1939) Notes sur quelques *Palpicornia* de la République Argentine. Revista de la Sociedad entomológica Argentina 10: 253–264.
- Orchymont A d' (1941) Palpicornia (Coleoptera). Notes diverses et espèces nouvelles I. Bulletin du Musée royal d'Histoire naturelle de Belgique 17 (1): 1–23.
- Orchymont A d' (1942) Contribution à l'étude de la tribu Hydrobiini Bedel, spécialement de sa sous-tribu Hydrobiae (Palpicornia–Hydrophilidae). Mémoires du Musée royal d'Histoire naturelle de Belgique 24 (2): 1–68.
- Orchymont A d' (1943) Faune du Nord-Est Brésilien (récoltes du Dr O. Schubart). Palpicornia. Mémoires du Musée royal d'Histoire naturelle de Belgique 28 (2): 1–85.
- Peck SB (2006) The beetle fauna of Dominica, Lesser Antilles (Insecta: Coleoptera): Diversity and distribution. Insecta Mundi 20 (3–4): 165–210.
- Régimbar M (1901) Revision des grands hydrophiles. Annales de la Société entomologique de France 70: 188–232.
- Say T (1835) Descriptions of new North American coleopterous insects, and observations on some already described. Boston Journal of Natural History 1: 151–203.
- Shaan EA, Canyon DV (2009) Aquatic insect predators and mosquito control. Tropical Biomedicine 26 (3): 223–261.
- Sharp D (1882 [1882–87]). Insecta. Coleoptera. Vol. 1, part 2 (Haliplidae, Dytiscidae, Gyrinidae, Hydrophilidae, Heteroceridae, Parnidae, Georissidae, Cyathoceridae, Staphylinidae). In: Godman FD, Salvin O (Eds) Biologia Centrali-Americana 16: xv + 824 pp. Taylor and Francis, London. [Only pp. 1–144 issued in 1882].
- Sharp D (1883) Revision of the species included in the genus *Tropisternus* (fam. Hydrophilidae). Transactions of the Entomological Society of London 1883: 91–117.
- Short AEZ (2004) Review of the *Enochrus* Thomson of the West Indies (Coleoptera: Hydrophilidae). Koleopterologische Rundschau 74: 351–361.
- Short AEZ (2005) A review of the subtribe Acidocerina of Central America with special reference to Costa Rica (Coleoptera: Hydrophilidae). Koleopterologische Rundschau 75: 191–226.
- Short AEZ, Fikáček M (2011) World catalogue of the Hydrophiloidea (Coleoptera): additions and corrections II (2006–2010). Acta Entomologica Musei Nationalis Pragae 51 (1): 83–122.
- Short AEZ, Fikáček M (2013) Molecular phylogeny, evolution and classification of the Hydrophilidae (Coleoptera). Systematic Entomology 38 (4): 723–752. <https://doi.org/10.1111/syen.12024>
- Short AEZ, García M, Girón JC (2017) Revision of the Neotropical water scavenger beetle genus *Globulosis* García, 2001 (Coleoptera: Hydrophilidae: Acidocerinae). Zootaxa 4232 (2): 271–281. <https://doi.org/10.11646/zootaxa.4232.2.10>
- Short AEZ, Hebauer F (2006) World catalogue of Hydrophiloidea (Coleoptera)—additions and corrections, 1 (1999–2005). Koleopterologische Rundschau 76: 315–359.
- Short AEZ, Torres PJ (2006) Review of the *Hemiosus* Sharp and *Derallus* Sharp of North America, Central America, and the Caribbean (Coleoptera: Hydrophilidae: Berosini). Zootaxa 1369: 1–17.
- Smetana A (1975) Revision of the New World genera of the tribe Omicrini trib. nov. of the hydrophilid subfamily Sphaeriidiinae (Coleoptera). Studies on Neotropical Fauna and Environment 10 (2): 153–182.
- Smetana A (1978) Revision of the subfamily Sphaeriidiinae of America north of Mexico (Coleoptera: Hydrophilidae). Memoirs of the Entomological Society of Canada, 110 (S105): 1–292.
- Smetana A (1984) Remarks on the Sphaeriidiinae genera *Pelosoma* Mulsant and *Deltostethus* Sharp (Coleoptera: Hydrophilidae). The Coleopterists Bulletin 38 (2): 192–200.
- Solier AJJ (1834) Observations sur la tribu des Hydrophiliens, et principalement sur le genre *Hydrophilus* de Fabricius. Annales de la Société entomologique de France 3: 299–318.
- Spangler PJ (1981) Supplement to the aquatic and semiaquatic Coleoptera of Cuba collected by the biospeleological expeditions to Cuba by the Academies of Science of Cuba and Romania. In: Orghidan T, et al. (Eds) Résultats des Expéditions Biospéologiques Cubano-Roumaines à Cuba, Vol. 3. Academiei Republicii Socialiste România, Bucharest, 145–171.
- Winters FC (1926) Notes on the Hydrobiini (Coleoptera-Hydrophilidae) of boreal America. Pan-Pacific Entomologist 3: 49–58.
- Winters FC (1927) Key to the subtribe Helochara: Orchym. (Coleoptera-Hydrophilidae) of boreal America. Pan-Pacific Entomologist 4: 19–29.
- Wooldridge DP (1973) New *Paracymus* from South America (Coleoptera: Hydrophilidae). Journal of the Kansas Entomological Society 46 (1): 116–123.
- Young FN (1954) The water beetles of Florida. University of Florida Studies, Biological Science Series 5(1): ix + 238 pp.
- Young FN (1967) Studies on the color patterns of crosses of *Tropisternus* from western Mexico with other forms of the *Tropisternus collaris* Complex. Proceedings of the Indiana Academy of Science 76: 272–278.
- Zaitzev FA (1908) Catalogue des coléoptères aquatiques des familles Dryopidae, Georyssidae, Cyathoceridae, Heteroceridae et Hydrophilidae. Horae Societatis entomologicae Rossiae 38: 283–420.
- Zimmermann C (1869) Synonymical notes on Coleoptera of the United States, with descriptions of new species, from the MSS. of the late Dr C. Zimmermann. (LeConte JL, Ed.) Transactions of the American entomological Society 2: 243–259.