



Recent occurrence of *Tinamus solitarius* (Vieillot, 1819) (Tinamiformes, Tinamidae) in central Rio Grande do Sul state, southern Brazil

Ivan Luiz Brondani,^{1,3} Everton Rodolfo Behr,² Veronica Pereira Damasceno,³ Marilise Mendonça Krügel⁴

1 Universidade Federal de Santa Maria, Departamento de Zootecnia, Av. Roraima, nº 1000, CEP 97105-900, Santa Maria, RS, Brazil.

2 Universidade Federal de Santa Maria, Departamento de Zootecnia e Programa de Pós-graduação em Biodiversidade Animal, CEP 97105-900, Santa Maria, RS, Brazil. **3** Universidade Federal de Santa Maria, Programa de Pós-graduação em Zootecnia, CEP 97105-900, Santa Maria, RS, Brazil.

4 Universidade Federal de Santa Maria, Departamento de Engenharia Sanitária e Ambiental, Av. Roraima, nº 1000, CEP 97105-900, Santa Maria, RS, Brazil.

Corresponding author: Ivan Luiz Brondani, ivanbrondani@gmail.com

Abstract

Tinamus solitarius (Vieillot, 1819), the Solitary Tinamou, was observed in the central region of Rio Grande do Sul state, on the western portion of the Serra Geral escarpment, in the municipality of Itaara. This population is the most austral of the species and is isolated from the 2 other populations that inhabit Rio Grande do Sul: 1 in Turvo State Park and the other in the eastern portion of the Serra Geral escarpment.

Key words

Solitary Tinamou; new record; Itaara; Serra Geral escarpment; threatened species.

Academic editor: Rafael Antunes Dias | Received 3 July 2017 | Accepted 10 March 2018 | Published 11 May 2018

Citation: Brondani IL, Behr ER, Damasceno VP, Krügel MM (2018) Recent occurrence of *Tinamus solitarius* (Vieillot, 1819) (Tinamiformes, Tinamidae) in central Rio Grande do Sul state, southern Brazil. Check List 14 (3): 505–508. <https://doi.org/10.15560/14.3.505>

Introduction

Tinamus solitarius (Vieillot, 1819), the Solitary Tinamou, is a large (length 52 cm, weight 2 kg), terrestrial game bird with solitary habits. It was formerly distributed throughout the entire Atlantic Forest region; however, today it is mainly confined to the largest habitat remnants within its original range and is extirpated in some places (del Hoyo et al. 1992, Sick 1997, Straube et al. 2004).

The species is globally (IUCN 2016) and nationally (ICMBio 2014) classified as Near Threatened. In several Brazilian states of the Atlantic Forest domain it is currently a threatened species: Critically Endangered in

Espírito Santo (Simon et al. 2007), Endangered in Minas Gerais (Copam 2010) and Rio de Janeiro (Alves et al. 2000), and Vulnerable in Paraná and São Paulo (Straube et al. 2004, Silveira et al. 2009). In Rio Grande do Sul, it is currently classified as Vulnerable (FZB 2014) but had been listed as Critically Endangered in a previous assessment (Marques et al. 2002, Bencke et al. 2003).

At the end of the nineteenth century, *T. solitarius* was commonly found in the forests of the eastern section of the escarpment in northeastern Rio Grande do Sul (Ihering 1899). However, in the early 1980s, this species had almost completely disappeared from that state (Belton 2003). In the more recent past, it was thought




Figure 1. Map showing current (●) and historical (▲) records of *Tinamus solitarius* in Rio Grande do Sul state, Brazil.

to be restricted to a few large reserves in northern Rio Grande do Sul (e.g., Turvo State Park) and to the remaining forests along the escarpment between São Francisco de Paula and Canela (Belton 2003, Bencke et al. 2003). Other historical records include specimens collected at Taquara and Marcelino Ramos over 70 years ago (Berlepsch and Ihering 1885, Gliesch 1930), as well as birds reported to have been hunted around Santa Cruz do Sul in the 1940s and 1950s or claimed to have been recorded near Agudo in the 1990s (Bencke 1997) (Fig. 1).

Recent records are from the eastern portion of the escarpment in the Pró-Mata Nature Research and Conservation Center, Serra Geral State Biological Reserve and Aratinga State Ecological Station (Bencke and Kindel 1999, Bencke et al. 2003, Fontana et al. 2000). Other photographic and audio records are available on online databases such as the Xeno-canto, eBird, and the Macaulay Library at the Cornell Lab of Ornithology. All these are from Maquiné (<http://www.xeno-canto.org/XC353537> and <http://www.xeno-canto.org/XC353525>) and São Francisco de Paula (<http://www.ebird.org/List/S39838626e>; <http://www.macaulaylibrary.org/ML71573911>), in areas already covered by the known records.

Breeding of *T. solitarius* in Rio Grande do Sul was documented in Turvo State Park in October 2003 (Maurício et al. 2013). Eggs were also found at Serra Geral State Biological Reserve in Maquiné (December 2004) and at Osório (January 2007) (Maurício et al. 2013).

Based on this information, we report a new locality for *T. solitarius* in central Rio Grande do Sul, which confirms the continued existence of this species at the southern edge of its historical range.

Methods

Sporadic expeditions were carried out to areas of the escarpment in central Rio Grande do Sul encompassing the municipalities of Santa Maria and Itaara. Reports by the local community about the presence of *T. solitarius* motivated the accomplishment of periodic expeditions aimed at finding and documenting the species in this region. The study area (centered at 29°33'49" S, 053°43'46" W) is in private properties in the municipality of Itaara, on the Serra Geral escarpment. The original vegetation is semideciduous forest. For the visual and photographic records, we used a fixed



Figure 2. Photographic record of the *Tinamus solitarius* in the municipality of Itaara, Rio Grande do Sul State, Brazil (<http://www.wikiaves.com.br/WA2227823>).

blind installed inside the forest to allow the view of 2 nature trails, upstream and downstream. The observations occurred twice a week for 2 months (April and May 2014), between 2:00 pm and 04:45 pm, and the observer remained sitting in the blind with a 55–300 mm zoom camera. After this first period of observations, efforts continued in the study area sporadically.

Results

New records. Brazil: Rio Grande do Sul: Itaara: near the Arroio Grande stream (29°33'50.47" S, 053°43'48.00" W), Ivan Luiz Brondani, 9 May 2014 (first record); Brazil: Rio Grande do Sul: Itaara: near the Arroio Grande stream (29°33'22.17" S, 053°42'26.36" W), Ivan Luiz Brondani, 3 August 2016 (Fig. 2); Brazil: Rio Grande do Sul: Itaara: near the Arroio Grande stream (29°33'22.17" S, 053°42'26.36" W), Ivan Luiz Brondani, 8 August 2016 (last record).

Identification. On all occasions, birds were identified based on their large size, ochre line along side of the neck, and olivaceous-gray back, barred with dark color (del Hoyo 1992). Size was the main characteristic used to distinguish the species from other species of Tinamidae

present in the study area (e.g., *Crypturellus obsoletus* (Temminck, 1815) and *Crypturellus tataupa* (Temminck, 1815) personal observation).

Discussion

Specimens were seen in a private property in the forested valley of the Arroio Grande stream, which is a tributary of the Vacacaí-Mirim River. The municipality of Itaara is extensively covered with native forests. The mapping of permanent preservation areas revealed their importance, since forests correspond to 43.2% of the whole territory (Fensterseifer et al. 2014). The highly rugged relief possibly helped prevent these forests from being removed in the past and the large size of some remnants may be contributing to the maintenance of this population of *T. solitarius*.

Historical records of *T. solitarius* along the escarpment extended west to Santa Cruz do Sul and possibly Agudo (Bencke 1997, Belton 2003). Itaara is located more or less at the same latitude and about 50 km to the west of these locations and currently represents the extreme southern limit of this species' geographical distribution. The persistence of isolated populations of *T. solitarius*

in central Rio Grande do Sul was suggested by Bencke (1997). This population is isolated from the 2 other existing populations of Rio Grande do Sul, 1 inhabiting Turvo State Park and the other in the eastern portion of the Serra Geral escarpment, between the São Francisco de Paula National Forest and the Serra Geral National Park.

It is worth noting that an isolated population of another threatened tinamid, *Crypturellus noctivagus noctivagus* (Wied, 1820), which was thought extinct in the state (Bencke et al. 2003), was also discovered recently in central Rio Grande do Sul (municipality of São Sepé; Corrêa et al. 2010). Historical data suggest that this species, like *T. solitarius*, occurred along the Serra Geral escarpment from Torres to Santa Maria (Belton 2003). These recent findings make the central region of Rio Grande do Sul an important area for research and survey of these and other species of conservation concern.

As *T. solitarius* is a large and relatively well-known species in rural areas, it would be of great interest to use it in environmental education programs for forest conservation and to prevent hunting as suggested by Straube et al. (2004). The conservation plan for *T. solitarius* in Paraná assigns high priority and crucial importance to the creation of protected areas where the species has significant populations (IAP 2009). Our new data of *T. solitarius* on the western escarpment of the Serra Geral indicate the need for continued monitoring of the species as well as for the establishment of public and/or private reserves for its protection.

Acknowledgements

We thank Bruna Biassi and Alberto Senra Gonçalves for helping with the map, and two anonymous reviewers and the academic editor for improving the manuscript.

Authors' Contributions

ILB collected the data, ERB, VPD and MMK wrote the manuscript.

References

- Alves MAS, Pacheco JF, Gonzaga LAP, Cavalcanti RB, Raposo MA, Yamashita C, Maciel NC, Castanheira M (2000) Aves. In: Bergallo HG, Rocha CFD, Alves MAS, van Sluys M (Eds) A fauna ameaçada de extinção do Estado do Rio de Janeiro. UERJ, Rio de Janeiro, 113–124.
- Belton W (2003) Aves do Rio Grande do Sul: Distribuição e Biologia. UNISINOS, São Leopoldo, 584 pp.
- Bencke GA (1997) Sobre a coleção de aves do Museu do Colégio Mauá, Santa Cruz do Sul (RS). *Biociências* 5 (1): 143–164.
- Bencke GA, Fontana CS, Dias RD, Maurício GN, Mähler Jr JKF (2003) Aves. In: Fontana CS, Bencke GA, Reis RE (Eds) Livro Vermelho da Fauna Ameaçada de Extinção no Rio Grande do Sul. EDIPUCRS, Porto Alegre, 189–479.
- Bencke GA, Kindel A (1999) Bird counts along an altitudinal gradient of Atlantic forest in northeastern Rio Grande do Sul, Brazil. *Ararajuba* 7 (2): 91–107.
- Berlepsch H von, Ihering H von (1885) Die Vögel der Umgegend von Taquara do Mundo Novo, Prov. Rio Grande do Sul. *Zeitschrift für die gesammte Ornithologie* 2: 1–88.
- Conselho de Política Ambiental. Deliberação Normativa COPAM nº 147, de 30 de abril de 2010. Aprova a Lista de Espécies Ameaçadas de Extinção da Fauna do Estado de Minas Gerais. Diário do Executivo do Estado de Minas Gerais, Belo Horizonte, 04 maio 2010. 18 pp.
- Corrêa LLC, Silva DE, Cappellari LH (2010) Aves, Tinamidae, *Crypturellus noctivagus noctivagus* (Wied, 1820): southward range extension and rediscovery in Rio Grande do Sul, Brazil. *Check List* 6 (4): 585–586. <https://doi.org/10.15560/6.4.585>
- Del Hoyo J, Elliott A, Sargatal J (Eds) (1992) Handbook of the Birds of the World: Ostrich to Ducks. Lynx Edicions, Barcelona, 696 pp.
- Fensterseifer SLB, Gonçalves IHP, Donadel LP (2014) Mapeamento de áreas de preservação permanente no município de Itaara – RS pelo Spring 5.1.6. *Geografia* 23 (1): 5–19.
- Fontana CS, Mähler Jr JKF, Joenck CM, Mendonça-Lima A de (2000) Lista comentada da avifauna do Centro de Pesquisas e Conservação da Natureza Pró-Mata (CPCN), São Francisco de Paula, Rio Grande do Sul, VIII Congresso Brasileiro de Ornitologia. Universidade do Sul de Santa Catarina, Sociedade Brasileira de Ornitologia, Curitiba, 266–267.
- FZB (2014) Lista vermelha das espécies da fauna ameaçadas de extinção no Rio Grande do Sul. http://www.fzb.rs.gov.br/conteudo/2403/?Informações_da_Biodiversidade. Accessed on 2016-12-15.
- Gliesch R (1930) Lista das aves colligidas e observadas no Estado do Rio Grande do Sul. *Egatea* 15: 276–292.
- IAP (2009) Planos de conservação para aves e mamíferos ameaçados do Paraná—Planos completos. IAP/Projetos Paraná Biodiversidade, Curitiba, 72 pp.
- ICMBio. Lista das espécies consideradas quase ameaçadas (NT). <http://www.icmbio.gov.br/portal/faunabrasileira/lista-de-especies-dados-insuficientes>. Accessed on 2018-02-15.
- IUCN. 2016. The IUCN Red List of Threatened Species. Version 2017.3. International Union for Conservation of Nature. <http://www.iucnredlist.org>. Accessed on 2018-01-15.
- Ihering H von (1899) As aves do Estado do Rio Grande do Sul. In: Anuário do Estado do Rio Grande do Sul para o ano de 1900, Porto Alegre, 113–154.
- Marques AAB, Fontana CS, Vélez E, Bencke GA, Schneider M, Reis RE dos (Orgs) (2002) Lista das espécies da fauna ameaçadas de extinção no Rio Grande do Sul. Publicações Avulsas, FZB, 11. FZB/MCT–PUCRS/PANGEA, Porto Alegre, 52 pp.
- Maurício GN, Glayson AB, Repenning M, Machado DB, Dias RA, Bugoni L (2013) Review of the breeding status of birds in Rio Grande do Sul, Brazil. *Iheringia* 103 (2): 163–184.
- Silveira LF, Benedicto GA, Schunck F, Sugieda AM (2009) Aves. In: Bressan PM, Kierulff MCM, Sugieda AM (Coords) Livro vermelho da fauna ameaçada de extinção no Estado de São Paulo: Vertebrados. Fundação Parque Zoológico de São Paulo: Secretaria do Meio Ambiente, São Paulo, 87–283.
- Simon JE, Antas PTZ, Pacheco JF, Efe MA, Ribon R, Raposo MA, Laps RR, Musso C, Passamani JA Pacagnella SG (2007) As aves ameaçadas de extinção no Estado do Espírito Santo. In: Passamani M, Mendes SL (Eds) Espécies da fauna ameaçadas de extinção no Estado do Espírito Santo. Instituto de Pesquisas da Mata Atlântica, Vitória, 46–64.
- Straube FC, Urben-Filho A, Kajiwarra D (2004) Aves. In: Mikich SB, Bérnils RS (Eds) Livro Vermelho da Fauna Ameaçada de Extinção no Estado do Paraná. Instituto Ambiental do Paraná, Curitiba, 146–496.
- Sick H (1997) Ornitologia Brasileira. Editora Nova Fronteira, Rio de Janeiro, 912 pp.