



The first country record of the Mexican Blind Snake, *Anomalepis mexicanus* Jan, 1860 (Serpentes, Anomalepididae), in Colombia

Juliett M. González-Carvajal^{1,2}, Guido Fabian Medina-Rangel^{3,4}, Luis Eduardo Rojas-Murcia^{1,2,4}

1 Universidad Pedagógica y Tecnológica de Colombia. Avenida Central del Norte 39-115, 150003 Tunja, Tunja, Boyacá. **2** Laboratório de Biologia de Vertebrados, Faculdade de Ciências, Universidade Estadual do Centro-Oeste, Paraná, Brazil. Rua Salvatore Renna, 875 - Santa Cruz, Guarapuava - PR, 85015-430, Brazil. **3** Grupo de Biodiversidad y Sistemática Molecular, Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Ciudad Universitaria, Edificio 425, Bogotá D.C., Colombia. **4** Grupo de Biodiversidad y Conservación, Reptiles, Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Ciudad Universitaria, Edificio 425, Bogotá D.C., Colombia.

Corresponding author: Guido Fabian Medina-Rangel, guidofabianmedina@gmail.com

Abstract

We report the first Colombian record of the Blind Snake *Anomalepis mexicanus* Jan, 1860, based on a single specimen from Cantagallo municipality, department of Bolívar. Our new record fills a large gap and extends this species' geographical distribution by approximately 630 km in a straight line east-southeast from its nearest previously known Panamanian locality, and approximately 1295 km in a straight-line north-northeast from its nearest previously known Peruvian locality.

Key words

Cantagallo; department of Bolívar; distribution; dry forest; range extension; Serranía de San Lucas; South America.

Academic editor: Sebastian Lotzkat | Received 11 July 2018 | Accepted 31 October 2018 | Published 9 November 2018

Citation: González J, Medina-Rangel GF, Rojas-Murcia LE (2018) The first country record of the Mexican Blind Snake, *Anomalepis mexicanus* Jan, 1860 (Serpentes, Anomalepididae), in Colombia. Check List 14 (6): 1047–1052. <http://doi.org/10.15560/14.6.1047>

Introduction

Belonging to the so-called blind snakes (Scolecophidia), the Neotropical snake family Anomalepididae is composed of 3 genera with 18 species, of which 8 are found in Colombia (Uetz et al. 2018). The anomalepidid genus *Anomalepis* Jan, 1860, is currently contains 4 species: *Anomalepis aspinosus* Taylor, 1939, *A. colombia* Marx, 1953, *A. flavapices* Peters, 1957, and *A. mexicanus* Jan, 1860. All these species are fossorial, adapted for burrowing, and most have been recorded below 600 m above sea level in moist and dry forest of lowlands and foothills. Only *A. colombia* has been recorded between 1000 and

1700 m above sea level (Marx 1953, Arroyo-Sánchez et al. 2016). In Colombia, until now, only 1 species has been reported: *A. colombia*, which is currently believed to be endemic to the country (Peters 1957, Peters and Orejas-Miranda 1970, Kofron 1988, Pérez-Santos and Moreno 1988, Arroyo-Sánchez et al. 2016, Uetz et al. 2018).

Three or fewer specimens each of *A. aspinosus*, *A. colombia*, and *A. flavapices* are currently known in collections (Dunn 1941, Peters 1957, Kofron 1988, Arroyo-Sánchez et al. 2016, Cisneros-Heredia 2016, GBIF 2018). In contrast, 23 specimens of *A. mexicanus* specimens have been reported in biological collections (Appendix 1) to date, collected at about 10 different loca-




Figure 1. Distribution map for the species of the genus *Anomalepis*. Green circles: *A. aspinosus* (Taylor 1939, Dunn 1941, Kofron 1988). Yellow circles: *A. colombia* (Marx 1953, Arroyo-Sánchez et al. 2016). Blue circles: *A. flavapices* (Peters 1957, Kofron 1988). Red circles: *A. mexicanus* in Central America and Peru (Dunn 1941, Kofron 1988). Red star: new record of *A. mexicanus* was found at Cantagallo municipality, department of Bolívar, Colombia.

tions in 5 countries. *Anomalepis mexicanus* has 2 disjunct populations, 1 distributed from the west of Honduras to the center of Panama below 500 m above sea level, and another east of the Peruvian Andes in the Amazonas department below 200 m above sea level (Fig. 1; Dunn 1941, Taylor 1951, Kofron 1988, Savage 2002, Köhler et al. 2004, Wilson and Townsend 2006, Batista et al. 2013, Sunyer 2014). This species has been found in lowland rainforests and foothill moist forests from under rotten logs and in leaf litter (McCranie and Wilson 2002). *Anomalepis mexicanus* has been assessed as Data Deficient (DD) for the IUCN Red List of Threatened Species (Batista et al. 2013) because, although its documented range is relatively broad, it is rarely found and little is known about its full geographical distribution, abundance, and ecology. Moreover, there is uncertainty about the threats it faces. Here, we report the first confirmed record of the Mexican Blind Snake, *A. mexicanus*, from Colombia.

Methods

A field survey, by means of active searching, was carried out in the Cimitarra River Valley Rural Reserve Zone (ZRC-VRC) in Cantagallo municipality, department of Bolívar, Colombia. The specimen was euthanized in 2% Roxicaine, fixed in 10% formalin solution, and preserved in 70% ethanol. We identified specimens of *A. mexicanus*

by comparison to the original description of the species (Jan 1860) and the latest revision of the genus (Kofron 1988). The specimen's identification was verified by J.E. Carvajal-Cogollo. The specimen was deposited at the reptile collection of the Instituto de Ciencias Naturales, Universidad Nacional de Colombia, Bogotá, Colombia (ICN-R), under the catalogue number ICN-R 13040, and collected under the permits #524 of 17 October 2017 and #664 of 7 December 2017 of the Asociación Campesina del Valle del río Cimitarra, Antioquia-Bolívar-Santander departments, Colombia. Scale counts follow the methodology established by Kofron (1988).

Head scale photos were taken with a Leica M205 C stereo microscope with Fusion Optics type. We obtained the geographic coordinates of the capture site using a Garmin Map 62S GPS receiver and locality records for other species of *Anomalepis* (Appendix, Table A1) were obtained from Kofron (1988), Sunyer (2014), Arroyo-Sánchez et al. (2016), Serrano et al. (2016), and GBIF 2018. In cases where the coordinates of the specimen were not explicitly provided, we inferred them using the closest point to the referred locality by searching it in Google Earth™. Institutional acronyms for museum collections follow those of Sabaj-Perez (2016), except for that of the herpetological reference collection of the Universidad del Quindío (ARUQ). The map was made using the open source software QGIS v. 2.18.4 (QGIS Development Team 2017).



Figure 2. Adult female of *Anomalepis mexicanus* (ICN-R 13040) from Vereda Yanacué, Colombia, in life. Photo JG.

Results

New record. Colombia: Bolívar department: Cantagallo municipality: Vereda Yanacué: Cimitarra River Valley Rural Reserve Zone (07°17.23' N, 074°01.78' W, ca 205 m a.s.l.; Fig. 1), collector Juliatt González, 18 October 2016, 1 individual (ICN-R 13040; Fig. 2).

The single specimen was found at noon under a fallen log during the rainy season. The forest fragment (Fig. 3A, B) had been burnt and was undergoing regeneration.

Identification. *Anomalepis mexicanus* has ventral and dorsal scales equal in size (Fig. 2), a pair of polygonal prefrontals in contact on midline behind rostral, followed by pentagonal frontal (Fig. 4A), and the eye scarcely visible under a scale (Fig. 4B). Apart from matching these characteristics, our specimen's low number of dorsal scales (265) and its scale-row formula of 24 / 22 / 20 are within the range expected for the species and distinguish it from the other 3 species of *Anomalepis* (Table 1; Dunn 1941, Taylor 1939, Marx 1953, Peters 1957, Peters and Orejas-Miranda 1970, Kofron 1988, Pérez-Santos and Moreno 1988).

Discussion

The distribution and even more the natural history of *Anomalepis* species, because of their secretive habits and small size, are poorly known (Dunn 1941, Peters 1957, Kofron 1988). The most recent review of the genus was 30 years ago (Kofron 1988) and included the different populations of *A. mexicanus* from Central America and Peru. *Anomalepis mexicanus* is an uncommon fossorial and seldom seen snake and the type species of the genus *Anomalepis*. It was first described from 1 specimen; however, the type locality ("Mexico") was in error, and the specimen is now believed lost (Kofron 1988).




Figure 3. Forest fragment in regeneration where the first Colombian specimen *Anomalepis mexicanus* was found. **A.** Overview. **B.** Detail of the interior. Photos: JG.

Our record is the first of *A. mexicanus* for Colombia and extends its known geographic distribution by approximately 630 km in a straight line southeast from its nearest previously known locality in Panama, Corregimiento de Cristóbal, Barro Colorado Island (Kofron 1988). From the nearest previously known Peruvian locality, vicinity of La Poza, Santiago river, Amazonas

Table 1. Selected morphological characters for the genus *Anomalepis*. Information was obtained from the following literature: *A. aspinosus* data from Taylor (1939), Dunn (1941), and Kofron (1988); *A. colombia* data from Marx (1953) and Kofron (1988); *A. flaviceps* data from Peters (1957) and Kofron (1988); *A. mexicanus* data from Taylor (1939), Dunn (1941), Kofron (1988), and this study.

Species	No. of specimens examined in the papers consulted	No. of dorsal scales around the body*	No. of scale rows
<i>A. aspinosus</i>	3	(27 or 28) / 24 / 24	329–342
<i>A. colombia</i>	1	30 / 30 / 27	363
<i>A. flaviceps</i>	2	26 / 26 / 24	308–317
<i>A. mexicanus</i> , Central America	9	(24 or 26) / (20 or 22 or 23) / (20 or 21 or 22)	258–274
<i>A. mexicanus</i> , new record (ICN-R 13040)	1	24 / 22 / 20	265
<i>A. mexicanus</i> , Peru	7	(24 or 25 or 26) / (22 or 23 or 24) / (21 or 22)	265–299

department, by straight-line distance 1295 km to the northeast (Peters 1957, Peters and Orejas-Miranda 1970, Kofron 1988).

This is a rarely found species of snake that can be severely affected by deforestation and habitat loss; its populations are fragmented and distribution records are scattered. However, insufficient data prevent any Red List categorization other than Data Deficient (DD; Batista et al. 2013). We highlight the importance of our new record, which opens the possibility that the geographic range of *A. mexicanus* is still slightly larger and may even occupy future regionally protected areas in Colombia south of the Serranía de San Lucas.

However, the Cis-Andean distribution of the Peruvian specimens and the separation of these populations from Central American populations, as well as the differences in some scale counts (Table 1; Taylor 1939, Kofron 1988), suggest the possibility that there are 2 distinct species involved. The specimen we collected has scale counts between the recorded values for both populations (Table 1). Additional material for morphological comparisons and tissue samples allowing for molecular analysis will be necessary to clarify the uncertainty about this species and its populations.

Acknowledgements

We thank the Asociación Campesina del Valle del río Cimitarra, Antioquia-Bolívar-Santander departments, Colombia, for their constant commitment and dedication in the conservation of its natural environment, funding the study of biological diversity under the mark its project “Evaluation biodiversity of the wetland complex stream Negro and river Cimitarra, in the Cimitarra River Valley Rural Reserve Zone, Colombia”. We appreciate the collaboration of our guides Melquiades and Helmer. We thank the communities of Buenos Aires, Cagüí, Yanacué, Trasmallo, and the La Poza in Cantagallo municipality (Bolívar). We thank the Wildlife Conservation Society, Fondo Acción ONG, Regional Autonomous Corporation of South Bolívar, Regional Autonomous Corporation of Central Antioquia, and finally, all the people and entities involved in the declaration of Serranía de San Lucas as a new regional protected area. We also thank the Laboratorio




Figure 4. Close-ups of the head of the preserved specimen ICN-R 13040 of *Anomalepis mexicanus*. **A.** Dorsal head scale details. **B.** Lateral head scale details. Photos: GFM.

de Equipos Ópticos Compartidos, Departamento de Biología, Universidad Nacional de Colombia–Bogotá for allowing us the use of equipment for photographing the specimen. We appreciate the valuable comments and suggestions made by reviewers Abel Batista, Javier Sunyer, Edgar Lehr, and Sebastian Lotzkat that contributed to improving the text.

Authors' Contributions

JG conducted the field survey. JG and GFM took the photographs. GFM wrote the text, made map. JG, GFM and LER made the identification, tables, and took the counts and measurements. All authors read and approved the final manuscript.

References

- Arroyo-Sánchez SB, Bernal-González VA, Quiñones-Betancur E, Viloria Rivas J (2016) Anfibios y Reptiles. In: Prieto-Cruz A, Rodríguez J, Aguirre J (Eds) Guía ilustrada del Proyecto Hidroeléctrico Ituango – Antioquia. Vol. II. Serie de Guías del Instituto de Ciencias Naturales de la Universidad Nacional de Colombia N. 18. Banco Interamericano de Desarrollo, Empresas Públicas de Medellín y Universidad Nacional de Colombia. Bogotá, 19–42.
- Batista A, Saborío G, Vargas-Álvarez J, Köhler G, García-Rodríguez A, Ballesteros E, Acosta-Chaves V (2013) *Anomalepis mexicanus*. The IUCN Red List of Threatened Species 2013: eT203193A2761875. <https://doi.org/10.2305/IUCN.UK.2013-2.RLTS.T203193A2761875.en>. Accessed on: 2017-9-7.
- Cisneros-Heredia DF (2016) *Anomalepis flavapices*. The IUCN Red List of Threatened Species 2016: eT203192A2761868. <https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T203192A2761868.en>. Accessed on: 2017-9-7.
- Dunn ER (1941) Notes on the snake genus *Anomalepis*. Bulletin of the Museum of Comparative Zoology 87 (7): 509–526.
- GBIF (2018) GBIF Occurrence Download. <https://doi.org/10.15468/dl.bly5s8>. Accessed on: 2017-9-7.
- Jan G (1860) Iconographie générale des ophiidiens. 1. Livraison. J.B. Baillière et Fils, Paris, Pl 5. <https://doi.org/10.5962/bhl.title.4885>
- Kofron C (1988) The central and south american blindsnakes of the genus *Anomalepis*. Amphibia-Reptilia 9: 7–14. <https://doi.org/10.1163/156853888X00152>
- Köhler G, Quintana AZ, Buitrago F, Diethert H (2004) New and noteworthy records of amphibians and reptiles from Nicaragua. Salamandra 40 (1): 15–24.
- Marx H (1953) A new worm snake from Colombia, genus *Anomalepis*. Fieldiana Zoology 34 (17): 197–198.
- McCrane RJ, Wilson LD (2002) The amphibians of Honduras. Society for the Study of Amphibians and Reptiles, Ithaca, NY, 625 pp.
- Pérez-Santos C, Moreno AG (1988) Ofidios de Colombia. Museo regionale di Scienze Naturali, Monographie 6: 1–517.
- Peters JA (1957) Taxonomic notes on Ecuadorian snakes in the American Museum of Natural History. American Museum Novitates 1851: 1–13.
- Peters JA, Orejas-Miranda B (1970) Catalogue of the Neotropical Squamata. Part I. Snakes. United States National Museum Bulletin 297: 1–347. <https://doi.org/10.5479/si.03629236.297.1>
- QGIS Development Team (2017) QGIS Geographic Information System, version 2.18.4. Open Source Geospatial Foundation. <http://qgis.osgeo.org>.
- Sabaj-Perez MH (2016) Standard Symbolic Codes for Institutional Resource Collections in Herpetology and Ichthyology: An Online Reference. Version 6.5. American Society of Ichthyologists and Herpetologists, Washington, DC. <http://www.asih.org/standard-symbolic-codes>. Accessed on: 2018-2-15.
- Savage JM (2002) The Amphibians and Reptiles of Costa Rica: A Herpetofauna Between Two Continents, Between Two Seas. University of Chicago Press, Chicago and London, 934 pp.
- Serrano H, Rudas A, Prieto A, Arias-G. JC (2016) Componente geográfico. In: Prieto-Cruz A, Rodríguez J, Aguirre J (Eds) Guía ilustrada del Proyecto Hidroeléctrico Ituango - Antioquia. Vol. I. Geografía, vegetación y flora. Serie de Guías del Instituto de Ciencias Naturales de la Universidad Nacional de Colombia N. 18. Banco Interamericano de Desarrollo, Empresas Públicas de Medellín y Universidad Nacional de Colombia, Bogotá, 19–40.
- Sunyer J (2014) An updated checklist of the amphibians and reptiles of Nicaragua. Mesoamerican Herpetology 1 (2): 186–202.
- Taylor EH (1939) Two new species of the genus *Anomalepis* Jan, with a proposal of a new family of snakes. Proceedings of the New England Zoological Club 17: 87–96.
- Taylor EH (1951) A brief review to the snakes of Costa Rica. Kansas University Science Bulletin 34 (1): 3–188.
- Uetz P, Freed P, Hošek J (Eds) (2018) The Reptile Database. <http://www.reptile-database.org>. Accessed on: 2018-2-7.
- Wilson LD, Townsend JH (2006) The herpetofauna of the rainforests of Honduras. Caribbean Journal of Science 42 (1): 88–113.

Appendix

Table A1. Locality records for species of the genus *Anomalepis*. The geodetic datum is WGS84.

Species	Specimen voucher	Country	Province/ Department	Locality	Latitude	Longitude	Source
<i>A. aspinosus</i>	LSUMZ 32591	Peru	Amazonas	22 km SE Bagua Grande	05°53.83'S	78°18.07'W	Kofron (1988)
<i>A. aspinosus</i>	MCZ 14781–85, 17401–03, USNM 76295	Peru	Cajamarca	Perico	05°20.63'S	78°47.68'W	GBIF (2018), Kofron (1988)
<i>A. aspinosus</i>	CM 90254, LSUMZ 19450–54	Peru	Cajamarca	Perico, 28 km N of Santa Cruz	06°21.32'S	78°55.61'W	GBIF (2018), Kofron (1988)
<i>A. colombia</i>	In process of being deposited at the ICN-R collection	Colombia	Antioquia	Toledo municipality, corregimiento El Valle, Tacuí Cuní	07°02.90'N	75°41.90'W	Arroyo-Sánchez et al. (2016), Serrano et al. (2016)
<i>A. colombia</i>	FMNH 54986	Colombia	Caldas	Pueblo Rico municipality, La Selva	05°07.88'N	75°31.59'W	Kofron (1988)
<i>A. colombia</i>	ARUQ-106	Colombia	Quindío	Armenia municipality, Universidad de Quindío campus, Botanical Garden	04°33.18'N	75°39.79'W	GBIF (2018)
<i>A. flavapices</i>	USNM 196349	Ecuador	Esmeraldas	near Esmeraldas	00°58.99'N	79°42.00'W	Kofron (1988)
<i>A. flavapices</i>	AMNH R-6966	Ecuador	Manabí	No data	00°56.06'S	80°22.64'W	Kofron (1988)
<i>A. mexicanus</i>	ZFMK 57773	Costa Rica	Guanacaste	Guanacaste National Park, Quebrada Zompota, south of Estación Biológica Maritza	10°57.41'N	85°29.69'W	GBIF (2018)
<i>A. mexicanus</i>	KU 35512	Costa Rica	Guanacaste	4 km ESE Los Ángeles de Tilarán	10°28.47'N	85°02.99'W	Kofron (1988)
<i>A. mexicanus</i>	UMMZ 132190	Costa Rica	Guanacaste	Highway 19, 5,5 km W Tilarán	10°26.23'N	84°56.88'W	Kofron (1988)
<i>A. mexicanus</i>	USNM 565512	Honduras	Gracias a Dios	Cañones Cay	15°49.20'N	84°39.00'W	GBIF (2018)
<i>A. mexicanus</i>	USNM 563482–83	Honduras	Gracias a Dios	San Hil	14°56.56'N	84°40.45'W	GBIF (2018)
<i>A. mexicanus</i>	USNM 562870, 564062–63	Honduras	Gracias a Dios	Warunga Tingni Kiamip	14°56.56'N	84°40.46'W	GBIF (2018)

Table A1. *Continued.*

Species	Specimen voucher	Country	Province/ Department	Locality	Latitude	Longitude	Source
<i>A. mexicanus</i>	UF 141964–65	Honduras	Gracias a Dios	Urus Tingni Kiangp	14°55.00'N	84°40.99'W	GBIF (2018)
<i>A. mexicanus</i>	SMF 82845	Nicaragua	Atlántico Norte	Parque Nacional Saslaya	13°55.39'N	84°51.82'W	Sunyer (2014)
<i>A. mexicanus</i>	AMNH R-119069	Panama	Bocas del Toro	No data	09°22.65'N	82°31.43'W	GBIF (2018)
<i>A. mexicanus</i>	MCZ 191201	Panama	Panamá	Canal Zone, Gatún	09°12.41'N	79°52.04'W	GBIF (2018)
<i>A. mexicanus</i>	FMNH 8224, 22853, 94731, MCZ 29220, KU 75663, 116883	Panama	Panamá	Corregimiento de Cristóbal, Canal Zone, Barro Colorado Island	09°09.29'N	79°50.73'W	Kofron (1988)
<i>A. mexicanus</i>	AMNH R-103750	Panama	Panamá	No data	08°59.96'N	79°33.23'W	GBIF (2018)
<i>A. mexicanus</i>	MVZ 174999	Peru	Amazonas	vicinity of La Poza, Santiago river	04°01.32'S	77°45.06'W	Kofron (1988)
<i>A. mexicanus</i>	MVZ 163245	Peru	Amazonas	vicinity of Huampami (Aguaruna village), Cenepa river	04°27.34'S	78°09.67'W	Kofron (1988)
<i>A. mexicanus</i> , new record	ICN-R 13040	Colombia	Bolívar	Cantagallo municipality, Vereda Yanacué	07°17.23'N	73°59.74'W	This study