



Panstrongylus diasi Pinto & Lent, 1946 (Hemiptera, Reduviidae, Triatominae): first record in Mato Grosso, Brazil

Mirian Francisca Martins^{1,2}, Sinara Cristina de Moraes¹, Marcos Takashi Obara³, Cleber Galvão⁴

1 Departamento de Vigilância em Saúde Ambiental, Secretaria de Estado de Saúde de Mato Grosso, Amaro Leite, 474, Barra do Garças, MT, 78600-000 Brazil. **2** Programa de Pós-Graduação em Biodiversidade e Saúde, Instituto Oswaldo Cruz, FIOCRUZ, Av. Brasil 4365, Rio de Janeiro, RJ, 21040-360 Brazil. **3** Departamento de Saúde Coletiva, Universidade de Brasília, Campus Ceilândia, QNN 14, Área Especial Ceilândia Sul, Brasília, DF, 72220-140 Brazil. **4** Laboratório Nacional e Internacional de Referência em Taxonomia de Triatomíneos, Instituto Oswaldo Cruz, FIOCRUZ Av. Brasil 4365, Pavilhão Rocha Lima, sala 505, Rio de Janeiro, RJ, 21040-360 Brazil.

Corresponding author: Mirian Francisca Martins, miranfranmartins@gmail.com

Abstract

Panstrongylus diasi Pinto & Lent, 1946, previously known from throughout the Brazilian center-west, is reported for the first time in Mato Grosso state, Brazil. We found 3 specimens, 1 of them from an urban area of Barra do Garças. The new record of *P. diasi* is important for the understanding of the epidemiology of Chagas disease, mainly because this species is found naturally infected with *Trypanosoma cruzi* (Chagas, 1909). Studies on the ecology, biogeography, and vector–host/parasite interactions of *P. diasi* are essential.

Key words

Chagas disease, ecology, new records, surveillance programs, *Trypanosoma cruzi*.

Academic editor: Hélcio R. Gil-Santana | Received 26 October 2018 | Accepted 30 November 2018 | Published 1 March 2019

Citation: Martins MF, Moraes SC, Obara MT, Galvão C (2018) *Panstrongylus diasi* Pinto & Lent, 1946 (Hemiptera, Reduviidae, Triatominae): first record in Mato Grosso, Brazil. Check List 15 (2): 259–264. <https://doi.org/10.15560/15.2.259>

Introduction

The blood-sucking insects of the subfamily Triatominae are vectors of Chagas disease, an infection caused by the protozoan *Trypanosoma cruzi* (Chagas, 1909). The subfamily is composed of 155 described species, assigned to 5 tribes and 19 genera (Souza et al. 2016, Justi and Galvão 2017, Rosa et al. 2017, Poinar-Jr 2019). The genus *Panstrongylus* Berg, 1879 was described based on *Panstrongylus guentheri* Berg, 1879, and has 15 species, 9 recorded in Brazil. In the state of Mato Grosso 4 species of this genus have been recorded: *P. megistus* (Burmeister, 1835), *P. geniculatus* (Latreille, 1811), *P. rufotuberculatus* (Champion, 1899), and *P. lignarius* (Walker, 1873) (Paula et al. 2013, Galvão 2014).

The genus *Panstrongylus* is important for public health because some species are associated with the transmission of *T. cruzi* to humans and other mammals. (Santos et al. 2003, Patterson et al. 2009). According to Maeda et al. (2012), among the 5 most collected triatomine species in the Federal District of Brazil, 3 belong to the genus *Panstrongylus*: *P. megistus*, *P. geniculatus*, and *P. diasi*.

Panstrongylus diasi has been reported in Brazil in the states of Bahia, Espírito Santo, Goiás, Maranhão, Mato Grosso do Sul, Minas Gerais, Rio Grande do Norte, São Paulo, and Tocantins and in the Distrito Federal (Sherlock and Serafim 1972, Rêbelo et al. 1998, Marassá and Barata 2000, Santos et al. 2004, Costa et al. 2003, Oliveira and Silva 2007, Almeida et al. 2008, Mendes et




Figure 1. Municipality of Barra do Garças, Mato Grosso, Brazil. Source: "Instituto Brasileiro de Geografia e Estatística".

al. 2008, Maeda et al. 2012, Gil-Santana et al. 2014, Barreto et al. 2017). Here, we report the first record of this species in Mato Grosso.

Methods

Barra do Garças is a municipality, in Mato Grosso, with a population of over 56,000 inhabitants and a territorial area of 9079 km² (IBGE 2017). It is located on the banks of the Garças and Araguaia rivers and is an important area for tourism. The municipality is within the Cerrado biome, which is characterized by spatial heterogeneity and high endemism of plants and insects (Silva and Bates 2002, MMA 2002); the biome is a biodiversity hotspot. The climate is characterized by having 2 seasons (dry winter and rainy summer), corresponding to type Aw according to the Köppen classification (Köppen 1948); the average annual humidity is 60% and the annual averages of temperature range from 20 °C to 27 °C.

In 2011, during the routine work by the Chagas Disease Control Program of Environmental Health Surveillance, 2 specimens of triatomines were collected by hand by agents of endemic diseases in the rural settlements of Pontal do Araguaia, Mato Grosso, Brazil. The materials used were tweezers and lanterns to inspect shaded places. In 2017, a resident collected a third specimen, this time inside a dwelling in Barra do Garças (Fig. 1).

The 3 female specimens were studied at the Entomology Laboratory of the Escritório Regional de Saúde de Barra do Garças (ERSBG) of the Secretaria de Estado de Saúde de Mato Grosso, Brazil (SESMT), and identified as *P. diasi*. The identification was made using the dichotomous keys of Lent and Wygodzinsky (1979) and Galvão and Dale (2014).

The specimens were deposited in the Entomology Laboratory ERSBG/SESMT and in the Triatominae Collection at the Oswaldo Cruz Institute (CTIOC).

Results

New records. Brazil: Mato Grosso: Barra do Garças (15°51.98' S, 052°16.29' W, 395 m alt.), collected by Moraes SC, March 2017 (female total length 27 mm, CTIOC/9641; females, lengths 27 mm ERSBG/SESMT 053; female, total length 26.8 mm, ERSBG/SESMT 054 (Fig. 2). Brazil: Mato Grosso: Pontal do Araguaia (15°40.72' S, 052°00.55' W, 370 m alt.), collected by agents of control of endemic diseases of Pontal do Araguaia, 2011.

Identification. For identification of this genus the main criteria is the position of the antennae, inserted close to the eyes (Fig. 3A). The species can be diagnosed as follows: anterolateral angles of pronotum very short; anterolateral margins of the pronotum forming a continuous line, almost straight (Fig. 3B, C); upper face of the head in lateral view straight; rostrum with second segment longer than first (Fig. 3D, E); femora with 2 or 3 denticles and scutellum process elongated, cylindrical and tapered at the apex (Fig. 3F, G) (Gurgel-Gonçalves et al. 2012b, Galvão and Dale 2014).



Figure 2. Female specimen of *Panstrongylus diasi* collected in Pontal do Araguaia (SESMT/ERSBG 053), Mato Grosso, Brazil in 2011.




Figure 3. Main criteria for identification of *Panstrongylus diiasi*: **A.** Antennal insertion, close to the eyes. **B.** Pronotum, anterolateral angles. **C.** Anterolateral margins in a continuous line. **D.** Upper face of straight head, lateral. **E.** Second segment of rostrum longer than the first. **F.** Femora with 2 denticles. **G.** Scutellum elongated and tapered.

Discussion

We extend the geographical distribution of *P. diiasi* to Mato Grosso state. These new records are consistent with the prediction map for the potential distribution of this species in Brazil published by Gurgel-Gonçalves et al. (2012a).

Lent and Wygodzinsky (1979: fig. 216) figured the type specimen of *P. diiasi* with the indication that it was a female from Mato Grosso, Brazil. The legend is discordant with the labeling of the specimen, which is not from Mato Grosso. We examined the specimen deposited at CTIOC and can verify that it was collected in Minas Gerais (Fig. 4). A similar situation was detected by Gil-Santana and Galvão (2013), that recorded an apparent discordance between the labels attached to the holotype of *Parabelminus yurupucu* Lent & Wygodzinsky, 1979 in the CTIOC and the information published by Lent and Wygodzinsky (1979). The figure of the holotype by these

authors confirms that it was this specimen type, not the paratype, which would have the information on the labels found with the holotype. Apparently, Lent and Wygodzinsky (1979) switched the information on the labels of the holotype and the paratype in their paper.

In Brazil, *P. diiasi* was recorded in the states of Bahia, Espírito Santo, Goiás, Maranhão, Mato Grosso do Sul, Minas Gerais, Rio Grande do Norte, São Paulo, and Tocantins, as well as the Distrito Federal. This species is frequently collected in artificial habitats but the sylvatic habitats as well as aspects of its biology, ecology, and genetics remain unknown (Patterson et al. 2009, Galvão et al. 2014, Barreto et al. 2017). The 2 cities in Mato Grosso, where the new records were found are adjacent to the state of Goiás, where a record of *P. diiasi* was collected in 2000 (Oliveira and Silva 2007).

Paula et al. (2010) and Brito et al. (2017) reported the natural infection of this species with *T. cruzi*. It is,




Figure 4. Type specimen and labels of a specimen of *P. diiasi* deposited in the Triatominae Collection at the Oswaldo Cruz Institute (CTIOC), Rio de Janeiro, Brazil.



Figure 5. Panoramic view of Barra do Garças municipality, Mato Grosso, Parque Estadual da Serra Azul, Pontal do Araguaia (Mato Grosso) and Aragarças (Goiás), Brazil.

therefore, important to know the ecology, biogeography, and parasite/host interactions to propose effective interventions to reduce the risk of Chagas disease.

Barra do Garças has a different ecological context, given its proximity to the Parque Estadual da Serra Azul,

which may facilitate the contact of sylvatic triatomines, potentially vectors of Chagas disease, with the human population (Fig. 5). The occurrence of sylvatic species that sporadically invade human dwellings are a major difficulty for vector surveillance programs (Caranha et al. 2011).

Acknowledgments

This research was supported by the Brazilian National Research Council (Conselho Nacional de Desenvolvimento Científico e Tecnológico). We thank Mirian Lacerda Golembiouski from the Secretaria de Estado de Saúde de Mato Grosso for supporting the surveillance actions, Rodrigo Gurgel-Gonçalves, University of Brasília, for confirming the identifications, Carolina Dale, Instituto Oswaldo Cruz-FIOCRUZ for reviewing the text, and Craig Woollard by revising the English.

Authors' Contributions

MCM conducted the fieldwork; MFM and OMT contributed to the preparation of the manuscript; GC studied the specimen and reviewed whole text.

References

- Almeida PS, Ceretti-Junior W, Obara MT, Santos HR, Barata JMS, Facenda O (2008) Levantamento da fauna de Triatominae (Hemiptera: Reduviidae) em ambiente domiciliar e infecção natural por Trypanosomatidae no Estado de Mato Grosso do Sul. Revista da Sociedade Brasileira de Medicina Tropical 41 (4): 374–380. <http://doi.org/10.1590/S0037-86822008000400010>
- Barreto MAF, Cavalcanti MAF, Andrade CM, Nascimento EGC, Oliveira-Pereira W (2017) Indicadores entomológicos de triatomíneos no Estado do Rio Grande do Norte, Brasil. Ciência Saúde Coletiva. Accessed on: 2018-4-07.
- Brito RN, Gorla DE, Diotauiti L, Gomes ACF, Souza RCM, Abad-Franch F (2017) Drivers of house invasion by sylvatic Chagas disease vectors in the Amazon-Cerrado transition: a multi-year, state-wide assessment of municipality-aggregated surveillance data. PLOS Neglected Tropical Diseases 11 (11): e0006035. <https://doi.org/10.1371/journal.pntd.0006035>
- Caranha L, Gurgel-Gonçalves R, Ramalho RD, Galvão C (2011) Novos registros e mapa de distribuição geográfica de *Triatoma petrocchiae* Pinto and Barreto, 1925 (Hemiptera: Reduviidae: Triatominae). Check List 7 (4): 508–509. <http://doi.org/10.15560/7.4.508>
- Instituto Brasileiro de Geografia e Estatística (IBGE) (2017). <http://www.ibge.gov.br/cidadesat/topwindow.htm?1>. Accessed on: 2017-8-21.
- Costa J, Almeida CE, Dotson EM, Lins A, Vinhaes M, Silveira AC, Beard CB (2003) The Epidemiologic Importance of *Triatoma brasiliensis* as a Chagas disease vector in Brazil: a revision of domiciliary captures during 1993–1999. Memórias Instituto Oswaldo Cruz 98 (4): 443–449. <http://doi.org/10.1590/S0074-02762003000400002>
- Galvão C (Ed.) (2014) Vetores da doença de Chagas no Brasil. Série Zoologia, Guia e manuais de identificação. Sociedade Brasileira de Zoologia, Curitiba, 289 pp.
- Galvão C, Dale C (2014) Chaves de identificação para adultos. In: Galvão C (Ed.) Vetores da doença de Chagas no Brasil. Série Zoologia, Sociedade Brasileira de Zoologia, Curitiba, 171–208.
- Gil-Santana HR, Galvão C (2013) Description of the male genitalia of *Belminus rugulosus* Stål and *Belminus corredori* Galvão & Angulo, and comments on the holotype of *Parabelminus yurupucu* Lent & Wygodzinsky (Hemiptera: Heteroptera: Reduviidae: Triatominae: Bolboderini). Zootaxa 3746 (4): 587–596. <https://doi.org/10.11646/zootaxa.3746.4>
- Gil-Santana HR, Galvão C, Mielke CGC (2014) *Cavernicola pilosa* Barber, 1937 (Hemiptera: Reduviidae: Triatominae): first report in the state of Maranhão, Brazil. Check List 10 (4): 944–946. <https://doi.org/10.15560/10.4.944>
- Gurgel-Gonçalves R, Galvão C (2014) Vetores conhecidos no Brasil. In: Galvão C (Ed.) Vetores da doença de Chagas no Brasil. Série Zoologia, Sociedade Brasileira de Zoologia, Curitiba, 171–208.
- Zoologia, Sociedade Brasileira Zoologia, Curitiba, 88–170.
- Gurgel-Gonçalves R, Galvão C, Costa J, Peterson AT (2012a) Geographic distribution of Chagas disease vectors in Brazil based on ecological niche modeling. Journal of Tropical Medicine 2012 (1): 705326. <https://doi.org/10.1155/2012/705326>
- Gurgel-Gonçalves R, Galvão C, Mendonça J, Costa-Neto EM (2012b) Vetores conhecidos no Brasil. In: Guia de triatomíneos da Bahia. Ed. UEFS, Feira de Santana, 19–26.
- Justi SA, Galvão C (2017) The evolutionary origin of diversity in Chagas disease vectors. Trends in Parasitology 33 (1): 42–52. <https://doi.org/10.1016/j.pt.2016.11.002>
- Köppen W (1948) Climatología: comun estudio de los climas de la tierra. Fondo de Cultura Económica, México, 478 pp.
- Lent H, Wygodzinsky PW (1979) Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. Bulletin of the American Museum of Natural History 163: 520–529.
- Maeda MH, Knox MB, Gurgel-Gonçalves R (2012) Occurrence of synanthropic triatomines (Hemiptera: Reduviidae) in the Federal District of Brazil. Revista da Sociedade Brasileira de Medicina Tropical 45 (1): 71–76.
- Marassá AM, Barata JMS (2000) Nota sobre tipos e espécimes de triatomíneos depositados na Coleção Entomológica do Instituto Butantan, São Paulo (Hemiptera: Reduviidae). Revista da Sociedade Brasileira de Medicina Tropical 33: 227–230.
- Mendes PC, Lima SC, Paula MBC, Souza AA, Rodrigues EAS, Limongi JE (2008) Doença de Chagas e a distribuição espacial de triatomíneos capturados em Uberlândia, Minas Gerais – Brasil. Hygeia 3 (6): 176–204.
- Ministério do Meio Ambiente (MMA) (2002) Secretaria de biodiversidade e florestas. Avaliação e identificação de áreas e ações prioritárias para conservação, utilização sustentável e repartição dos benefícios da biodiversidade nos biomas brasileiros. Ministério do Meio Ambiente, Brasília, 404 pp.
- Oliveira AWS, Silva IG (2007) Distribuição geográfica e indicadores entomológicos de triatomíneos sinantrópicos capturados no Estado de Goiás. Revista da Sociedade Brasileira de Medicina Tropical 40 (2): 204–208.
- Paula AM, Castrillon MI, Maciel GBML, Galvão C (2013) Registro de ocorrência de *Panstrongylus lignarius* (Walker, 1837) (Hemiptera: Reduviidae: Triatominae) no Estado de Mato Grosso, Brasil. Acta Amazônica 43 (1): 117–119. <http://doi.org/10.1590/S0044-59672013000100015>
- Paula MBC, Costa INPAF, Limongi JE, Pajuaba-Neto AA, Pinto RMC, Gonçalves ALR, Costa-Cruz JM (2010) Occurrence of positivity for *Trypanosoma cruzi* in triatomine from municipalities in southeastern Brazil, from 2002 to 2004. Revista da Sociedade Brasileira de Medicina Tropical 43 (1): 9–14. <https://doi.org/10.1590/S0037-86822010000100003>
- Patterson JS, Barbosa SE, Feliciangeli MD (2009) On the genus *Panstrongylus* Berg 1879: evolution, ecology and epidemiological significance. Acta Tropica 110 (2–3): 187–199. <https://doi.org/10.1016/j.actatropica.2008.09.008>
- Ponair-Junior G (2019) A primitive triatomine bug, *Paleotriatoma metaxytaxa* gen. et sp. nov. (Hemiptera: Reduviidae: Triatominae), in mid-Cretaceous amber from northern Myanmar. Cretaceous Research 93: 90–97. <https://doi.org/10.1016/j.cretres.2018.09.004>
- Rêbelo JMM, Barros VLL, Mendes WA (1998) Espécies de Triatominae (Hemiptera: Reduviidae) do Estado do Maranhão, Brasil. Cadernos de Saúde Pública 14 (1): 187–192.
- Rosa JA, Justino HHG, Nascimento D, Mendonça VJ, Rocha CS, Blanco de Carvalho D, Falcone R, Azeredo-Oliveira MTV, Alevi KCC, Oliveira J (2017) A new species of *Rhodnius* from Brazil (Hemiptera, Reduviidae, Triatominae). ZooKeys 675: 1–25. <https://doi.org/10.3897/zookeys.675.12024>
- Santos CM, Jurberg J, Galvão C, Rocha D, Fernandez JIR (2003) Estudo Morfométrico do Gênero *Panstrongylus* Berg, 1879 (Hemiptera, Reduviidae, Triatominae). Memórias do Instituto Oswaldo Cruz 98

- (7): 939–944.
- Santos CB, Rangel CV, Leite CR (2004) Ocorrência de *Panstrongylus diasi* Pinto & Lent, 1946 (Hemiptera: Reduviidae: Triatominae) no estado do Espírito Santo, Brasil. Entomología y Vectores 11 (2): 363–367.
- Sherlock IA, Serafim EM (1972) Fauna triatominae do Estado da Bahia, Brasil. I – As espécies e distribuição geográfica. Revista da Sociedade Brasileira de Medicina Tropical 6 (5): 266–276.
- Silva JMC, Bates JM (2002) Biogeographic patterns and conservation

in the South American Cerrado: a tropical savanna hotspot: the Cerrado, which includes both forest and savanna habitats, is the second largest South American biome, and among the most threatened on the continent. BioScience 52 (3): 225–234.

- Souza ES, Von Atzingen NCB, Furtado MB, Oliveira J, Nascimento JD, Vendrami D, Gardim S, Rosa JA (2016) Description of *Rhodnius marabaensis* sp. n. (Hemiptera, Reduviidae, Triatominae) from Pará state, Brazil. ZooKeys 621: 45–62. <https://doi.org/10.3897/zookeys.621.9662>