



Rediscovery and expansion of the distribution of *Phyllanthus itatiaiensis* Brade (Phyllanthaceae) in southeastern Brazil

Jone Clebson Ribeiro Mendes¹, Sarah Maria Athiê-Souza^{1*}, Rayane de Tasso Moreira Ribeiro¹, Caio Baez Gomes², Tatiana Tavares Carrijo², Margareth Ferreira de Sales¹

1 Programa de Pós-Graduação em Biodiversidade, Universidade Federal Rural de Pernambuco, Recife, PE, Brazil • JCRM: jhonnie321@hotmail.com  <https://orcid.org/0000-0002-9696-1978> • SMAS: sarah_athie@yahoo.com.br  <https://orcid.org/0000-0002-6090-981X> • RTMR: rayane.tasso@gmail.com  <https://orcid.org/0000-0001-6006-598X> • MFS: mfsales65@hotmail.com  <https://orcid.org/0000-0002-9587-1209>

2 Centro de Ciências Agrárias, Universidade Federal do Espírito Santo, Alegre, ES, Brazil • CBG: caiobg@cnclflora.net • TTC: tcarrijo@gmail.com  <https://orcid.org/0000-0002-8366-0738>

* Corresponding author

Abstract

We report here the rediscovery of *Phyllanthus itatiaiensis* Brade (Phyllanthaceae) in Rio de Janeiro after 50 years. We also report new occurrence records of this species in Minas Gerais state, southeastern Brazil. We provide a taxonomic diagnosis, *in situ* photographs, a distribution map, and a reassessment of its conservation status, which will improve the knowledge of this rare, endemic species as well as the ability to protect it. *Phyllanthus itatiaiensis* is currently classified as Endangered under IUCN criteria B1ab(iii), B2ab(iii).

Keywords

Atlantic domain, biodiversity hotspot, endemism, *Phyllanthus* subsect. *Clausseniani*, taxonomy

Academic editor: Matias Köhler | Received 15 January 2021 | Accepted 14 April 2021 | Published 22 April 2021

Citation: Mendes JCR, Athiê-Souza SM, Ribeiro RTM, Gomes CB, Carrijo TT, Sales MF (2021) Rediscovery and expansion of the distribution of *Phyllanthus itatiaiensis* Brade (Phyllanthaceae) in southeastern Brazil. Check List 17 (2): 695–700. <https://doi.org/10.15560/17.2.695>

Introduction

The Atlantic Forest is considered a global biodiversity hotspot. It extends along the eastern coast of South America, with 90% of its area within Brazilian territory. Thus, it represents one of the principal phytogeographic domains in Brazil (Ribeiro et al. 2009; Colla-Silva et al. 2020; Ostroski et al. 2020). The forest retains enormous species richness and high endemism of both plant and animal taxa (Myers et al. 2000). Recent estimates indicate that the Brazilian Atlantic Forest holds approximately 15,000 native species of angiosperms, half of which are exclusive to this phytogeographic domain (BFG 2018).

In addition to being a hotspot for plant diversity, the Brazilian Atlantic Forest is also associated with a wide variety of geomorphological formations, including highland grasslands (Rizzini 1997; Pereira et al. 2020). Those mountain phytophysiognomies associated with the Atlantic Forest combine unique abiotic and biotic conditions that favor high species diversity (Segadas-Viana and Dau 1965; Rizzini 1997; Safford 1999).

The Itatiaia Massif in the Serra da Mantiqueira Range stands out among the many mountain formations found within the Atlantic Forest domain and has been the target of plant research and collection campaigns by prominent

European and Brazilian naturalists, including Auguste de Saint-Hilaire, Ernesto Ule, Friedrich Sellow, José M. da C. Vellozo, and Per Karl Dusén (Martinelli et al. 1989; Coelho et al. 2017; Carrijo et al. 2018). Those historic expeditions, which took place in the 19th and early 20th centuries, contributed to the discovery of many plant species, as well as the development of the *Flora fluminensis* (Coelho et al. 2017; Colli-Silva et al. 2020).

Recent discoveries of new taxa (Cardoso et al. 2019; Vanderplank 2020) and biogeographic records (Santiago et al. 2018), as well as the rediscovery of numerous species (Miranda and Menini Neto 2010; Guimarães et al. 2020) in the Serra da Mantiqueira region, reinforce the view that the area comprises significant floristic richness and diversity, and needs to be further explored with floristic and phytosociological focuses (Pompeu et al. 2014).

Species of many different plant groups can be found in the Serra da Mantiqueira mountains, including *Phyllanthus* L. Fifty-four species of the genus have been recorded in the Atlantic Forest, although taxa restricted to highland grasslands in that domain have only been poorly studied and are represented by few collected specimens, such as *Phyllanthus pinifolius* Baill. and *Phyllanthus itatiaiensis* Brade (Martins et al. 2014, 2017; Orlandini et al. 2021).

Phyllanthus itatiaiensis, a taxon established by Brade (1957), was only known from the type collection. According to Martins et al. (2014), the species is endangered, and only limited information is available on its geographic distribution and conservation status. Recently, however, populations of *P. itatiaiensis* were found in Serra da Mantiqueira, in the Itatiaia National Park (INP), during the project “Rediscovering endangered species in Conservation Units (UCs) in the Atlantic Forest: bases for management, conservation and access to information”. Additionally, specimens of *P. itatiaiensis* that had mistakenly been identified as *Phyllanthus niruri* L. were found among HUCS Herbarium collections. Despite relatively constant floristic studies in the type locality, this species has not been reported for more than 50 years.

We report here the rediscovery of *P. itatiaiensis*, 50 years after its last collection in the type locality, as well as a new occurrence record for Minas Gerais state in southeastern Brazil. We include a taxonomic diagnosis, *in situ* photographs, a distribution map, and a reassessment of the conservation status of *P. itatiaiensis*, which will all contribute to a better understanding of its distribution and conservation status in INP in the Serra da Mantiqueira.

Methods

Area characterization. The INP, comprises the Itatiaia massif, in the Serra da Mantiqueira, and includes municipalities in the states of Rio de Janeiro (RJ) and Minas Gerais (MG) (Moreira et al. 2020). The area has a mesothermal climate and a rugged and rocky landscape ranging from 700 to 2,787 m a.s.l. (Barreto et al. 2013).

That geomorphological situation, as already mentioned, results in the establishment of different phytophysiognomies, depending on the elevation (Segadas-Vianna and Dau 1965; Geise et al. 2004).

Sampling and data treatment. The present study was based on the study of collections held in the F, HUEFS, MBM, MG, NY, RB, SP, and HUCS herbaria (acronyms according to Thiers 2021), consultations of the specialized literature (Brade 1957; Webster 2002; Martins et al. 2014), and field expeditions carried out between October and December/2019 in INP in the municipalities of Resende (RJ) and Itamonte (MG). The specimens observed in the field were photographed using a digital camera (Nikon Coolpix P500) and collected following Fidalgo and Bononi (1989). Field specimens were deposited in the RB herbarium.

The species' conservation status was determined following IUCN (2012) criteria and the use of the Geocat tool (<http://geocat.kew.org>; Bachman et al. 2011). The distribution map was prepared using Quantum GIS v. 2.8.2 (Quantum GIS development team 2018).

Results

Phyllanthus itatiaiensis Brade

Figure 1C–F

New records (Fig. 2). BRAZIL • Minas Gerais, Itamonte; 22°21'37"S, 044°40'07"W; 16 Dec. 2019; fl., fr.; J. Kükamp et al. 1117 (RB 01426341) • Rio de Janeiro state, Itatiaia; 13 March 2010; fr. J.M. Silva 7550 (HUCS 36267). • Rio de Janeiro state, Resende; 22°22'27"S, 044°40'22"W; 31 Oct. 2019; fl., fr. C. Baez & A. Sá 1780 (RB 01423645) • Rio de Janeiro state, Resende; 22°22'28"S, 044°40'33"W; 3 Nov. 2019; fl., fr.; C. Baez & A. Sá 1800 (RB 01423665).

Additional material examined. BRAZIL • Rio de Janeiro state, Itatiaia; Mar. 1937; fl. A.C. Brade 15575 (RB 00569788, holotype) • Rio de Janeiro state, Itatiaia; May 1950; fl. A.C. Brade 20362 (RB 00545529, paratype).

Identification. *Phyllanthus itatiaiensis* can be distinguished from other species of the genus by having oblong to falcate leaves, and stipules 1–1.5 mm long, triangular-lanceolate, showy, and pink. It resembles *P. niruri* L. in terms of their shared herbaceous habits, as well as by having staminate and pistillate flowers with five sepals, and the androecium formed by three stamens with free filaments; it differs from *P. niruri* by having oblong to falcate leaves with slightly revolute margins (vs. oblong leaves with flat margins in *P. niruri*), stipules 1–1.5 mm long, triangular-lanceolate, showy, pink (vs. 0.5–1 mm long, linear to lanceolate, not showy, greenish), and the pedicels of the male flowers 1–2 mm long. (vs. 2.5–4 mm long).

Habitat and distribution. *Phyllanthus itatiaiensis* is




Figure 1. *Phyllanthus itatiaiensis*. **A, B.** Habitat, with emphasis on the Pedra do Altar rock formation. **C, D.** Habitat. **E.** Staminate flower. **F.** Pistillate flower and fruits. Photographs: Caio Baez (A, B); J. Kükamp (C–F).

endemic to Brazil, with its distribution restricted to the Atlantic Forest in the southeastern region of the country. It is endemic to highland grasslands on the Itatiaia Massif, in the municipalities of Itatiaia and Resende (RJ) and Itamonte (MG) (Figs. 1A, B, 2). The species is often found in wetlands and along lakesides where it is associated with *Chusquea pinifolia* Nées (Poaceae) (Martins et al. 2014).

Phenology. Flowering and fruiting between March and December.

Conservation status. The species is classified as Endangered under criteria Blab(iii), B2ab(iii) (IUCN 2012). The area of occupation is 80.5 km² and the extent of occurrence is 164.9 km². The main threats to *P. itatiaiensis* are related to intense losses of areas of occupation as well as habitat quality degradation, the latter due to constant forest fires and tourism in the INP.

Comments. Following the classification proposed by Webster (2002), *P. itatiaiensis* was circumscribed under *P. subsect. Clausseniani* Webster for having stamens




Figure 2. Distribution of *Phyllanthus itatiaiensis* in Brazil. New record from Rio de Janeiro state (circles) and Minas Gerais (star); the type collection (square).

with free filaments, anthers deeply emarginated, and a distinct theca. The species is morphologically related to *P. mocotensis* G.L Webster and *P. carvalhoi* G.L Webster, but can be distinguished from the former because it is an herb with typically phyllanthoid branches (vs. sub-shrub, with rigid branches with fissures), and from the latter by having oblong leaves, rarely falcate (vs. falcate-asymmetrical). From a phylogenetic point of view, until now, the studies carried out (Samuel et al. 2005; Kathriarachchi et al. 2006; Bouman et al. 2020, Hidalgo et al. 2020) have not included *P. itatiaiensis*. For this reason, one of us (J.C.R. Mendes) is conducting molecular phylogenetic analyses of the species of the subsection *Claussenianii*.

Discussion

Although specimens of *Phyllanthus itatiaiensis* were collected in highland grasslands within a protected area (the INP), those grasslands have undergone both natural and anthropic threats to the species (Assis and Matos 2016; Gonçalves and Santos 2018). Among the anthropic threats are fires and unregulated tourism, while changes in environmental conditions reflect intensifying climate change (Aximoff and Rodrigues 2011; Assis and Matos 2016). The INP experiences major fires almost every year, as well as increasing tourism throughout its extent (Lima 2000; Aximoff and Rodrigues 2011). To avoid or mitigate losses in biodiversity, it will be important to

establish proactive environmental actions and policies within this and other conservation areas in the Atlantic Forest—and especially in highland grasslands—focusing on stricter enforcement and the implementation of measures to prevent and combat fires and other threats. Additionally, although many recent studies have focused on cataloging the richness of the INP flora (e.g., Morim and Barroso 2007; Giannerini et al. 2007, 2015; Barberena et al. 2008; Pereira and Mansano 2008; Monteiro and Guimarães 2009; Barros et al. 2010; Ramos and Sylvestre 2010; Mezabarba et al. 2013; Rollim and Trovó 2016; Freitas and Trovó 2017; Gonzaga et al. 2017), additional floristic surveys will be necessary for searching for *Phyllanthus* species in highland grasslands.

Many species of the genus have been recorded in the Serra da Mantiqueira region, and some of them are endemic (e.g., *P. acutifolius* Poir. ex Spreng. and *Phyllanthus pinifolius*). Field expeditions in little- or unexplored areas of highland grasslands may reveal new occurrence records, new species, or even new populations of *P. itatiaiensis*, which would contribute to the conservation of this species and other rare and endemic species in the Brazilian Atlantic Forest.

Acknowledgments

We are grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) for the PhD scholarship awarded to the first author. Additionally, we extend our thanks to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) – Finance

Code 001, and the Fundação de Amparo a Ciência e Tecnologia de Pernambuco (FACEPE); T.T. Carrijo and C. Baez are grateful to “Instituto Chico Mendes de Conservação da Biodiversidade” (ICMBio), “Conselho Nacional de Desenvolvimento Científico e Tecnológico” (CNPq 421276/2017-7) and “Fundação de Amparo à Pesquisa e Inovação do Espírito Santo” (FAPES 34/2018) for the research grant (“Chamada CNPq/ICMBio/FAPs Nº 18/2017 - Pesquisa em Unidades de Conservação da Caatinga e Mata Atlântica”); T.T. Carrijo is also grateful to CNPq for the grant “Bolsa de Produtividade em Pesquisa”, and to FAPES for the research grant “Taxa de Pesquisa”. We thank the reviewers for their critical comments and suggestions to improve this article and Josimar Kükamp for his valuable field contributions and photographs.

Authors' Contributions

CBG collected the specimen and identified the specimen; JCRM designed the study and wrote the manuscript; AMT contributed to the writing of the manuscript; SMAS contributed to the discussion and review; RTMR contributed to the writing, reading, review, and correction of the manuscript; MFS contributed to the correction and research funding; TTC contributed to the correction of the manuscript and research funding.

References

- Assis V, Mattos EA. (2016) Vulnerabilidade da Vegetação de Campos de Altitude às Mudanças Climáticas. *Oecologia Australis* 20: 24–36. <http://dx.doi.org/10.4257/oeco.2016.2002.03>
- Aximoff I, Rodrigues RC (2011) Histórico dos incêndios florestais no Parque Nacional do Itatiaia. *Ciência Florestal* 21: 83–92. <https://doi.org/10.5902/198050982750>
- Bachman S, Moat J, Hill AW, Torre J, Scott B (2011) Supporting Red List threat assessments with GeoCAT: geospatial conservation assessment tool. *ZooKeys* 150: 117–126. <https://doi.org/10.3897/zookeys.150.2109>
- Barberena FF, Villela A, Baumgratz JFA, Chiavegatto B (2008) Melastomataceae no Parque Nacional do Itatiaia, Sudeste do Brasil: tribos Bertoloniaceae e Merianiae. *Rodriguésia* 59: 381–392. <https://doi.org/10.1590/2175-7860200859210>
- Barreto CG, Campos JB, Roberto DM, Schwarzstein NT, Alves GSG, Coelho WC (2013) Plano de manejo: Parque Nacional do Itatiaia. Instituto Chico Mendes de Conservação da Biodiversidade, Ministério do Meio Ambiente, Brasília, Brazil, 215 pp. https://www.icmbio.gov.br/portal/images/stories/docs-planos-de-manejo/pm_parna_itatiaia_encl.pdf. Accessed on 2020-12-01.
- Barros MJG, Mansano VF, Chautems A (2010) Comparações florísticas e taxonomia da família Gesneriaceae no Parque Nacional do Itatiaia, Brasil. *Hoehnea* 37: 131–145.
- BFG (2018) Brazilian Flora 2020 innovation and collaboration to meet Target 1 of the Global Strategy for Plant Conservation (GSPC). *Rodriguésia* 69: 1513–1527. <http://doi.org/10.1590/2175-7860201869402>.
- Bouman RW, Keßler PJA, Telford IRH, Bruhl JJ, Strijk JS, Saunders RMK, van Welzen PC (2020) Molecular phylogenetics of Phyllanthus sensu lato (Phyllanthaceae): towards coherent monophyletic taxa. *Taxon* 70: 72–98. <https://doi.org/10.1002/tax.12424>
- Brade AC (1957). *Phyllanthus itatiaiensis*. *Archivos do Jardim Botânico do Rio de Janeiro* 15: 9.
- Cardoso PH, Neto LM, Salimena FRG (2019) *Lippia mantiqueirae* (Verbenaceae), a new species from Minas Gerais, Brazil. *Phytotaxa* 420: 249–254. <https://doi.org/10.11646/phytotaxa.420.3.5>
- Carrijo TT, Alves-Araújo AG, Amorim AMA, Barbosa DEF, Barcellos LB, Baumgratz JF, Bueno VR, Coelho RLG, Costa DP, Couto DR, Delgado CN, Dutra VF, Flores TB, Furtado SG, Giacomini LL, Goldenberg R, Gomes M, Gonzaga DR, Guimarães EF, Heiden G, Kameyama C, Labiak PHE, Lírio EJ, Lohmann LG, Matos, FB, Moraes PLR, Meireles LD, Menini-Neto L, Monteiro D, Moreira MM, Morim MP, Mota MCA, Oliveira, JRPM, Pastore JFB, Pederneiras LC, Pereira LC, Rapini A, Salimena FRG, Silva AV, Silva-Neto SJ, Sobral MEG, Souza MC, Sylvestre LS, Trovó M, Viana PL, Forzza RC (2018) Lista de espécies de plantas terrestres do Parque Nacional do Itatiaia. In: *Catálogo de Plantas das Unidades de Conservação do Brasil. Jardim Botânico do Rio de Janeiro*. <https://catalogo-ucs-brasil.jbrj.gov.br>. Accessed on: 2020-12-31.
- Coelho, MAN, Baumgratz JFA, Lobão AQ, Sylvestre LS, Trovó M, Silva, LAE (2017) Flora do estado do Rio de Janeiro: avanços no conhecimento da diversidade. *Rodriguésia* 68: 1–11. <https://doi.org/10.1590/2175-7860201768101>
- Colli-Silva M, Reginato M, Cabral A, Forzza RC, Pirani JR., Vasconcelos TNDC (2020). Evaluating shortfalls and spatial accuracy of biodiversity documentation in the Atlantic Forest, the most diverse and threatened Brazilian phytogeographic domain. *Taxon* 69 (3): 567–577. <https://doi.org/10.1002/tax.12239>
- Fidalgo O, Bononi VL (1984) Técnicas de coleta, preservação e herborização de material botânico. Instituto de Botânica, São Paulo, 62 pp.
- Freitas SNS, Trovó M (2017) Eriocaulaceae no Parque Nacional do Itatiaia, Brasil. *Rodriguésia* 68: 1387–1395. <https://doi.org/10.1590/2175-7860201768419>
- Geise L, Pereira LG, Bossi DEP, Bergallo HG (2004) Pattern of elevational distribution and richness of non volant mammals in Itatiaia National Park and its surroundings, in Southeastern Brazil. *Brazilian Journal of Biology* 64: 599–612. <https://doi.org/10.1590/S1519-69842004000400007>
- Giannerini AC, Quinet A, Andreata RHP (2007) O gênero *Ocotea* Aubl. (Lauraceae) no Parque Nacional do Itatiaia, Brasil. *Pesquisas, Botânica* 58: 283–329.
- Giannerini AC, Quinet A, Andreata RHP (2015) Lauraceae no Parque Nacional do Itatiaia, Brasil. *Rodriguésia* 66: 863–880. <https://doi.org/10.1590/2175-7860201566314>
- Gonçalves MTA, Santos ND (2018) Campos de altitude do Parque Nacional do Itatiaia: um hotspot para briófitas. *Diversidade e Gestão* 2: 90–105.
- Gonzaga DR, Neto LM, Peixoto AL (2017) Cactaceae no Parque Nacional do Itatiaia, Serra da Mantiqueira, Brasil. *Rodriguésia* 68: 1397–1410. <https://doi.org/10.1590/2175-7860201768420>
- Guimarães PJF, Baez C, Rosa P, Martinelli G, Silva DN (2020) The importance of rediscovering rare and endemic plants: two species of *Pleroma* D. Don (Melastomataceae), an update of conservation status, and improved descriptions. *Webbia* 75: 293–304. <https://doi.org/10.36253/jopt-8902>
- Gonzaga DR, Menini Neto L, Peixoto AL (2017) Cactaceae no Parque Nacional do Itatiaia, Serra da Mantiqueira, Brasil. *Rodriguésia* 68: 1397–1410. <https://doi.org/10.1590/2175-7860201768420>
- Hidalgo BF, Bazan SF, Iturrealde RB, Borsch T (2020) Phylogenetic relationships and character evolution in neotropical Phyllanthus (Phyllanthaceae), with a focus on the Cuban and Caribbean taxa. *International Journal of Plant Sciences* 181: 284–305. <https://doi.org/10.1086/706454>
- IUCN (2012) The IUCN Red List categories and criteria: version 3.1. International Union for the Conservation of Nature, Gland, Switzerland and Cambridge, UK, 32 pp.
- Kathriarachchi H, Samuel R, Hoffmann P, Mlinarec J, Wurdack KJ, Ralimanana H, Stuessy TF, Chase MW (2006) Phylogenetics of tribe Phyllantheae (Phyllanthaceae; Euphorbiaceae sensu lato) based on nrITS and plastid matK DNA sequence data. *Ameri-*

- can Journal of Botany 93: 637–655. <https://doi.org/10.3732/ajb.93.4.637>
- Lima GS (2000) Os incêndios florestais no Estado de Minas Gerais. Revista Ação Ambiental 2: 15–18.
- Martinelli G, Bandeira J, Bragança JO (1989) Campos de Altitude. Editora Index, Rio de Janeiro, Brazil, 160 pp.
- Martins ER, Lima LR, Cordeiro I (2014). *Phyllanthus* (Phyllanthaceae) no estado do Rio de Janeiro. Rodriguésia 65: 405–424. <https://doi.org/10.1590/S2175-78602014000200007>
- Martins ER, Lima LR, Cordeiro I (2017) Flora do Espírito Santo: *Phyllanthus* (Phyllanthaceae). Rodriguésia 68: 1813–1828. <https://doi.org/10.1590/2175-7860201768517>
- Mezabarba V, Vianna Filho MDM, Borges RAX, Mansano VF (2013) Ericaceae do Parque Nacional do Itatiaia, RJ, Brasil. Hoehnea 40: 115–130.
- Miranda MR, Menini Neto L (2010) Redescoberta de *Centroglossa macroceras* Barb. Rodr. e *Thysanoglossa jordanensis* Porto & Brade (Orchidaceae), duas micro-orquídeas consideradas presumivelmente extintas no Estado de São Paulo, Brasil. Hoehnea 37: 403–405. <https://doi.org/10.1590/S2236-89062010000200012>
- Monteiro D, Guimarães EF (2009) Flora do Parque Nacional do Itatiaia – Brasil: *Manekia* e *Piper* (Piperaceae). Rodriguésia 60: 999–1024. <https://doi.org/10.1590/2175-7860200960413>
- Moreira MM, Carrijo TT, Alves-Araújo A, Amorim AMA, Rapini A, Silva AVS, Cosenza BAP, Lopes CR, Delgado CN, Kameyama C, Couto DR, Barbosa DEF, Monteiro D, Gonzaga DR, Dalcin EC, Guimarães EF, Lírio EJ, Matos FB, Salimena FRG, Oliveira FA, Heiden G, Lanna JM, Baumgratz JF, Pastore JFB, Oliveira JRPM, Barcelos LB, Sylvestre LS, Freitas L, Giacomini LL, Pederneiras L, Meireles LD, Lohmann LG, Pereira LC, Silva LAE, Neto LM, Souza MC, Trovó M, Sobral MEG, Garbin ML, Gomes M, Morim MP, Mota MCA, Labiak PH, Viana PL, Moraes PLR, Goldenberg R, Coelho RLG, Furtado SG, Silva-Neto SJ, Flores TB, Dutra VF, Bueno VR, Forzza RC (2020) Using online databases to produce comprehensive accounts of the vascular plants from the Brazilian protected areas: the Parque Nacional do Itatiaia as a case study. Biodiversity Data Journal 8: e50837. <https://doi.org/10.3897/BDJ.8.e50837>
- Morim MP, Barroso GM (2007) Leguminosae arbustivas e arbóreas da floresta atlântica do Parque Nacional do Itatiaia, sudeste do Brasil: subfamílias Caesalpinoideae e Mimosoideae. Rodriguésia 58: 423–468. <https://doi.org/10.1590/2175-7860200758216>
- Myers N, Mittermeier RA, Mittermeier CG, Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities, Nature 403: 853–858. <https://doi.org/10.1038/35002501>
- Orlandini P, Torres AM, Mendes JCR, Silva MJ. *Phyllanthus* in Flora do Brasil 2020 em construção. Jardim Botânico do Rio de Janeiro. <http://floradobrasil.jbrj.gov.br/reflora/floradobrasil/FB38526>. Accessed on 2021-01-11.
- Ostroski P, Saiter FZ, Amorim AM, Fiaschi P (2020) Angiosperm endemism in a Brazilian Atlantic Forest biodiversity hot-point. Brazilian Journal of Botany 43: 397–404. <https://doi.org/10.1007/s40415-020-00603-w>
- Pereira PH, Mansano VF (2008) Estudos taxonômicos da tribo Tecomeae (Bignoniacae) no Parque Nacional do Itatiaia, Brasil. Rodriguésia 59: 265–289. <https://doi.org/10.1590/2175-7860200859201>
- Pereira LC, Chautems A, Menini Neto L (2020) Biogeography and Conservation of Gesneriaceae in the Serra da Mantiqueira, Southeastern Region of Brazil. Brazilian Journal of Botany 43: 1–10. <https://doi.org/10.1007/s40415-020-00671-y>
- Pompeu PV, Fontes MAL, Santos RM, Garcia PO, Batista TA, Carvalho WAC, Oliveira Filho AT (2014) Floristic composition and structure of an upper montane cloud forest in the Serra da Mantiqueira Mountain Range of Brazil. Acta Botanica Brasiliensis 28: 456–464. <https://doi.org/10.1590/0102-33062014abb3239>
- QGIS – Development Team (2018) Sistema de Informações Geográficas do QGIS. Projeto Código Aberto Geospatial Foundation. <http://qgis.osgeo.org>. Accessed on 2020-12-1.
- Ramos CGV, Sylvestre LS (2010) Lycopodiaceae no Parque Nacional do Itatiaia, RJ e MG, Brasil. Acta Botanica Brasiliensis 24: 25–46. <https://doi.org/10.1590/S0102-33062010000100004>
- Ribeiro MC, Metzger JP, Martensen AC, Ponzoni FJ, Hirota MM (2009) The Brazilian Atlantic Forest: how much is left, and how is the remaining forest distributed? Implications for conservation. Biological Conservation 142: 1141–1153.
- Rizzini CT (1997) Tratado de fitogeografia do Brasil: aspectos ecológicos, sociológicos e florísticos. 2ed. Âmbito Cultural, Rio de Janeiro, Brazil, 746 pp.
- Rollim IM, Trovó M (2016) Campanulaceae no Parque Nacional do Itatiaia, Brasil. Rodriguésia 67: 1025–1030. <https://doi.org/10.1590/2175-7860201667412>
- Safford HD (1999) Brazilian Páramos I: an introduction to the physical environment and vegetation of the campos de altitude. Journal of Biogeography 26: 693–712.
- Samuel R, Kathriarachchi H, Hoffman P, Barfuss KJ, Wurdack MH, Davis CC, Chase MW (2005) Molecular phylogeny of Phyllanthaceae: evidence from plastid matK and nuclear PHYC sequences. American Journal of Botany 92: 132–141. <https://doi.org/10.3732/ajb.92.1.132>
- Santiago DS, Oliveira Filho AT, Menini Neto L, Carvalho FA, Salimena FRG (2018) Floristic composition and phytogeography of an Araucaria Forest in the Serra da Mantiqueira, Minas Gerais, Brazil. Rodriguésia 69: 1909–1925. <https://doi.org/10.1590/2175-7860201869426>
- Segadas-Vianna F, Dau L (1965) Ecology of the Itatiaia Range, Southeastern Brazil – Altitudinal Zonation of the Vegetation. Arquivos do Museu Nacional 53: 7–30.
- Thiers B (2021, continuously updated) Index herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/>. Accessed on: 2020-12-1.
- Vanderplank J (2020) *Passiflora itatiaiensis*—Passifloraceae: a new species of *Passiflora* from Brazil. Curtis's Botanical Magazine 37: 139–143. <https://doi.org/10.1111/curt.12326>
- Webster GL (2002). A synopsis of the Brazilian taxa of *Phyllanthus* section *Phyllanthus* (Euphorbiaceae). Lundelia 5: 1–26.