



New records of two teleost species from two insular areas of the western equatorial Atlantic

Rodrigo Xavier Soares^{1*}, Karlla Danielle Jorge Amorim¹, Amanda Torres Borges¹, Wagner Franco Molina¹, José Garcia Júnior²

1 Departamento de Biologia Celular e Genética, Centro de Biociências, Universidade Federal do Rio Grande do Norte, Natal, RN, Brazil • RXS: roxsoares@gmail.com  <https://orcid.org/0000-0002-3735-3649> • KDJA: karlla_danielle@msn.com  <https://orcid.org/0000-0003-3441-4298> • ATB: amanda_borges6@yahoo.com.br  <https://orcid.org/0000-0003-0776-1980> • WFM: molinawf@yahoo.com  <https://orcid.org/0000-0002-6695-0952>

2 Instituto Federal de Educação, Ciência e Tecnologia do Rio Grande do Norte, Campus Currais Novos, Currais Novos, RN, Brazil; jose.garcia@ifrn.edu.br  <https://orcid.org/0000-0002-2608-4877>

* Corresponding author

Abstract

We report the first record of two teleost species from two archipelagos in the western equatorial Atlantic. We recorded the occurrence of *Cantherhines pullus* (Ranzani, 1842) (Monacanthidae, Tetraodontiformes) from the Fernando de Noronha Archipelago, a group of volcanic islands 345 km off the northeastern coast of Brazil. We also report the first regional record of *Hemiramphus brasiliensis* (Linnaeus, 1758) (Hemiramphidae, Beloniformes) from the Saint Peter and Saint Paul's Archipelago, which is a small and isolated group of rocky islands 520 km from the Fernando de Noronha Archipelago on the Mid-Atlantic Ridge.

Keywords

Brazil, Fernando de Noronha Archipelago; ichthyofauna; Saint Peter and Saint Paul's Archipelago

Academic editor: Arturo Angulo | Received 26 August 2021 | Accepted 27 October 2021 | Published 10 November 2021

Citation: Soares RX, Amorim KDJ, Borges AT, Molina WF, Garcia Júnior J (2021) New records of two teleost species from two insular areas of the western equatorial Atlantic. Check List 17 (6): 1515–1520. <https://doi.org/10.15560/17.6.1515>

Introduction

The inventory of the ichthyofauna of oceanic island environments is of considerable importance in various fields of knowledge, including evolution and biogeography, and has direct relevance to conservation (Pinheiro et al. 2018). Insular environments are known to have a high rate of endemism, as certain evolutionary processes, such as inbreeding and loss of genetic variability, are more pronounced in limited populations disconnected from the continental environments (Roberts et al. 2002; Rocha 2003).

The Fernando de Noronha Archipelago (FNA) comprises an isolated group of volcanic islands located in the western equatorial Atlantic Ocean, ~350 km off the northeastern coast of Brazil (03°50'S, 032°25'W; Fig. 1). This insular region is under the influence of the South Equatorial Current, with an average water surface temperature of 26 °C and a tropical oceanic climate characterized by a rainy season during February–July and a dry season during the rest of the year (Maida and Ferreira 1997).




Figure 1. Geographical location of Fernando de Noronha Archipelago (FNA) and Saint Peter and Saint Paul's Archipelago (SPSPA) in the western equatorial Atlantic.

Most of the area around the FNA constitutes a marine national park where fishing is prohibited and tourist activities are regulated (Brasil 1988; Ibama 1990). The remaining area of the archipelago falls under the Fernando de Noronha Environmental Protection Area – Rocas – São Pedro and São Paulo, where fishing, land occupation, and tourist activities are allowed (Brasil 1986; ICMBio 2017).

The Saint Peter and Saint Paul's Archipelago (SPSPA) consists of 10 small rocky islets in Brazilian territorial waters and are located on the Mid-Atlantic Ridge ($00^{\circ}55'01''\text{N}$, $029^{\circ}20'76''\text{W}$; Fig. 1), ~1100 km from the northeastern coast of Brazil. The SPSPA is under the influence of the South Equatorial Current that flows westwards and the Equatorial Countercurrent that flows eastwards at a depth of 40–150 m (Edwards and Lubbock 1983).

In 2018, the Brazilian government declared the SPSPA an Environmental Protection Area, where fishing and the use of natural resources are permitted over $407,052 \text{ km}^2$ around the archipelago. SPSPA was also given the status as a marine natural marine monument,

an integral protection area with $47,263 \text{ km}^2$ of territorial sea (Brasil 2018).

The ichthyofauna of island environments is dynamic, undergoing colonization and extinction events over time (Mellin et al. 2010). Additionally, knowledge about the species diversity of remote oceanic islands is often incomplete, primarily due to difficulties in access and the inherent limitations of collection methods. The high degree of endemism of reef fish (Edwards and Lubbock 1983; Rocha 2003; Floeter et al. 2008; Hachich et al. 2015; Macieira et al. 2015; Pinheiro et al. 2018; Pimentel et al. 2020; Pinheiro et al. 2020), the presence of divergent genetic lineages (Rocha et al. 2005; Neves et al. 2016; Anderson et al. 2017), unusual behaviors (Luiz 2005; Gasparini et al. 2008), and the existence of populations with specific color morphs (Feitoza et al. 2003; Luiz 2003) reinforce the need for a more accurate investigation of the fish diversity in such locations.

The most complete and up-to-date information on ichthyofauna for the FNA reported the occurrence of 250 species (Pimentel et al. 2020), while for SPSPA, 225 species have been reported (Pinheiro et al. 2020). The

present study adds two unreported fish to the species inventories for these archipelagos, providing data for a better assessment of the biogeographic patterns of the oceanic islands of the western equatorial Atlantic.

Method

Specimens were collected with permission from the Chico Mendes Institute for Biodiversity Conservation (ICMBio 19135-8). The specimens were collected with a hand net, fixed in 10% formalin, transferred to 70% ethanol, and deposited in the fish collection of the Fisheries Biology Laboratory of the Federal University of Rio Grande do Norte (LABIPE). Morphometric measurements were performed using digital calipers with 1mm precision and the weight was measured with an electronic scale to the nearest 1 g. The individuals were identified, based on meristic and morphometric characters, following the diagnosis proposed by Berry and Vogele (1961) and McEachran and Fechhelm (1998).

Results

Cantherhines pullus (Ranzani, 1842)

Orangespotted Filefish

Figure 2

New record. BRAZIL • PE, Fernando de Noronha Archipelago, Praia do Sancho; 03°51'07"S, 032°26'38"W, depth of ~2 m from a rock bottom; 26.I.2018; R.X.

Soares, A.T. Borges and K.D.J. Amorim leg.; hand net; LABIPE 1257.

Identification. The individual was identified based on meristic and morphometric characteristics (Table 1) described in the diagnostic key by Berry and Vogele (1961) (Table 1).

Hemiramphus brasiliensis (Linnaeus, 1758)

Ballyhoo Halfbeak

Figure 3

New record. BRAZIL • Saint Peter and Saint Paul's Archipelago; 00°56'N, 029°18'W, ~5.5 km northeast of the main rocky islets of SPSPA; 13.I.2018; R.X. Soares, A.T. Borges and K.D.J. Amorim leg.; hand net; LABIPE 1256.

Identification. The individual was identified based on the diagnosis proposed by McEachran and Fechhelm (1998) after confirming the meristic and morphometric characters (Table 1).

Discussion

The existence of an ichthyofaunal species shared between both sides of the Atlantic and the SPSPA is likely a result of the dispersal, colonization, and larval settlement capacity of some species (Feitoza et al. 2003).

The documented distribution of *C. pullus* covers the East Atlantic from the Gulf of Guinea (Carpenter and De Angelis 2016a), including the islands of São Tomé and Príncipe (Wirtz et al 2007) and in the Western




Figure 2. *Cantherhines pullus*, LABIPE-1257, 141.8 mm SL. Scale bar = 30 mm.

Table 1. Morphometric and meristic characteristics of *Cantherhines pullus* and *Hemiramphus brasiliensis* specimens deposited in the fish collection of the LABIPE.

<i>Cantherhines pullus</i>	
Morphometric (mm)	LABIPE 1257
Total length	168
Standard length	141.8
Head length	43.2
Snout length	31.2
Pectoral length	15.9
Body depth	73.1
Eye diameter	10.5
Eye to dorsal spine distance	9.3
Meristic	
Dorsal rays	33
Anal rays	29
Pectoral rays	13
Weight (g)	95

<i>Hemiramphus brasiliensis</i>	
Morphometric (mm)	LABIPE 1256
Total length*	245.8
Standard length*	181.4
Distance between pectoral insertion to pelvic insertion	76.6
Distance between pelvic insertion to caudal base	57.1
Pectoral length	30.7
Body depth at pectoral origin	24.8
Body depth at pectoral tip	28
Meristic	
Dorsal rays	14
Anal rays	13
Pectoral rays	10
Pelvic rays	6
Caudal rays	20
Gill rakers	12+18
Weight (g)	102

* Beak partially broken

Atlantic from Massachusetts (USA), Gulf of Mexico and Caribbean Sea, to the Laje de Santos marine park, on the coast of the state of São Paulo in the southeastern Brazil (Menezes et al. 2007; Luiz et al. 2008; Fricke et al. 2021), including the Trindade and Martin Vaz Archipelago (Pereira-Filho et al. 2011). The occurrence of *C. pullus* in Bermuda (Carpenter 2002) and in the Trindade and Martin Vaz Archipelago represent the dispersive capacity of the species throughout its geographic distribution in remote island habitats. This may explain the extensive distribution of *C. pullus*, with certain oceanic islands serving as favorable habitats for settlement and

connectivity between the two sides of the Atlantic.

Hemiramphus brasiliensis is distributed in the Eastern Atlantic from Mauritania to Luanda (Angola), including the Cape Verde Islands (Carpenter and De Angelis 2016b). In the Western Atlantic, it is recorded from Massachusetts (USA), Gulf of Mexico, and Caribbean Sea to southeastern Brazil (Menezes et al. 2007; Fricke et al. 2021), including the FNA (Soto 2001) and the Trindade Archipelago and Martin Vaz (Gasparini and Floeter 2001). The occurrence of *H. brasiliensis* in the SPSPA, located ~1100 km from the northeast coast of Brazil and ~1700 km from the Cape Verde Islands, reinforce the role of this region as a support point to some amphi-Atlantic species.

Despite the common occurrence of endemic species (*Bodianus insularis* Eschmeyer, 1971, *Herpetoichthys regius* (Richardson, 1848), *Pontinus nigropunctatus* (Günther, 1868), *Scorpaenodes insularis* Eschmeyer, 1971) among Ascension Island, Saint Helena, and the SPSPA suggest a faunal link between these archipelagos, *H. brasiliensis* is reported exclusively for the last mid-Atlantic insular region (Wirtz et al. 2017; Brown et al. 2019). Thus, the consistent increase in the number of species recorded over time in the FNA and SPSPA demonstrates their biogeographic importance as stepping-stone habitats for some Atlantic fish species, and emphasizes the conservation importance of these island areas.

Acknowledgements

We are grateful to the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq - #442664/2015-0; #442626/2019-3), for the financial support and to the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for the doctoral fellowship granted to R.X. Soares. We also thank Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for the collection licenses (#19135-1, #19135-8).

Authors' Contributions

Conceptualization: RXS, JGJ. Data curation: JGJ. Formal analysis: ATB, JGJ. Funding acquisition: WFM. Investigation: RXS, KDJ, ATB. Methodology: RXS, KDJ. Project administration: WFM. Visualization: RXS, JGJ. Writing – original draft: RXS, KDJ, WFM, JGJ. Writing – review and editing: ATB, WFM, JGJ.



Figure 3. *Hemiramphus brasiliensis*, LABIPE-1256, 181.4 mm SL. Scale bar = 30mm.

References

- Anderson WD, Baldwin CC, Carvalho-Filho A, Júnior TV (2017) Re-description of the Jeweled Gemfish, *Anthias asperilinguis* (Serranidae: Anthiadinae), with comments on its ontogeny, phylogeny, and ecology. *Aqua, International Journal of Ichthyology* 23: 73–95.
- Brasil (1986) Decreto no 92.755, de 5 de junho de 1986. Diário Oficial da União, Poder Executivo, Brasília, Brazil. http://www.planalto.gov.br/ccivil_03/decreto/1980-1989/1985-1987/D92755.htm. Accessed on: 2021-08-01.
- Brasil (1988) Decreto no 96.693, de 14 de setembro de 1988. Diário Oficial da União, Poder Executivo, Brasília, Brazil. <https://www2.camara.leg.br/legin/fed/decret/1988/decreto-96693-14-setembro-1988-447461-publicacaooriginal-1-pe.html>. Accessed on: 2021-08-01.
- Brasil (2018) Decreto n. 9.313, de 19 de março de 2018. Cria a Área de Proteção Ambiental do Arquipélago de São Pedro e São Paulo e o Monumento Natural do Arquipélago de São Pedro e São Paulo. Diário Oficial da União, Brasília, Brazil. https://www.gov.br/icmbio/pt-br/assuntos/biodiversidade/unidade-de-conservacao-unidades-de-biomas/marinho/lista-de-ucs/area-de-protecao-ambiental-do-sao-pedro-e-sao-paulo/arquivos/decreto_9313_de_19mar2018_cria_apo_do_arquipelago_de_sao_pedro_e_sao_paulo.pdf. Accessed on: 2021-08-01.
- Brown J, Beard A, Clingham E, Fricke R, Henry L, Wirtz P (2019) The fishes of St Helena Island, central Atlantic Ocean—new records and an annotated check-list. *Zootaxa* 4543 (2): 151–194. <https://doi.org/10.11646/zootaxa.4543.2.1>
- Carpenter KE (2002) The living marine resources of the western central Atlantic. Volume 2: Bony fishes part 1 (Acipenseridae to Grammatidae). FAO Species Identification Guide for Fishery Purposes and American Society of Ichthyologists and Herpetologists Special Publication No. 5. FAO, Rome, Italy, 601–1374.
- Carpenter KE, De Angelis N (2016a) The living marine resources of the eastern central Atlantic. Volume 4: Bony fishes part 2 (Perciformes to Tetraodontiformes) and Sea turtles. FAO Species Identification Guide for Fishery Purposes. FAO, Rome, Italy, 2343–3124.
- Carpenter KE, De Angelis N (2016b) The living marine resources of the eastern central Atlantic. Volume 3: Bony fishes part 1 (Elopiformes to Scorpaeniformes). FAO Species Identification Guide for Fishery Purposes. FAO, Rome, Italy, 1511–2350.
- Edwards A, Lubbock R (1983) Marine zoogeography of St. Paul's Rocks. *Journal of Biogeography* 10: 65–72. <https://doi.org/10.2307/2844583>
- Feitoza BM, Rocha LA, Luis-Júnior OJ, Floeter SR, Gasparini JL (2003) Reef fishes of St. Paul's Rocks: new records and notes on biology and zoogeography. *Aqua, Journal of Ichthyology and Aquatic Biology* 7: 61–82.
- Floeter SR, Rocha LA, Robertson DR, Joyeux JC, Smith-Vaniz WF, Wirtz P, Edwards AJ, Barreiros JP, Ferreira CEL, Gasparini JL, Brito A, Falcón JM, Bowen BW, Bernardi G (2008) Atlantic reef fish biogeography and evolution. *Journal of Biogeography* 35: 22–47. <https://doi.org/10.1111/j.1365-2699.2007.01790.x>
- Fricke R, Eschmeyer WN, Van der Laan R (2021) Eschmeyer's catalog of fishes: genera, species, references. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. Accessed on 2021-06-18.
- Gasparini JL, Floeter SR (2001) The shore fishes of Trindade Island, western South Atlantic. *Journal of Natural History* 35 (11): 1639–1656. <https://doi.org/10.1080/002229301317092379>
- Gasparini J, Luiz O, Sazima I (2008) Cleaners from the underground. *Coral Reefs* 27: 143. <https://doi.org/10.1007/s00338-007-0326-7>
- Hachich NF, Bonsall MB, Arraut EM, Barneche DR, Lewinsohn TM, Floeter SR (2015) Island biogeography: patterns of marine shallow-water organisms in the Atlantic Ocean. *Journal of Biogeography* 42: 1871–1882. <https://doi.org/10.1111/jbi.12560>
- IBAMA (Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis) (1990) Plano de Manejo do Parque Nacional Marinho de Fernando de Noronha. IBAMA, Brasília, Brazil, 253 pp.
- ICMBio Instituto Chico Mendes de Conservação da Biodiversidade (2017) Plano de Manejo da Área de Proteção Ambiental de Fernando de Noronha – Rocas – São Pedro e São Paulo. ICMBio, Brasília, Brazil, 156 pp.
- Luiz OJ (2003) Colour morphs in a Queen angelfish *Holacanthus ciliaris* (Perciformes: Pomacanthidae) population of St. Paul's Rocks, NE Brazil. *Tropical Fish Hobbyist* 51: 82–90.
- Luiz OJ (2005) Unusual behavior of moray eels on an isolated tropical island (St. Paul's Rocks, Brazil). *Coral Reefs* 24: 501. <https://doi.org/10.1007/s00338-005-0500-8>
- Luiz OJ, Carvalho-Filho A, Ferreira CE, Floeter SR, Gasparini JL, Sazima I (2008) The reef fish assemblage of the Laje de Santos Marine State Park, Southwestern Atlantic: annotated checklist with comments on abundance, distribution, trophic structure, symbiotic associations, and conservation. *Zootaxa*, 1807: 1–25. <https://doi.org/10.11646/zootaxa.1807.1.1>
- Macieira RM, Simon T, Pimentel CR, Joyeux JC (2015) Isolation and speciation of tidepool fishes as a consequence of Quaternary sea-level fluctuations. *Environmental Biology Fishery* 98: 385–93. <https://doi.org/10.1007/s10641-014-0269-0>
- Maida M, Ferreira BP (1997) Coral reefs of Brazil: an overview. *Proceedings of the 8th International Coral Reef Symposium* 1: 263–274.
- McEachran JD, Fechhelm JD (1998) Fishes of the Gulf of Mexico. Vol. 1. University of Texas Press, Austin, USA, 1112 pp.
- Mellin C, Huchery C, Caley MJ, Meekan MG, Bradshaw CJA (2010) Reef size and isolation determine the temporal stability of coral reef fish populations. *Ecology* 91: 3138–3145. <https://doi.org/10.1890/10-0267.1>
- Neves JMM, Lima SMQ, Mendes LF, Torres RA, Pereira RJ, Mott T (2016) Population structure of the Rockpool Blenny *Entomacrodus vomerinus* shows source-sink dynamics among ecoregions in the tropical southwestern Atlantic. *PLoS ONE* 11: e0157472. <https://doi.org/10.1371/journal.pone.0157472>
- Pereira-Filho G, Amado-Filho G, Guimarães S, Moura RL, Sumida P, Abrantes DP, Bahia RG, Güth AZ, Jorge RR, Francini Filho RB (2011) Reef fish and benthic assemblages of the Trindade and Martin Vaz Island group, Southwestern Atlantic. *Brazilian Journal of Oceanography* 59: 201–212.
- Pimentel CR, Rocha LA, Shepherd B, Phelps TAY, Joyeux J-C, Martins AS, Stein CE, Teixeira JB, Gasparini JL, Reis-Filho JA, Garla RC, Francini-Filho RB, Delfino SDT, Mello TJ, Giarrizzo T, Pinheiro HT (2020) Mesophotic ecosystems at Fernando de Noronha Archipelago, Brazil (South-western Atlantic), reveal unique ichthyofauna and need for conservation. *Neotropical Ichthyology* 18 (4): e200050. <https://doi.org/10.1590/1982-0224-2020-0050>
- Pinheiro HT, Rocha C, Rocha LA (2018) *Tosanoides aphrodite*, a new species from mesophotic coral ecosystems of St. Paul's Rocks, Mid Atlantic Ridge (Perciformes, Serranidae, Anthiadinae). *ZooKeys* 786: 105–115. <https://doi.org/10.3897/zookeys.786.27382>
- Pinheiro HT, Macena BCL, Francini-Filho RB, Ferreira CEL, Albuquerque FV, Bezerra NPA, Carvalho-Filho A, Ferreira RCP, Luiz OJ, Mello TJ, Mendonça SA, Nunes DM, Pimentel CR, Pires AMA, Soares-Gomes A, Viana DL, Hazin FHV, Rocha LA (2020) Fish biodiversity of Saint Peter and Saint Paul's Archipelago, Mid-Atlantic Ridge, Brazil: new records and a species database. *Journal of Fish Biology* 97 (4): 1143–1153. <https://doi.org/10.1111/jfb.14484>
- Roberts CM, McClean CJ, Veron JEN, Hawkins JP, Allen GR, McAlister DE, Mittermeier CG, Schueler FW, Spalding M, Wells F, Vynne C, Werner TB (2002) Science marine biodiversity hotspots and conservation priorities for tropical reefs. *Science* 295 (5558): 1280–1284. <https://doi.org/10.1126/science.1067728>
- Rocha LA (2003) Patterns of distribution and processes of speciation

- in Brazilian reef fishes. *Journal of Biogeography* 30: 1161–1171. <https://doi.org/10.1046/j.1365-2699.2003.00900.x>
- Rocha LA, Robertson DR, Roman J, Bowen BW (2005) Ecological speciation in tropical reef fishes. *Proceedings of the Royal Society B: Biological Sciences* 272: 573–579. <https://doi.org/10.1098/rspb.2004.3005>
- Soto JMR (2001) Peixes do Arquipélago Fernando de Noronha. *Mare Magnum* 1: 147–169.
- Wirtz P, Bingeman J, Bingeman J, Fricke R, Hook T, Young J (2017). The fishes of Ascension Island, central Atlantic Ocean – new records and an annotated checklist. *Journal of the Marine Biological Association of the United Kingdom* 97 (4): 783–798. <https://doi.org/10.1017/S0025315414001301>
- Wirtz P, Ferreira C, Floeter S, Fricke R, Gasparini J, Iwamoto T, Rocha L, Sampaio C, Schliewen KU (2007) Coastal Fishes of São Tomé And Príncipe Islands, Gulf of Guinea (Eastern Atlantic Ocean) - An Update. *Zootaxa* 1523: 1–48. <https://doi.org/10.5281/zenodo.177470>