

# Fishes of the Salonga National Park (Congo basin, central Africa): A list of species collected in the Luilaka, Salonga, and Yenge Rivers (Equateur Province, Democratic Republic of Congo)

Raoul J.C. Monsembula Iyaba<sup>1</sup> and Melanie L.J. Stiassny<sup>2\*</sup>

<sup>1</sup> Université de Kinshasa, Faculté des Sciences, Département de Biologie, B.P 190 Kin XI, Democratic Republic of Congo.

<sup>2</sup> American Museum of Natural History, Department of Ichthyology, Central Park West at 79th Street, New York, NY 10024, USA.

\* Corresponding author e-mail: [mljs@amnh.org](mailto:mljs@amnh.org)

**ABSTRACT:** A list of fishes collected in and around the periphery of the Salonga National Park in Equateur Province of the Democratic Republic of Congo is provided. While noteworthy for a wide array of rare and endangered terrestrial vertebrates, the fish fauna of the Salonga National Park, the largest protected area in Africa, is very poorly known. One hundred and fifty two species distributed in 24 families are reported on here; of which 9 represent putatively undescribed taxa currently under investigation. Lack of up-to-date taxonomic and distributional knowledge of African fishes is particularly acute in the central Congo basin and the present report, while undoubtedly incomplete, provides a foundation for much needed further work in the region.

## INTRODUCTION

Created by presidential decree in 1970, the Salonga National Park (SNP) is the largest tropical rainforest preserve in Africa (UNESCO 2012). Covering an area of ca. 36,000 km<sup>2</sup>, the SNP is entirely located within the sedimentary “cuvette centrale” of the Congo basin (Evrard 1968), where it is divided into two large forest blocks, each delimited by major waterways (Figure 1). Most of the park is situated about 300 meters above sea level with more-or-less continuous low relief cut by broad, meandering rivers with extensive seasonally flooded, or permanently inundated, riparian zones. The climate is equatorial, hot and humid, with an average annual temperature of 25.5°C, and rainfall of around 1800-2200 mm per year (Hughes and Hughes 1992). The densely forested landscape (Figure 2A) encompasses a large and complex aquatic network that traverses much of the terrestrial area of the park. Despite the high importance of the region's fishes for food security and to the local economy, most of the SNP remains remarkably poorly documented ichthyologically. This is undoubtedly a serious impediment for fisheries management in the region, where increasing fishing pressure, and the use of highly destructive fishing techniques is of growing concern both outside and inside the park's boundaries (Inogwabini 2005; pers obs.).

For fishes, a single reference (Inogwabini 2005) reports on 56 species collected at two sites in the SNP; one in the northern block at Etate on the Salonga River, and the second in the southern block at Lokofa on the Luilaka River. Historical literature for the area is equally sparse. Matthes (1964) reported on a series of collections from around Ikela (a settlement on the Tshuapa River some 270 km west of Etate; Figure 1) and that work remains the primary published reference for the entire region. However, a recent assessment of the status and

distribution of fishes throughout central Africa (Stiassny *et al.* 2011) provides useful comparative data. That IUCN assessment compiled point data from legacy collections housed in major museums (Darwell and Smith 2011), and in particular the large historical holdings of fishes from collections made around Mbandaka, Boende, and Ikela (Figure 1) housed at the Africa Museum, Tervuren, and these provide the best available comparative data for fishes in the vicinity of the SNP.

## MATERIALS AND METHODS

The list of species provided here is based on collections made during three field expeditions to the SNP and adjacent regions: the first, from July 3-August 3, 2006 (during the main low water season), the second from October 28-November 17, 2007 (during the primary flood season), and the third between May 19-May 28, 2010 (towards the end of the secondary flood season). Transportation within the SNP is restricted to passage along waterways in local pirogues. Due to difficulties of access, collecting was primarily restricted to main channel sites and only a few collections were made in peripheral habitats, or in the many small tributaries and affluent streams. Sites were selected to sample accessible regions of the main river channels, both inside and outside of the SNP, and subjected to varying degrees of anthropogenic impact, from pristine (e.g., Figure 2B) to marginally degraded and heavily fished (e.g., Figure 2C). Fifteen sites along a total stretch of approximately 650 km of the Luilaka, Salonga and Yenge Rivers are reported on (Table 1, Figure 1).

Fishes were collected and euthanized prior to preservation in accordance with recommended guidelines for the use of fishes in research (AFS/AIFRB/ASIH 2003). All materials are housed in the Ichthyology Department of the American Museum of Natural History (AMNH), New

York with a database accessible at <http://entheros.amnh.org/db/emuwebamnh/>.

#### Notes on Collection Sites

Sampled sites share similar physical characteristics, being situated along the banks of the heavily forested Salonga, Yenge, and Luilaka Rivers, with most in seemingly pristine condition. The waters of all three rivers are tea-colored, highly humic, and notably acidic, with low dissolved oxygen and conductivity (Table 2). The Luilaka River, which delimits the northern boundary of the SNP southern block, has an average width of 45 m, varying from 30–60 m between Ifumu (Site 1) and Ikala (Site 9). Depths at collection sites range between 2–10 m (depending on season), the substrate is predominately fine mud, sand, and detritus, with strong flow in the main channel, particularly during the dry season when waters are restricted within the river channel. At its maximum extent, the zone of inundation rarely reaches more than 100 m into the forest on either side of the main channel. The Salonga River has

an average width of 60 m, but varies from 40–80 m between Etate (Site 14) and Bomputu (Site 15). Depths at collection sites varies from 3–11 m (depending on season), again with water flow strongest in the main channel during the dry season. The substrate consists primarily of sand, fine mud and detritus, with many large sand bars exposed in the dry season. During the height of the rainy season a zone of inundation of about 100 m extends into the forests on either side of the main channel. The Yenge River, a large left bank tributary of the Salonga, is located entirely within the SNP. Although the smallest of the three rivers sampled in terms of channel width, the zone of inundation of the Yenge is by far the greatest, with waters reaching upwards of 400 m into the forest during the height of the rains. Site 3 (Boangi, Figure 2C), Site 11 (Efefa, Figure 2F), and Site 15 (Bomputu, Figure 2H) are located near village docks, with evidence of forest clearing (mainly for housing and fishing hut construction), and with grassy banks replacing riparian vegetation.

FIGURE 1. Coordinates and site data.

SITE	LOCALITY	COORDINATES	MAIN RIVER CHANNEL	COVER
1	Ifumu	1°47'31" S, 20°41'18" E	Luilaka	Dense forest, undisturbed
2	Bekongo	1°54'42" S, 20°51'11" E	Luilaka	Dense forest, undisturbed
3	Boangi	1°55'19" S, 20°52'56" E	Luilaka	Dense forest, some localized disturbance
4	Kema Asondji	2°2'12" S, 20°59'48" E	Luilaka	Dense forest, undisturbed
5	Bosombangwa	2°13'27" S, 2°11'6" E	Luilaka	Dense forest, undisturbed
6	Ikala (confluence with Lofombo)	2°36'47" S, 21°13'32" E	Luilaka	Dense forest, undisturbed
7	Mboko Mboko	2°30'45" S, 21°22'1" E	Luilaka	Dense forest, undisturbed
8	Ilenge	2°35'47" S, 21°34'36" E	Luilaka	Dense forest, undisturbed
9	Komba Dumbe	2°40'15" S, 21°43'15" E	Luilaka	Dense forest, undisturbed
10	Watshikengo	0°54'45" S, 2°32'15" E	Salonga	Dense forest, undisturbed
11	Efefa	1°1'48" S, 20°43'41" E	Yenge/Salonga confluence	Dense forest, some localized disturbance
12	Yenge 1	1°2'45" S, 20°43'59" E	Yenge	Dense forest, undisturbed
13	Boyenga	1°3'50" S, 20°44'44" E	Yenge	Dense forest, undisturbed
14	Etate	1°3'18" S, 20°48'38" E	Salonga	Dense forest, undisturbed
15	Bomputu	0°28'5" S, 20°8'32" E	Salonga	Dense forest, some urbanization and disturbance




FIGURE 1. The Salonga National Park (shaded grey) and associated drainages in Central Africa, with the locations of Sites 1–15 indicated.




### Fish sampling

Standard fishing techniques were employed (Lang and Baldwin 1996), and included deployment of monofilament gill nets, traps, dip nets, cast nets, hook and line, and locally made barrage traps (Figure 2H). Additional specimens were purchased from fishermen along the river and at village markets.

Taxonomic nomenclature used herein is based primarily on Eschmeyer (2012) but with a few modifications

to reflect recent taxonomic changes. All fishes were collected and exported with permission of the République Démocratique du Congo, Ministère de l'Environnement, Conservation de la Nature, Eaux et Forêts, Direction des Ressources Faunistiques et Chasse (Permit #'s 004/SG/ECN-EF/DRFC/06 and 002/SG/ECN-EF/DRFC/08), and the Ministère de l'Agriculture, Secrétariat Général à l'Agriculture, Pêche et Elevage, Direction des Pêches (Permit # 003/2010), which are on file at AMNH.



**FIGURE 2.** Selected localities: A) View of SNP from the air near Site 1, Ifumu; B) Site 2, Bekongo; C) Site 3, Boangi; D) Site 4, Kema Asondji; E) Site 7, Mboko Mboko; F) Site 11, Efofa; G) Site 14, Etate; H) Site 15, Bomputu.



## RESULTS AND DISCUSSION

A total of 2582 individuals belonging to 152 species, distributed in 24 families and 8 orders are represented in Table 2, with their distribution among sites indicated in columns 1–15. In column 16, distribution data from an IUCN assessment of the status and distribution of central African fishes (Brooks *et al.* 2011) are used to indicate which SNP species have previously been known to occur in the vicinity of the park. While comparative data are minimal for the rivers of the SNP itself, Brooks *et al.* (2011) do provide data on the region's fishes, mostly with historical records from around Mbandaka on the Ruki River, and Boende and Ikela on the Tshuapa River, thereby providing the best available comparison for the SNP list provided here.

Based on these data we recognize 33 species that have not previously been recorded from the central cuvette of the Congo basin (exclusive of the main channel of the Congo River), or are known only from isolated regions distant from the rivers of the SNP. Most of these species (Table 3, column 16) represent range extensions within the Congo basin, and a selection of them is illustrated in Figures 3–5. While the present study has significantly increased the number of species known to occur within the boundaries of the SNP, with few exceptions, the species reported on here are widespread in the region and many occur throughout much of the rain forested central Congo basin. Regional endemism is seemingly low, possibly due to the extent of currently interconnected, and densely forested habitat throughout most of the cuvette (Chapman 2001), or as a result of repeated climatic shifts resulting in fluctuating forest contraction and expansion since the Middle Pleistocene (Campbell 2005). Despite the predominance of widespread species in our collections, a number of specimens collected during the course of the present study are not assignable to currently described taxa, and appear to represent undescribed species, indicating that significant regional endemism may remain to be discovered. A selection of these is illustrated in Figures 6–7, and includes a specimen of psammophilous, dorso-ventrally depressed, and depigmented clarioteid catfish (Figure 6A), that belongs to a new genus endemic to the Congo basin (Sullivan, pers. comm.). It is noteworthy that, with the exception of the *Alestopetersius* (Figure 6B) and *Eugnathichthys* (Figure 6C) species, most of the specimens unassignable to known species are of small size (Figure 7), and were collected in marginal habitats rather than in the main river channels. As noted previously, most collecting effort was focused in the main river channels and sampling in the many small affluent streams, black water swamps, and peripheral habitats was minimal. Therefore, while the present study represents the best available tally for the SNP and its immediate vicinity, targeted collecting in these marginal habitats will be necessary to provide a fully comprehensive listing for the park, and an accurate assessment of the actual levels of endemism in the region.

TABLE 2. Water chemistry data.

PARAMETER (AVERAGED OVER SITES)	SALONGA RIVER SITES	YENGE RIVER SITES	LUILAKA RIVER SITES
Air T° C	25	25	24.8
Water surface T° C	24	23.6	23.8
Turbidity (FTU)	13	11.4	12.3
Transparency (cm)	60	55	58
Depth at site (m)	7.2	5.9	6.5
pH	4.8	4.8	4.9
Conductivity (µs/cm)	26	22	24
O <sub>2</sub> (mg/l)	2.1	2.4	2.8
K <sup>+</sup> (mg/l)	0.28	0.3	0.3
Ca <sup>2+</sup> (mg/l)	0.22	0.21	0.2
Mg <sup>2+</sup> (mg/l)	0.16	0.22	0.21
Cl <sup>-</sup> (mg/l)	1.4	2.4	1.6
SO <sub>4</sub> <sup>2-</sup> (mg/l)	1.1	1.2	1.0
NO <sub>3</sub> <sup>-</sup> (mg/l)	0.4	0.8	0.5
PO <sub>4</sub> <sup>3-</sup> (mg/l)	1.25	1.7	1.5







FIGURE 3. Species whose occurrence in the SNP represents a significant range extension within the Congo basin: A) *Potamothrissa obtusirostris*, B) *Cyphomyrus weeksii*, C) *Mormyrus caballus*, D) *Petrocephalus odzalaensis*, E) *Petrocephalus zakoni*, F) *Pollimyrus maculipinnis*. Scale bars: 1 cm.



**FIGURE 4.** Species whose occurrence in the SNP represents a significant range extension within the Congo basin: A) *Clupeocharax schoutedeni*, B) *Hemigrammocharax uniozellatus*, C) *Nannocharax elongatus*, D) *Hemigrammocharax ocellicauda*, E) *Clypeobarbus bomokandi*. Scale bars: 1 cm.



**FIGURE 5.** Species whose occurrence in the SNP represents a significant range extension within the Congo basin: A) *Clariallabes teugelsi*, B) *Clariallabes variabilis*, C) *Phractura lindica*, D) *Ctenopoma gabonense*, E) *Kribia uellensis*. Scale bars: 1 cm.



**FIGURE 6.** Undescribed species from the region: A) "Salonga glass catfish" (Claroteidae, new genus and species), in lateral and dorsal views, B) *Alestopetersius* sp., C) *Eugnathichthys* sp. Scale bars: 1 cm.



**FIGURE 7.** Undescribed species from the region: A) *Microsynodontis* sp., B) *Congochromis* sp., male above, female, C) *Hylopanchax* sp., D) *Fenerbahce* sp. Scale bars: 1 cm.

**TABLE 3.** List of species collected at sites 1-15 along the Lulaka, Salonga and Yenge Rivers (Salonga National Park). In column 16 distribution data culled from an IUCN assessment of central African fishes (Brooks *et al.* 2011) indicates which of the reported SNP species have previously been known to occur in the region.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<b>LEPIDOSIRENIFORMES</b>																
Protopteridae (1)																
<i>Protopterus dolloi</i> Boulenger, 1900	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<b>POLYPTERIFORMES</b>																
Polypteridae (4)																
<i>Polypterus delhezi</i> Boulenger, 1899	-	-	-	-	-	-	-	-	-	-	-	-	x	-	-	x
<i>Polypterus retropinnis</i> Vaillant, 1899	x	-	-	-	-	-	x	-	-	-	-	-	-	-	-	x
<i>Polypterus weeksii</i> Boulenger, 1898	x	-	-	x	-	-	-	-	-	-	-	-	-	-	-	x
<i>Polypterus ornatipinnis</i> Boulenger, 1902	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<b>CLUPEIFORMES</b>																
Clupeidae (1)																
<i>Potamothrissa obtusirostris</i> (Boulenger, 1909)	x	-	-	x	-	x	-	-	x	x	-	x	x	-	-	-
<b>OSTEOGLOSSIFORMES</b>																
Pantodontidae (1)																
<i>Pantodon buchholzi</i> Peters, 1877	x	x	x	-	-	-	-	-	x	-	-	x	x	-	-	x
<b>Notopteridae (2)</b>																
<i>Papyrocranus congoensis</i> (Nichols and La Monte, 1932)	-	-	-	-	-	-	-	x	-	-	x	-	x	-	-	x
<i>Xenomystus nigri</i> (Günther, 1868)	-	x	x	x	-	-	x	x	-	-	-	x	x	x	-	x
<b>Mormyridae (28)</b>																
<i>Campylomormyrus tamandua</i> (Günther, 1864)	-	-	-	-	-	-	x	-	-	-	-	-	-	-	-	x
<i>Cyphomorphus weeksii</i> (Boulenger, 1902)	-	-	-	-	-	-	x	x	-	-	-	-	-	-	-	-
<i>Cyphomorphus ghesquieri</i> (Poll, 1945)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Genyomorphus donnyi</i> Boulenger, 1898	-	-	-	-	-	-	x	-	-	-	-	-	-	-	-	x
<i>Gnathonemus echidnorhynchus</i> Pellegrin, 1924	-	-	-	-	-	-	-	x	-	-	-	x	x	-	-	x
<i>Gnathonemus petersii</i> (Günther, 1862)	-	x	-	x	x	x	x	x	-	-	-	-	-	x	-	x
<i>Marcusenius friteli</i> (Pellegrin, 1904)	x	-	-	-	-	x	-	-	-	-	-	-	-	-	-	x
<i>Marcusenius greshoffii</i> (Schilthuis, 1891)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Marcusenius intermedius</i> Pellegrin, 1924	-	-	x	x	-	-	-	-	x	-	-	x	-	x	-	-
<i>Marcusenius kutuensis</i> (Boulenger, 1899)	-	x	-	x	-	x	x	-	-	-	-	x	-	-	-	x
<i>Marcusenius moorii</i> (Günther, 1867)	-	-	x	-	x	-	-	-	-	-	-	-	x	-	-	x
<i>Mormyrops anguilloides</i> (Linnaeus, 1758)	x	-	-	-	x	-	-	-	-	-	-	-	-	-	x	x
<i>Mormyrops attenuatus</i> Boulenger, 1898	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Mormyrops nigricans</i> Boulenger, 1899	-	x	-	-	x	-	-	-	-	-	-	x	-	-	-	x
<i>Mormyrus caballus</i> Boulenger, 1898	x	-	-	-	x	-	-	-	-	-	-	-	-	-	-	-
<i>Oxymormyrus boulengeri</i> (Pellegrin, 1900)	-	-	-	x	x	-	x	-	-	-	-	-	-	-	-	x
<i>Petrocephalus balayi</i> Sauvage, 1883	-	-	-	x	-	x	x	-	-	-	-	-	-	-	-	x
<i>Petrocephalus binotatus</i> Pellegrin, 1924	-	-	-	x	-	-	x	x	x	-	-	-	-	x	-	x
<i>Petrocephalus odzalaensis</i> Lavoué <i>et al.</i> , 2010	x	-	x	x	-	x	x	x	-	-	-	x	-	-	-	-
<i>Petrocephalus zakoni</i> Lavoué <i>et al.</i> , 2010	x	-	x	-	-	x	x	x	-	-	-	x	x	-	-	-



TABLE 3. CONTINUED.

	AMNH Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<i>Pollimyrus osbornii</i> (Nichols and Griscom, 1917)	AMNH 244118	-	-	-	-	-	-	-	-	-	-	-	-	X	-	-	X
<i>Pollimyrus maculipinnis</i> (Nicholas and La Monte, 1934)	AMNH 244119	-	-	-	-	-	X	-	-	-	-	-	-	-	-	-	-
<i>Pollimyrus nigripinnis</i> (Boulenger, 1899)	AMNH 241899, 241900	-	-	X	X	-	-	-	-	-	-	-	-	-	-	-	X
<i>Pollimyrus schreyeni</i> Poll, 1972	AMNH 242746, 241897	-	-	X	X	-	-	X	-	X	-	-	-	-	-	-	X
<i>Stomatorhinus ater</i> Pellegrin, 1924	AMNH 244116	-	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-
<i>Stomatorhinus humilior</i> Boulenger, 1899	AMNH 252201	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-
<i>Stomatorhinus kununguensis</i> Poll, 1945	AMNH 243545, 252214	-	-	-	-	-	-	-	X	X	-	-	-	-	-	-	X
<i>Stomatorhinus patrizii</i> Vinciguerra, 1928	AMNH 244117	-	-	-	-	-	-	X	-	-	-	-	-	-	-	-	X
GONORYNCHIFORMES																	
Phractolaemidae (1)																	
<i>Phractolaemus ansorgii</i> Boulenger, 1901	AMNH 252190, 252227	X	-	-	-	-	-	-	X	-	-	-	-	X	-	-	X
CHARACIFORMES																	
Hepsetidae (1)																	
<i>Hepsetus odoe</i> (Bloch, 1794)	AMNH 240821, 252244	X	-	-	X	X	X	X	X	-	X	-	-	-	-	X	X
Alestidae (21)																	
<i>Alestes liebrechtsii</i> Boulenger, 1898	AMNH 240753, 244187	X	X	-	X	-	-	X	X	X	-	-	-	-	-	-	X
<i>Alestopetersius bifasciatus</i> (Poll, 1967)	AMNH 244034	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	X
<i>Alestopetersius hilgendorfi</i> (Boulenger, 1899)	AMNH 240791, 244114	-	-	-	X	-	-	X	X	-	-	-	X	X	X	X	X
<i>Alestopetersius</i> sp.	AMNH 252852, 25251	-	-	-	-	-	-	X	-	-	-	-	-	-	-	X	-
<i>Bathyaethiops greeni</i> Fowler, 1949	AMNH 240800, 252261	X	-	X	-	X	X	X	-	-	X	-	-	-	-	X	X
<i>Brachypetersius altus</i> (Boulenger, 1899)	AMNH 240485, 244030	-	-	-	X	-	-	X	X	X	-	-	-	-	-	X	X
<i>Brycinus bimaculatus</i> (Boulenger, 1899)	AMNH 240759, 240762	-	-	X	X	-	-	X	X	-	-	-	-	-	-	-	X
<i>Brycinus grandisquamis</i> (Boulenger, 1899)	AMNH 240763, 244112	X	-	-	X	-	-	X	X	-	-	-	-	-	-	-	X
<i>Brycinus poptae</i> (Pellegrin, 1906)	AMNH 240766, 240768	X	-	-	-	-	-	-	-	-	-	-	-	X	-	-	X
<i>Bryconaeethiops boulengeri</i> Pellegrin, 1900	AMNH 240771, 244051	X	X	X	X	-	-	X	X	X	X	-	-	X	-	-	X
<i>Bryconaeethiops microstoma</i> Günther, 1873	AMNH 240769, 240770	-	-	-	X	-	-	-	-	X	-	-	-	-	-	-	X
<i>Clupeocharax schoutedeni</i> Pellegrin, 1926	AMNH 243540	-	-	-	-	-	-	-	-	-	X	-	-	-	-	-	-
<i>Hydrocynus goliath</i> Boulenger, 1898	observed	X	X	-	-	-	-	-	-	-	-	-	-	-	-	X	-
<i>Hydrocynus vittatus</i> Castelnan 1861	observed	X	X	X	X	-	-	-	-	-	-	-	-	-	-	X	X
<i>Micralestes congicus</i> Poll, 1967	AMNH 244027, 252259	X	-	-	-	X	-	-	-	-	-	-	-	-	-	-	X
<i>Micralestes humilis</i> Boulenger, 1899	AMNH 240811, 240815	-	-	X	X	-	-	X	X	X	-	-	-	-	-	-	X
<i>Phenacogrammus aurantiacus</i> (Pellegrin, 1930)	AMNH 240808, 240809	-	-	-	-	-	-	-	X	X	-	-	-	-	-	-	X
<i>Phenacogrammus deheynei</i> Poll, 1945	AMNH 244045, 252251	-	-	X	-	X	X	-	-	X	-	-	-	-	-	-	X
<i>Phenacogrammus interruptus</i> (Boulenger, 1899)	AMNH 240752, 247923	X	X	X	X	-	X	X	X	X	-	-	X	-	-	-	X
<i>Phenacogrammus polli</i> Lambert, 1961	AMNH 240793, 244044	X	X	X	X	-	-	-	-	X	X	-	-	-	-	-	X
<i>Tricuspidalestes caeruleus</i> (Matthes, 1964)	AMNH 252193	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	X
Citharinidae (2)																	
<i>Citharinus gibbosus</i> Boulenger, 1899	AMNH 244111	-	-	-	-	-	-	-	-	-	X	-	-	-	-	-	X
<i>Citharinus macrolepis</i> Boulenger, 1899	observed	-	-	-	-	-	-	-	-	-	-	X	-	-	-	-	X

TABLE 3. CONTINUED.

	AMNH Number	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<b>Distichodontidae (15)</b>																	
<i>Belonophago hutschboudi</i> Giltay, 1929	AMNH 241850, 241853	-	-	X	X	-	-	-	-	-	X	-	-	-	-	-	X
<i>Distichodus decemmaculatus</i> Pellegrin, 1926	AMNH 241855, 252263	X	-	-	-	X	-	X	X	-	X	-	-	-	-	-	X
<i>Distichodus fasciolatus</i> Boulenger, 1898	observed	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-	X
<i>Distichodus noboli</i> Boulenger, 1899	AMNH 241862, 247930	X	-	-	X	-	-	-	X	-	-	X	X	-	X	-	X
<i>Distichodus sexfasciatus</i> Boulenger, 1897	AMNH 241854	-	-	-	-	-	-	-	-	-	-	-	X	-	-	-	X
<i>Eugnathichthys</i> sp.	AMNH 241647, 241650	-	-	-	-	-	-	-	-	X	-	-	X	-	-	-	-
<i>Hemigrammocharax ocellicauda</i> (Boulenger, 1907)	AMNH 243547, 241886	X	-	-	-	-	-	-	X	X	-	-	-	-	-	-	X
<i>Hemigrammocharax unioellatus</i> (Pellegrin, 1926)	AMNH 247007	-	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-
<i>Mesoborus crocodilus</i> Pellegrin, 1900	AMNH 241669, 252242	X	-	X	X	X	X	X	X	-	-	-	X	X	-	X	X
<i>Microstomatichthysborus bashforddeani</i> Nichols and Griscom, 1917	AMNH 244115, 257162	-	-	-	X	-	-	X	-	-	-	-	-	-	-	-	-
<i>Nannocharax elongatus</i> Boulenger, 1900	AMNH 247935, 241895	X	-	-	-	-	-	-	-	-	-	-	X	-	-	-	-
<i>Nannocharax schoutedeni</i> Poll, 1939	AMNH 241889, 247932	X	-	X	-	-	-	-	-	X	-	-	-	X	-	-	X
<i>Neolebias gracilis</i> Matthes, 1964	AMNH 244084, 247928	-	-	X	-	-	-	-	-	-	-	-	-	-	-	-	X
<i>Neolebias trilineatus</i> Boulenger, 1899	AMNH 244084, 247928	-	X	X	X	-	X	-	-	X	-	-	X	-	-	-	X
<i>Xenocharax crassus</i> Pellegrin, 1900	AMNH 241651, 244024	X	X	X	X	X	X	X	X	X	X	X	-	-	X	-	-
<b>CYPRINIFORMES</b>																	
<b>Cyprinidae (4)</b>																	
<i>Barbus humeralis</i> Boulenger, 1902	AMNH 240446, 247929	-	-	-	X	-	-	X	-	-	-	-	-	-	-	-	X
<i>Barbus matthesi</i> Poll and Gosse, 1963	AMNH 240447	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	X
<i>Barbus miolepis</i> Boulenger, 1902	AMNH 239926, 252248	X	X	-	X	X	-	-	X	-	X	-	-	-	-	-	X
<i>Glypeobarbus bomakandi</i> (Myers, 1924)	AMNH 244090, 244093	-	-	-	X	-	-	X	X	-	-	-	-	-	-	-	-
<b>SILURIFORMES</b>																	
<b>Clariidae (9)</b>																	
<i>Anaspidoglanis macrostoma</i> (Pellegrin, 1909)	AMNH 244142, 252225	X	-	-	-	-	-	X	-	-	-	-	-	-	-	-	X
<i>Auchenoglanis wittei</i> Giltay, 1930	AMNH 241988	X	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Chrysichthys cranchii</i> (Leach, 1818)	AMNH 244177, 244179	-	-	-	-	-	-	X	X	-	-	-	-	-	-	-	X
<i>Chrysichthys delhezi</i> Boulenger, 1899	AMNH 247882	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	X
<i>Chrysichthys habereri</i> Steindachner, 1912	AMNH 244181	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-	X
<i>Chrysichthys ornatus</i> Boulenger, 1902	AMNH 239875, 241992	-	-	-	-	-	X	X	X	-	X	-	-	-	X	-	X
<i>Chrysichthys punctatus</i> Boulenger, 1899	AMNH 239246, 241990	X	-	X	X	-	X	X	X	-	-	-	-	-	X	-	X
Salonga "glass catfish" new genus and species	AMNH 239657	-	-	-	-	-	-	-	-	X	-	-	-	-	-	-	-
<i>Parauchenoglanis punctatus</i> (Boulenger, 1902)	AMNH 241987, 253357	X	-	-	-	X	-	-	-	-	-	-	-	X	-	-	X
<b>Schilbeidae (5)</b>																	
<i>Pareutropius debauwi</i> (Boulenger, 1900)	AMNH 241970, 244145	X	-	X	X	-	-	X	-	X	-	-	X	-	X	-	X
<i>Pareutropius mandevillei</i> Poll, 1959	AMNH 244146, 244147	X	-	-	-	-	-	-	-	-	-	-	X	-	-	-	X
<i>Schilbe grenfelli</i> (Boulenger, 1900)	AMNH 241958, 247881	-	-	-	-	-	-	-	X	-	X	-	-	-	-	-	X
<i>Schilbe laticeps</i> (Boulenger, 1899)	AMNH 241957, 244154	-	-	-	-	-	-	-	-	-	-	-	X	X	-	-	X
<i>Schilbe marmoratus</i> Boulenger, 1911	AMNH 241695, 252250	X	X	X	X	X	X	X	X	X	X	-	X	X	X	-	X



TABLE 3. CONTINUED.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<b>Clariidae (11)</b>																
<i>Channallabes apus</i> (Günther, 1873)	-	-	x	-	-	-	-	x	-	-	-	-	-	-	-	x
<i>Clariallabes centralis</i> (Poll and Lambert, 1958)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Clariallabes teugelsi</i> Ferraris, 2007	-	-	-	-	-	-	-	x	-	-	-	-	-	-	-	-
<i>Clariallabes variabilis</i> Pellegrin, 1926	-	-	-	-	-	-	x	x	x	-	-	-	-	-	-	-
<i>Clarias angolensis</i> Steindachner, 1866	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Clarias buthupogon</i> Sauvage, 1879	x	x	x	-	-	-	-	x	-	x	-	x	-	-	-	x
<i>Clarias gabonensis</i> Günther, 1867	-	-	x	-	-	x	x	x	-	x	-	-	-	-	-	x
<i>Clarias pachynema</i> Boulenger, 1903	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Clarias platycephalus</i> Boulenger, 1902	-	-	-	-	-	-	-	x	-	-	-	-	-	-	-	x
<i>Clarias theodora</i> Weber, 1897	-	-	-	-	-	-	-	x	-	-	-	-	x	-	-	-
<i>Heterobranchius longifilis</i> Valenciennes, 1840	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<b>Malapteruridae (3)</b>																
<i>Malapterurus melanocheir</i> Norris, 2002	-	-	-	-	-	-	-	-	-	-	x	-	-	-	-	x
<i>Paradoxoglanis caudivittatus</i> Norris, 2002	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Paradoxoglanis parvus</i> Norris, 2002	-	-	-	-	-	-	-	x	-	-	-	-	-	-	-	x
<b>Mochokidae (8)</b>																
<i>Microsynodontis</i> sp.	x	-	-	-	-	-	-	-	x	-	-	-	-	-	-	-
<i>Synodontis aterrima</i> Poll and Roberts, 1968	-	-	-	-	-	-	-	-	x	-	-	-	-	-	-	x
<i>Synodontis contracta</i> Vinciguerra, 1928	x	-	-	x	-	-	x	-	x	-	-	-	-	-	-	x
<i>Synodontis decora</i> Boulenger, 1899	x	-	-	x	-	-	-	-	-	-	-	-	-	-	-	x
<i>Synodontis flavitaeniata</i> Boulenger, 1919	-	x	x	x	-	x	x	x	-	-	-	-	x	-	-	x
<i>Synodontis greshoffi</i> Schilthuis 1891	-	x	-	x	-	-	-	-	-	-	-	-	x	-	-	x
<i>Synodontis nigriventris</i> David, 1936	x	-	x	x	-	-	-	x	-	-	-	x	-	-	-	x
<i>Synodontis schoutedeni</i> David, 1936	x	-	-	-	-	-	-	-	-	-	-	-	x	-	-	x
<b>Amphiliidae (3)</b>																
<i>Belonoglanis tenuis</i> Boulenger, 1902	x	-	-	-	-	x	-	-	x	-	-	-	-	-	-	x
<i>Phractura lindica</i> Boulenger, 1902	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Phractura scaphyrhychura</i> (Vaillant, 1886)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<b>CYPRINODONTIFORMES</b>																
<b>Nothobranchiidae (4)</b>																
<i>Aphyosemion</i> sp.	-	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Epiplatys chevalieri</i> (Boulenger, 1913)	x	-	-	x	-	-	-	x	x	-	-	-	x	-	-	x
<i>Epiplatys multifasciatus</i> (Boulenger, 1913)	x	-	-	-	-	-	-	-	x	-	-	-	-	x	-	x
<i>Fenerbahce</i> sp.	-	-	-	-	-	-	-	-	-	-	-	x	x	-	-	-
<b>Poeciliidae (2)</b>																
<i>Gongopanchax brichardi</i> (Poll, 1971)	x	-	-	-	-	-	-	-	-	x	-	-	-	-	-	x
<i>Hylopanchax</i> sp.	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<b>PERCIFORMES</b>																

TABLE 3. CONTINUED.

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
<b>Channidae (2)</b>																
<i>Parachanna insignis</i> (Sauvage, 1884)	x	-	-	-	x	-	-	-	-	-	-	-	-	-	-	x
<i>Parachanna obscura</i> (Günther, 1861)	x	-	-	x	-	-	-	-	-	-	x	-	-	x	-	x
<b>Anabantidae (9)</b>																
<i>Ctenopoma acutirostre</i> Pellegrin, 1899	x	-	-	-	-	-	-	-	x	-	x	-	x	-	-	x
<i>Ctenopoma gabonense</i> Günther, 1896	-	x	-	-	-	x	-	x	x	-	-	-	x	x	x	-
<i>Ctenopoma kingsleyae</i> Günther, 1896	x	x	-	x	-	x	x	x	-	x	-	-	x	-	-	x
<i>Ctenopoma nigropannosum</i> Reichenow, 1875	-	-	-	-	-	-	-	x	-	-	-	-	-	x	-	x
<i>Ctenopoma ocellatum</i> Pellegrin, 1899	-	-	-	-	-	-	-	-	-	x	-	-	-	-	-	x
<i>Ctenopoma weeksii</i> Boulenger, 1896	x	-	-	-	-	-	x	-	x	x	-	-	-	-	-	x
<i>Microctenopoma ansorgii</i> (Boulenger, 1912)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Microctenopoma fasciolatum</i> (Boulenger, 1899)	x	-	-	-	-	-	-	-	-	-	-	-	-	-	x	x
<i>Microctenopoma nanum</i> (Günther, 1896)	x	-	x	-	-	-	-	-	-	-	-	-	x	-	-	x
<b>Cichlidae (12)</b>																
<i>Congochromis dimidiatus</i> (Pellegrin, 1900)	-	-	x	-	x	-	-	-	-	-	-	-	x	-	-	x
<i>Congochromis</i> sp.	x	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-
<i>Hemichromis elongatus</i> (Guichenot, 1861)	x	x	x	x	x	x	x	-	x	-	-	x	x	x	x	x
<i>Hemichromis lifaliti</i> Loisel, 1979	x	-	-	-	-	-	-	-	x	x	-	-	x	x	-	x
<i>Heterochromis multidentatus</i> (Pellegrin, 1900)	x	-	-	x	-	x	x	-	x	-	-	-	-	-	-	x
<i>Pelmatochromis nigrofasciatus</i> (Pellegrin, 1900)	-	-	x	-	-	-	-	x	-	-	x	-	-	x	-	x
<i>Sarotherodon galilaeus boulengeri</i> (Pellegrin, 1903)	-	-	-	-	-	-	x	-	-	-	-	-	-	-	-	-
<i>Tilapia bilineata</i> Pellegrin, 1900	x	-	-	-	-	x	-	x	x	x	-	-	x	x	x	x
<i>Tilapia congica</i> Poll and Thys van den Audenaerde, 1960	-	-	-	x	-	-	-	-	-	x	-	-	-	-	-	x
<i>Tilapia tholloni</i> (Sauvage, 1884)	-	-	-	x	-	-	-	x	-	x	-	-	x	x	-	-
<i>Tilapia</i> sp.	x	-	-	x	-	-	x	-	-	-	-	-	-	-	-	-
<i>Tylochromis aristoma</i> Stiassny, 1989	x	-	-	-	-	-	-	x	x	x	-	-	-	-	x	x
<b>Eleotridae (2)</b>																
<i>Kribia nana</i> (Boulenger, 1901)	-	-	x	-	-	-	-	-	-	-	-	-	-	-	-	x
<i>Kribia uellensis</i> (Boulenger, 1913)	x	-	x	-	-	-	-	-	-	-	-	-	-	-	-	-

**ACKNOWLEDGMENTS:** Financial support for this study was provided by WWF/DRC and the AMNH Axelrod Research Curatorship. For logistical support we are grateful to L. Steel and I. Omari (WWF/DRC), and to M. Ipantwa (ICCN), and K. Mulumba (Direction des Pêches) for assistance with permits. The taxonomic input of J. van der Zee (Africa Museum, Tervuren), U. Schlieven (ZSM, Munich), J. Sullivan and J. Friel (Cornell University) (Cyprinodontiformes, Cichlidae, Claroteidae, and Mochokidae respectively) is gratefully acknowledged. Our thanks also to R. Schelly (AMNH), who accompanied the senior author on a trip to the SNP and helped with the initial identification of specimens, and to Daniel Ramos (AMNH) for his patience and help with the many aspects of this project. We are grateful to B. Brown and R. Arrindell (AMNH) for the accessioning and cataloging of all specimens.

# LITERATURE CITED

- AFS/AIFRB/ASIH. 2003. *Guidelines for the use of fishes in research*. Document accessible at <http://www.asih.org/files/fish%20guidelines.doc>. Captured on July 27, 2012.
- Brooks, E.G.E., D.J. Allen and W.R.T. Darwell. 2011. *The Status and Distribution of Freshwater Biodiversity in Central Africa*. Gland and Cambridge: IUCN. 126 p.
- Campbell, D. 2005. The Congo River basin; p. 149-166 *In* Fraser, L.H. and P.A. Keddy (ed.). *The World's Largest Wetlands: Ecology and Conservation*. New York: Cambridge University Press.
- Chapman, L.J. 2001. Fishes of African rain forests. Diverse adaptations to environmental challenge; p. 263-290. *In* Weber, W., L.J.T. White, A. Vedder and L. Naughton-Treves (ed.). *African Rain Forest Ecology and Conservation: An Interdisciplinary Perspective*. New Haven: Yale University Press.
- Darwell, W.R.T. and K.G. Smith. 2011. Assessment methodology; p. 21-26 *In* Brooks, E.G.E., D.J. Allen and W.R.T. Darwell (ed.). *The Status and Distribution of Freshwater Biodiversity in Central Africa*. Gland and Cambridge: IUCN.

- Eschmeyer, W.N. 2012. *Catalog of Fishes*. Electronic database accessible at <http://research.calacademy.org/ichthyology/catalog/fishcatmain.asp> Captured 15 December 2012.
- Evrard, C. 1968. *Recherches écologiques sur le peuplement forestier des sols hydromorphes de la cuvette central congolaise*. Série scientifique 110. Laboratoire d'Écologie Végétale de l'Université Catholique de Louvain, Belgium. 295 p.
- Hughes, R.H. and J.S. Hughes, 1992. *A directory of African wetlands*. Gland, Nairobi, and Cambridge: IUCN, UNEP, and WCMC. 820 p.
- Inogwabini, B.I. 2005. Fishes of the Salonga National Park, Democratic Republic of Congo; survey and conservation issues. *Oryx* 39: 78-81.
- Lang, M.A. and C.C. Baldwin 1996. *Methods and techniques of underwater research*. Washington: Proceedings of the American Academy of Underwater Sciences Sixteenth Annual Scientific Diving Symposium, Smithsonian Institution. 236 p.
- Matthes, H. 1964. Les Poissons du Lac Tumba et de la région d'Ikela. *Annales Musée royal de l'Afrique Centrale* 126: 1-204.
- Stiassny, M.L.J., R.E. Brummett, I.J. Harrison, R. Monsembula and V. Mamonekene. 2011. The status and distribution of freshwater fishes in central Africa; p. 27-46 *In* Brooks, E.G.E., D.J. Allen and W.R.T. Darwell (ed.). *The Status and Distribution of Freshwater Biodiversity in Central Africa*. Gland and Cambridge: IUCN.
- UNESCO, 2012. *UNESCO World Heritage List, 36<sup>th</sup> Session*. Electronic Database accessible at <http://whc.unesco.org/en/list/280/>. Captured on December 15, 2012.

RECEIVED: December 2012

ACCEPTED: February 2013

PUBLISHED ONLINE: April 2013

EDITORIAL RESPONSIBILITY: Pedro Hollanda Carvalho