



# New records of the Kinkajou, *Potos flavus* (Schreber, 1774) (Mammalia, Carnivora) in the Cerrado

Jefferson Eduardo Silveira Miranda,<sup>1</sup> Fabiano Rodrigues de Melo,<sup>1,2,3,4,5</sup> Marlucci Baldo Fachi,<sup>1</sup> Seixas Rezende Oliveira,<sup>6</sup> Ricardo Keichi Umetsu<sup>1</sup>

**1** State University of Mato Grosso, Nova Xavantina, Postgraduate Program in Ecology and Conservation, BR 158 - km 65, MT, Brazil 78690000.

**2** Associate Professor II, Institute of Biosciences, Federal University of Goiás, Regional Jataí, Jataí, GO, Brazil 75801615. **3** Visiting Professor, Viçosa Federal University, Department of Forest Engineering, Avenida Purdue, s/nº, Campus Universitário, Edif. Reinaldo de Jesus Araújo, Viçosa, MG, Brazil 36570000. **4** Muriqui Institute for Biodiversity, Euclides Etienne Arreguy Filho, 102/Apt 303 Centro, Caratinga, MG, Brazil 35-300372.

**5** Brazilian Coordinator of the Primate Specialist Group, Species Survival Commission, International Union for Conservation of Nature (PSG/SSC/IUCN). **6** Postgraduate Program in Biodiversity and Conservation, Goiano Federal Institute, Rio Verde, Campus Rio Verde, Rodovia Sul Goiana km 01, s/n, Zona rural, Rio Verde, GO, Brazil, 75906-844.

**Corresponding author:** Fabiano Rodrigues de Melo, [fabiano\\_melo@ufg.br](mailto:fabiano_melo@ufg.br)

## Abstract

We provide new records of *Potos flavus* from 2 localities in the states of Goiás and Mato Grosso, central Brazil. These records extend the geographical distribution of this species in the Cerrado biome and highlights the importance of future research on this species. These new distributional data may contribute to a re-evaluation of its conservation status.

## Key words

Goiás; Central Plateau; Cerrado; central Brazil; roadkill; Procyonidae.

**Academic editor:** Guilherme Garbino | Received 15 November 2017 | Accepted 15 January 2018 | Published 23 March 2018


**Citation:** Miranda JES, Melo FR, Fachi MB, Oliveira SR, Umetsu RK (2018) New records of the kinkajou, *Potos flavus* (Schreber, 1774) (Mammalia, Carnivora) in the Brazilian Savanna. Check List 14 (2): 357–361. <https://doi.org/10.15560/14.2.357>

## Introduction

The Kinkajou, *Potos flavus* (Schreber 1774) (Mammalia, Carnivora, Procyonidae), lives in the tropical rainforests and dry forests with closed canopy. It is a small procyonid having nocturnal and arboreal habits (Kays 1999, Sampaio et al. 2013). *Potos flavus* has rounded ears, a long and flexible tongue, prehensile tail, and woolly fur (Ford and Hoffmann 1988). Pelage coloration is predominantly reddish but some individuals may have a sagittal grey band along the back (Ford and Hoffmann 1988). It is a frugivore/omnivore (Emmons and Feer 1997, Paglia et al. 2012), because most of its diet consists of fruit, but

this is complemented by seeds, flowers, honey, and some insects (Ford and Hoffmann 1988, Kays 1999, Kays et al. 2008). Reproduction occurs at intervals of 1 or 2 years and females give birth to 1 offspring (Kays and Gittleman 2001, Wilson and Reeder 2005, Wilson and Mittermeier 2009, Sampaio et al. 2013).

Typical of tropical environments (Ford and Hoffmann 1988), this species occurs in forested areas in the Americas, from southern Mexico to the Atlantic Forest in southeastern Brazil (Ford and Hoffmann 1988, Kays et al. 2008, Helgen et al. 2016). It is apparently abundant in the Amazon, with its occurrences in anthropized areas (Sampaio et al. 2013). Gonzaga and Rajão (2002) compiled



**Figure 1.** Occurrences of *Potos flavus* in central Brazil. Localities 1–3 represent new or unpublished records reported herein. Localities 4–10 represent previously known occurrences. See Table 1 for locality names, geographic coordinates, and sources of the data.

records of *P. flavus* in the Atlantic Forest for the Brazilian states of Alagoas, Bahia, Espírito Santo, Minas Gerais, and Rio de Janeiro. Because there are a few historical records of *P. flavus* for the Cerrado of Mato Grosso (Cope 1989, Ford and Hoffmann 1998), this species was determined to be Data Deficient for the Cerrado (Sampaio et al. 2013).

The occurrence of *P. flavus* in the Cerrado is based only on historical records, some of them very imprecise, as shown by Vieira (1952). To make clearer the distribution of *P. flavus* in the Cerrado, we present new and historical records for the species in this biome.

## Methods

Our new record of *P. flavus* was made during a wildlife roadkill program in southeastern Goiás in March 2014. We add unpublished records of this species from the follow-

ing sources: the collection of the Ornithological Museum Foundation of Goiânia, and the Collection of Mammals of the Mato Grosso State University (UNEMAT).


The region where our roadkill specimen was found is classified as Aw of Köppen (Silva et al. 2008), with a rainy season (October to April) and a dry season (May to September) (Silva et al. 2006).

## Results

**New records** (Fig. 1). Goiás: near Mineiros city: roadkill, on the BR-364 highway (17°35'34" S, 052°28'59" W), coll. Jefferson E. S. Miranda, Fabiano R. Melo and Seixas R. Oliveira, March 2014 (1 individual, not vouchered). The body was badly damaged, but with intact paws and tail, which helped us with the identification (Fig. 2). The surroundings of the record area are formed

**Table 1.** Occurrences of *Potos flavus* in some regions of Brazil. The occurrence number corresponds to the numbered dots in Figure 1. Datum for geographic coordinates: WGS84.

No.	Locality	State	Geographic coordinates		Source
			Longitude	Latitude	
1	BR 364, Mineiros	Goiás	52°28'59" W	017°35'34" S	New record
2	BR 158, Aragarças	Goiás	52°14'30" W	015°54'09" S	Historical record
3	Nova Xavantina	Mato Grosso	52°20'41" W	014°41'24" S	Historical record
4	Ribeirão Cascalheira	Mato Grosso	51°46'00" W	012°49'00" S	Pine et al. 1970
5	Fazenda Tanguro, Querência	Mato Grosso	52°22'00" W	012°54'00" S	Oliveira et al. 2010
6	Parque Estadual Cristalino	Mato Grosso	55°54'41" W	009°31'38" S	Rocha et al. 2012
7	RPPN "Dr. Marcos Vidigal Vasconcelos", Tombos	Minas Gerais	42°04'19" W	020°53'44" S	Melo et al. 2005
8	Além Paraíba	Minas Gerais	42°43'38" W	021°49'46" S	Gonzaga and Rajão, 2002
9	Passos	Minas Gerais	45°37'00" W	020°43'00" S	Gonzaga and Rajão, 2002
10	Chapada dos Guimarães	Mato Grosso	55°40'28" W	015°28'04" S	Smith, 1885
11	Cocalinho	Mato Grosso	51°40'30" W	014°04'13" S	Muséum d'histoire naturelle de la Ville de Genève
12	Anápolis	Goiás	48°56'35" W	016°20'09" S	Gilmore 1936



**Figure 3.** Habitats and land use at the site of the new record (BR-364 highway, Goiás, Brazil).

by pasture, agriculture, and a seemingly well-preserved vegetation (Fig. 3).

**Historical records.** Goiás: Aragarças, coll. José Hidasi, 1953 (1 skin, Ornithological Museum Foundation, in the city of Goiânia 1120) (J. Hidasi pers. comm.). Mato Grosso: Nova Xavantina: roadkill, on the BR-158 near the urban area of the city ( $14^{\circ}41'24''$  S,  $052^{\circ}20'41''$  W), coll. P. Matos, April 2015 (1 skin, female, UNEMAT CM679).

**Identification.** The road-killed specimen was identified as *Potos flavus* from its predominantly reddish pelage coloration, including muzzle and legs. However, one should not only consider the pelage color alone when differentiating *Potos* from other procyonid genera (Sampaio et al. 2011). In addition, it had a prehensile tail, a distinct trait that allowed us to differentiate between *Potos flavus* and *Bassaricyon alleni* Thomas, 1880, a similar species that also occurs in Brazil and whose morphological characteristics usually lead to misidentifications in the field (Cheida et al. 2011, Sampaio et al. 2011, Sampaio et al. 2013).

## Discussion

Gonzaga and Rajão (2002) suggested that the limits of occurrence of *P. flavus* in the Atlantic Forest might be extended to the north and south with further studies. Melo et al. (2005) corroborated this hypothesis with records in



**Figure 2.** *Potos flavus*. Roadkill on the BR-364 highway ( $17^{\circ}35'34''$  S,  $052^{\circ}28'59''$  W), Goiás, Brazil, March 2014.

the Amazon and Atlantic Forest and that the central areas were important for the evolution of mammals; for some species, connections between the main biomes may have been more recent (Costa 2003).

The distribution of *P. flavus* described by Sampaio et al. (2013) differs from the IUCN map (Kays et al. 2008, Helgen et al. 2016), which does not place this species within the Cerrado. This is probably because at that time there were no recent records of *P. flavus* in this biome. In this sense, there are 2 points to be considered.

First, there are historical records from transition areas between the Cerrado and other biomes, such as

the records from Cocalinho and Chapada dos Guimarães (both in Mato Grosso), as well as 5 records from Anápolis (Goiás) in 1937. The record of Cocalinho, (Fig. 1, point 11), collected in 1774, is based on a skin deposited in the Muséum d'histoire naturelle de la Ville de Genève (Geneva, Switzerland). From Chapada dos Guimarães (Fig. 1, point 10), there are 2 specimens in the American Museum of Natural History (New York, USA), both collected in 1885, 1 male and 1 of an unidentified sex. The 5 records from Anápolis are based on specimens collected in 1937 deposited in the American Museum of Natural History (Fig. 1, point 12). These records should be considered in future assessments of the conservation status of *Potos flavus* in the Cerrado.

Second, our new records should be used to reassess the conservation status and distribution of *P. flavus*, as these records confirm the continued existence of this species in the Cerrado. Thus, a re-evaluation of the conservation status of *P. flavus* in the Cerrado might determine that this species in this biome may belong to one of the Red List categories for threatened species (IUCN 2012). In addition, the known distribution map of the species must be changed to include this biome as the current occurrence area.

Thus, our study confirms the continued existence of *P. flavus* in the Cerrado, indicating that despite being a region occupied by monoculture, we can still find areas in good state of preservation, which need to be further explored. In addition, this study also reinforces the hypothesis that deciduous forests in the interior of the country were and still are characterized as important connections from the recent past that greatly influenced the processes of speciation of the fauna present in our tropical forests (Willis 1992, Costa 2003).

Finally, we suggest that surveys for records of the species must be carried out, mainly in the region of the basins of the Tocantins and Araguaia rivers, which it seems that this region of the Cerrado includes the most records of *P. flavus* that it was found in the literature until now. However, searching the species in other forested areas of the Cerrado is important to enhance its knowledge and to evaluate how many populations there are and point out their sizes. Only after additional study will it be possible to reassess the conservation status of *P. flavus* and, consequently, to prioritize steps for its conservation in the Cerrado.

## Acknowledgements

We thank Dr José Hidasi, Dr Kleber Pereira (Federal University of Goiás, UFG), Dr Fabiana C.S.A. Melo (UFG), Diego A. Silva, and many other students from the UFG that helped us. We also thank FAPEG for financial support (process 201210267001062, FAPEG Edital no. 005/12) and the FGG for providing logistical support, in name of Alessandro Martins (formal director of Regional Jataí). Ricardo Sampaio, José Serrano, and Guilherme Garbino provided very helpful comments and suggestions.

## Authors' Contributions

JESM and FRM collected the data, MBF drew the map. JESM, FRM, MBF, SRO and RKU wrote the text.

## References

- Cheida CC, Nakano-Oliveira EC, Fusco-Costa R, Rocha-Mendes F, Quadros J (2011) Ordem Carnívora. In: Reis NR, Peracchi AL, Pedro WA, Lima IP (Eds) Mamíferos do Brasil. 2nd ed. Edição do Autor, Londrina, 233–286.
- Costa C (2003) The historical bridge between the Amazon and the Atlantic Forest of Brazil: a study of molecular phylogeography with small mammals. *Journal of Biogeography* 30 (1): 71–86. <https://doi.org/10.1046/j.1365-2699.2003.00792.x>
- Emmons L, Feer F (1997) Neotropical Rainforest Mammals: a Field Guide. 2nd ed. University of Chicago Press, Chicago, 396 pp.
- Ford LS, Hoffmann RS (1988) *Potos flavus*. *Mammalian Species* 321:1–9.
- Gilmore. 1936. Mammals Collection, M-134017; American Museum of Natural History Library. <http://entheros.amnh.org/db/emuwebamnh/index.php>, Accessed on: 2017-01-09.
- Gonzaga LP, Rajão H (2002) Distribution of the kinkajou *Potos flavus* (Procyonidae, Carnivora) in the Atlantic forest region of eastern Brazil. *Mammalia* 66: 123–12
- Helgen K, Kays R, Schipper J (2016) *Potos flavus*. The IUCN Red List of Threatened Species 2016: e.T41679A45215631. <https://doi.org/10.2305/IUCN.UK.2016-1.RLTS.T41679A45215631.en>
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. Second edition. Gland, Switzerland and Cambridge, UK: IUCN. iv + 32pp.
- Johnson MA, Saraiva PM, Coelho D (1999) The role of gallery forests in the distribution of Cerrado mammals. *Revista Brasileira de Biologia* 59 (3): 421–427. <http://doi.org/10.1590/S0034-71081999000300006>
- Kays RW (1999) Food preferences of kinkajous (*Potos flavus*): a frugivorous carnivore. *Journal of mammalogy* 80(2): 589–599. <https://doi.org/10.2307/1383303>
- Kays R, Gittleman JL (2001) The social organization of the kinkajou *Potos flavus* (Procyonidae). *Journal of Zoology* 253 (4): 491–504.
- Kays R, Reid F, Schipper J, Helgen K (2008) *Potos flavus*. The IUCN Red List of Threatened Species 2008: e.T41679A10507696. <http://doi.org/10.2305/IUCN.UK.2008.RLTS.T41679A10507696.en>
- Melo FR, Barbosa EF, Souza SLF, Ferraz DS, Rodes ER, Souza SM, Faria MB, Nery MS, Cosenza BAP, Lima FS (2005) Redescoberta do jupará, *Potos flavus* Schreber, 1774 (Carnívora: Procyonidae) no Estado de Minas Gerais, Sudeste do Brasil. *Boletim do Museu de Biologia Mello Leitão* 18: 49–57. [http://www.melloleitao.locaweb.com.br/boletim/arquivos/18/Boletim\\_18\\_Artigo2.pdf](http://www.melloleitao.locaweb.com.br/boletim/arquivos/18/Boletim_18_Artigo2.pdf). Accessed on: 2017-01-09.
- Muséum d'histoire naturelle de la Ville de Genève (2017) Mammals housed at MHNG, Geneva. Occurrence dataset. <https://doi.org/10.15468/lwu4fj>. Accessed via GBIF on: 2017-11-28.
- Oliveira ACM, Carvalho Junior O, Santos PGP, Maria SL, Lima RCS, Rossi RV, Saggin PG, Souza Junior AP, Dourado G (2010) Mamíferos não voadores. In: Oliveira ACM, Santos JB, Santos-Costa MC (Eds) Os animais da Tanguro, Mato Grosso: diversidade na zona de transição entre a Floresta Amazônica e o Cerrado. MPEG, UFPA, and IPAM, Belém, 57–63.
- Paglia AP, Fonseca GAB, Rylands AB, Herrmann G, Aguiar LMS, Chiarello AG, Leite YLR, Costa LP, Siciliano S, Kierulff MCM, Mendes SL, Tavares VC, Mittermeier RA, Patton JL (2012) Lista anotada dos Mamíferos do Brasil/Annotated Checklist of Brazilian Mammals. 2nd edition. Occasional Papers in Conservation Biology: 1–76.
- Pine RH, Bishop IR, Jackson RL (1970) Preliminary list of mammals of the Xavantina/Cachimbo Expedition (Central Brazil). *Transactions*

- of the Royal Society of Tropical Medicine and Hygiene 64 (5): 668–670. [https://doi.org/10.1016/0035-9203\(70\)90003-9](https://doi.org/10.1016/0035-9203(70)90003-9)
- Rocha EC, Silva E, Dalponte JC, Giúdice GML (2012). Efeito das atividades de ecoturismo sobre a riqueza e a abundância de espécies de mamíferos de médio e grande porte na região do Cristalino, Mato Grosso, Brasil. *Revista Árvore* 36 (6): 1061–1072. <http://doi.org/10.1590/S0100-67622012000600007>
- Sampaio R, da Silva MNF, Cohn-Haft M (2011) Reassessment of the occurrence of the kinkajou (*Potos flavus* Schreber, 1774) and olingo (*Bassaricyon beddardi* Pocock, 1921) in the northern Brazilian Amazon. *Studies on Neotropical Fauna and Environment* 46 (2): 85–90. <http://doi.org/10.1080/01650521.2011.572678>
- Sampaio R, Beisiegel BM, Pontes ARM (2013) Avaliação do risco de extinção do jupará *Potos flavus* (Schreber, 1774) no Brasil. *Biodiversidade Brasileira* 3 (1): 277–282.
- Silva FAM, Assad ED, Evangelista BA (2008) Caracterização Climática do bioma Cerrado. In: Sano SM, Almeida SP, Ribeiro JF (Eds) Cerrado: ecologia e flora. EMBRAPA-CPAC, Planaltina, 69–87.
- Silva SC, Santana NMP, Pelegrini JC (2006) Caracterização Climática do Estado de Goiás. Secretaria de Indústria e Comércio. Superintendência de Geologia e Mineração, Goiânia, 133 pp.
- Smith HH (1885) Mammals Collection, MO-353; American Museum of Natural History Library. <http://entheros.amnh.org/db/emuweb/amnh/index.php>. Accessed on: 2017-01-09.
- Willis E (1992) Zoogeographical origins of eastern Brazilian birds. *Ornitologia Neotropical* 3: 1–15.
- Wilson DE, Reeder DM (2005) Mammals species of the world: a taxonomic and geographic reference. 3rd edition. The Johns Hopkins University Press, Baltimore, 2142 pp. (2 vols.).
- Wilson DE, Mittermeier RA (2009) Handbook of the Mammals of the World. Vol. 1, Carnivores. Lynx Edicions, Barcelona, 728 pp.
- Vieira CC (1952). Sobre o “Jupará” do nordeste do Brasil (*Potos flavus nocturnus* (Wied.)). *Papéis Avulsos do Departamento de Zoologia, Secretaria da Agricultura, São Paulo* 11 (3): 33–36.