



First capture of *Diclidurus albus* Wied-Neuwied, 1820 (Mammalia, Chiroptera, Emballonuridae) from Nicaragua

Julio Loza^{1,2}, Javier Sunyer¹, Fiona A. Reid³, José Gabriel Martínez-Fonseca^{1,2,4}

1 Universidad Nacional Autónoma de Nicaragua (UNAN), Enel Central 2.5 km S, Villa Fontana, Managua, Nicaragua. **2** Programa de Conservación de Murciélagos de Nicaragua (PCMN), Parque Central ½ c. W, Santa Teresa, Carazo, Nicaragua. **3** Department of Mammalogy, Royal Ontario Museum, 100 Queen's Park, Toronto, Canada. **4** School of Forestry, Northern Arizona University, 200 E Pine Knoll Dr, Flagstaff, AZ 86011, USA.

Corresponding author: José Gabriel Martínez-Fonseca, jmartinezfonseca89@yahoo.es

Abstract

We report on the first capture and photographs of the Northern Ghost Bat, *Diclidurus albus* Wied-Neuwied, 1820, from Nicaragua, based on one individual from the Reserva Silvestre Privada Refugio Bartola, department of Río San Juan, southeastern Nicaragua. We provide basic morphological measurements of the individual as well as a distribution map of previous visual sightings and acoustic recordings of this species from Nicaragua.

Key words

Lowland Wet Forest; Northern Ghost Bat; Refugio Bartola; mammal; Río San Juan.

Academic editor: Annia Rodríguez-San Pedro | Received 14 August 2018 | Accepted 25 October 2018 | Published 9 November 2018

Citation: Loza J, Sunyer J, Reid FA, Martínez-Fonseca JG (2018) First capture of *Diclidurus albus* Wied-Neuwied, 1820 (Mammalia, Chiroptera, Emballonuridae) from Nicaragua. Check List 14 (6): 1021–1025. <https://doi.org/10.15560/14.6.1021>

Introduction

The Northern Ghost Bat, *Diclidurus albus* Wied-Neuwied, 1820, is an uncommon, solitary, medium-sized, aerial insectivorous white bat that is distributed from Nayarit, Mexico to eastern Brazil, from sea level to 1,700 m a.s.l. (Ceballos and Medellín 1988, Moscoso and Tirira 2009, Lim et al. 2016). It flies high in open areas and in the forest canopy and is therefore very seldom-caught using mist nets (Ceballos and Medellín 1988, Reid 2009). It roosts under palm or banana fronds and on tree trunks (McCarthy 1987, Rodríguez-Herrera et al. 2007, Pedrosa Ferreira et al. 2013) and prefers humid habitats such as riparian and tropical rainforests, although it has also been found in human-disturbed areas such as plantations, clearings, and villages (Ceballos and Medellín 1988, Moscoso and Tirira 2009). Two

subspecies are currently recognized for this species (Ceballos and Medellín 1988): *Diclidurus albus albus* Wied-Neuwied, 1820, distributed in the southern portion of its range, and *D. a. virgo* Thomas, 1903, occurring from Ecuador, Colombia, Venezuela, and the island of Trinidad, northwards.

Gray (1844) recorded 1 specimen of *Diclidurus albus*, as *D. freyreisii* Wied-Neuwied, 1821, a junior synonym of *D. albus* (Alston 1879, Carter and Dolan 1978), from “Pueblo Nuevo, Central America” (stored at the British Museum of Natural History, today the Natural History Museum, London, voucher number BMNH 1842.10.30.1; skin and skull; Fig. 1). The locality of this specimen was later changed to “South America” by Dobson (1878), and redefined to “Pueblo Nuevo, Nicaragua” by Alston (1879). Thereafter, Pueblo Nuevo was considered to be



Figure 1. *Diclidurus albus* (BMNH 1842.10.30.1) from “Costa Rica, comunidad de Pueblo Nuevo, cerca del río Terraba” (see text). Photograph by L. Tomsett, copyright Trustees of the Natural History Museum, UK.

in Nicaragua until 2010 (see below) and all Nicaraguan records of this species during this period (i.e., Alston 1879, Allen 1910, Jones et al. 1971, Ojasti and Linares 1971, Baker and Jones 1975, Jones and Owen 1986, Ceballos and Medellín 1988, Martínez Sánchez 2000, Reid 2009) are based on this specimen alone. Medina-Fitoria et al. (2010) and Medina-Fitoria (2013, 2014) later recorded other visual sightings and acoustic recordings of *D. albus* from the 4 Nicaraguan departments of Atlántico Sur, León, Managua, and Rivas.

Medina-Fitoria et al. (2010) also redefined the locality of BMNH 1842.10.30.1 from “Pueblo Nuevo, Nicaragua” to “Costa Rica, comunidad de Pueblo Nuevo, cerca del río Terraba (08°58' N, 083°32' W)” without further argumentation. Herein we justify this action. Gray (1844: 26) reported that BMNH 1842.10.30.1 “was caught by Mr. Barclay, the botanical collector who accompanied the Voyage [H.M.S. *Sulphur*], on the 27th of March 1837, at Pueblo Nuevo in Central America.” However, on March 27 1837, the *Sulphur* and its crew were still in Costa Rica and that very same day resumed sailing from Puerto Cortéz, close to where Pueblo Nuevo, near río Terraba, is located, north along the Costa Rican coast, and did not anchor until April 4, 1837 offshore at Isla El Cardón (12°28' N, 087°12' W), Nicaragua [T. McCarthy, pers. comm. to A. Medina-Fitoria 2010].

Therefore, *Diclidurus albus* has never been collected in Nicaragua and no photographs or voucher specimens have been ever provided from the country (Medina-Fitoria et al. 2010, Medina-Fitoria 2014). Here we report on the first captured and photographed individual of the Northern Ghost Bat, *D. albus*, from Nicaragua.

Methods

The bat was collected and photographed during an opportunistic find in the Reserva Silvestre Privada Refugio Bartola, Departamento de Río San Juan, southeastern

Nicaragua, which lies within one of Central America’s largest expanses of pristine Lowland Wet Forest (Holdridge 1967). The specimen’s identification was verified with the use of the dichotomous keys and descriptions provided by Reid (2009) and Medina-Fitoria (2014). We followed Reid (2009) for the selection of morphological characters, which were measured with the use of standard calipers and a 100 g Pesola spring scale. Locality records of other visual sightings and acoustic recordings of Nicaraguan *D. albus* were obtained from Medina-Fitoria et al. (2010) and Medina-Fitoria (2014).

Results

New record. Nicaragua: Río San Juan: Reserva de la Biosfera Sureste de Nicaragua: Reserva Privada Refugio Bartola (10.97239° N, 084.33947° W; 35 m above sea level, datum WGS84), Julio Loza observer, 6 January 2017 (1 individual; Fig. 2).

We photographed 1 adult female *Diclidurus albus* night roosting (21:15 h) ca 7 m above the ground in one of the corridors of the Refugio Bartola Private Reserve (Fig. 2A). The bat remained still for ca 1.5 hours before it was collected with a hand net and morphometric measurements and photographs were obtained (Fig. 2B).

The description of the individual is as follows: female; head and body length = 76 mm; tail length = 21 mm; hind foot length = 11 mm; ear length = 16 mm; forearm length = 67 mm; and weight = 21 g.

Identification. There are only 2 species of white bats in Nicaragua (Reid 2009, Medina-Fitoria 2014): *Ectophylla alba* (family Phyllostomidae), which is small and has a noseleaf, and *Diclidurus albus* (family Emballonuridae), which is characterized by the absence of a noseleaf, long snout (upper lip exceeds in length the lower lip), large eyes, and short, yellowish and rounded ears, which fits perfectly with the description of our individual (Fig. 2).



Figure 2. Adult female *Diclidurus albus* from Refugio Bartola Private Reserve, department of Río San Juan, Nicaragua. A) Night roosting in the ceiling of the hotel corridors. B) Portrait shot of the same individual. Photos: JL.

Discussion

Although *Diclidurus albus* has been considered a member of the Nicaraguan mastofauna since the 19th century (Alston 1879), and there have been visual sightings and acoustic recordings of the species in the country from the departments of Atlántico Sur (7 June 2007 in Bluefields and 12 June 2013 in Cruz de Río Grande), Managua (12 January 2009 in the city of Managua), León (24 February 2013 in the city of León), and Rivas (23 March 2014 in La Fé), the current record represents the first captured and photographed individual of this species from Nicaragua, as well as a new departmental record for Río San Juan (Fig. 3). However, there remains no museum voucher specimen of *D. albus* from Nicaragua (Medina-Fitoria et al. 2010, Medina-Fitoria 2013, 2014).

Diclidurus albus is a peculiar, characteristic, and rarely encountered bat species listed as a species of Least Concern under the IUCN (International Union for Conservation of Nature) Red List of Threatened Species categories, where no major threats to this widespread species have been reported (Lim et al. 2016). Because of its foraging habits, it is rarely caught using mist nets (Ceballos and Medellín 1988, Reid 2009). In addition, *D. albus* has highly variable calls, which makes it difficult to identify it from other emballonurid bats from acoustic recordings (Jung et al. 2007, Arias-Aguilar et al. 2018). Although this species is expected to occur across Nicaragua, sampling limitations associated with acoustic identification and physical detection may explain why its distribution has been historically underestimated in the country (Medina-Fitoria 2014; see Fig. 3). Further sampling in Nicaragua will likely fill in gaps in the species distribution throughout the country.

We suspect this opportunistic finding in a building could be a consequence of the extreme disturbance of the surrounding forest of Refugio Bartola caused by hurricane Otto in late November 2016 (Brown 2017), resulting in presumable lack of available roosts.

Acknowledgements

We thank Arnulfo Medina-Fitoria for valuable comments on the note and for sharing personal information with Timm McCarthy (R.I.P.). We thank Louise Tomsett from the British Natural History Museum for photographs and additional data on their specimen. We thank Sandra Castrillo and the Refugio Bartola staff for their support during this and several other field expeditions to the area.

Authors' Contributions

JL took the photographs and the morphometric data, FAR identified the specimen, JS and JGMF wrote the text and elaborated the map. All authors read and approved the final manuscript.

References

- Allen JA (1910) Additional mammals from Nicaragua. Bulletin of the American Museum of Natural History 28: 87–115.
- Alston ER (1879–1882) Biología Centrali-Americana. Mammalia. Taylor and Francis, London, England, 220 pp. <https://doi.org/10.5962/bhl.title.730>
- Arias-Aguilar A, Hintze F, Aguiar LMS, Rufray V, Bernard E, Pereira MJR (2018) Who's calling? Acoustic identification of Brazilian bats. Mammal Research 63 (3): 231–253. <https://doi.org/10.1007/s13364-018-0367-z>




Figure 3. Map of Nicaragua showing the localities where *Diclidurus albus* has been recorded. Green localities corresponds to historical records: both visual sightings and acoustic recordings (circles), only acoustic recordings (triangles), and only visual sightings (square). The red star corresponds to the new captured individual recorded herein. The upper left inset corresponds to the general distribution map of the species (Lim et al. 2016).

- Baker RJ, Jones Jr JK (1975) Additional records of bats from Nicaragua, with a revised checklist of Chiroptera. Occasional papers of the Museum Texas Tech University 32: 1–13. <https://doi.org/10.5962/bhl.title.142873>
- Brown DP (2017) Tropical cyclone report: Hurricane Otto. National Hurricane Center. National Oceanic and Atmospheric Administration, Miami, United States, 28 pp.
- Carter DC, Dolan PG (1978) Catalogue of type specimens of Neotropical bats in selected European museums. Special Publications, the Museum, Texas Tech University 15: 1–136. <https://doi.org/10.5962/bhl.title.142635>
- Ceballos G, Medellín RA (1988) *Diclidurus albus*. Mammalian species 316: 1–4. <https://doi.org/10.2307/3504140>
- Dobson GE (1878) Catalogue of the Chiroptera in the Collection of the British Museum. British Museum London: Order of the Trustees, London, England, 567 pp. <https://doi.org/10.5962/bhl.title.55341>
- Gray JE (1844) Mammalia. In Hinds RB (Ed.) The Zoology of the Voyage of HMS Sulphur under the command of Captain Edward Belcher. Hinds RB (Ed.) (Vol. 1). Smith, Elder & Co., London, England, 150 pp. <https://doi.org/10.5962/bhl.title.53764>
- Jones JK, Owen RD (1986) Checklist and bibliography of Nicaraguan Chiroptera. Occasional Papers of the Museum Texas Tech University 106: 1–13.
- Jones JK, Smith JD, Turner RW (1971) Noteworthy records of bats from Nicaragua, with a checklist of the chiropteran fauna of the country. Occasional Papers of the Museum of Natural History 2: 1–35.
- Jung K, Kalko EKV, von Helversen O (2007) Echolocation calls in Central American emballonurid bats: signal design and call frequency alternation. Journal of Zoology 272: 125–137. <https://doi.org/10.1111/j.1469-7998.2006.00250.x>
- Lim B, Miller B, Reid F, Arroyo-Cabral J, Cuarón AD, de Grammont PC (2016) *Diclidurus albus*. The IUCN Red List of Threatened Species 2016: e.T6561A21986615. <https://doi.org/10.2305/IUCN.UK.2016-2.RLTS.T6561A21986615.en>. Accessed on: 2017-5-23.
- Martínez Sánchez JC, Morales Velásquez S, Castañeda Mendoza EA (2000) Lista patrón de los mamíferos de Nicaragua. Alianza para las Áreas Silvestres (ALAS), Managua, Nicaragua, 35 pp.
- McCarthy TJ (1987) Distributional records of bats from the Caribbean lowlands of Belize and adjacent Guatemala and Mexico. Fieldiana Zoology 39: 137–162.

- Medina-Fitoria A (2013) Identificación de sitios y áreas importantes para la conservación de los murciélagos en Nicaragua. Managua: Programa para la conservación de los murciélagos de Nicaragua (PCMN) & Bat Conservation International (BCI), Managua, Nicaragua, 42 pp.
- Medina-Fitoria A (2014) Murciélagos de Nicaragua. Guía de campo. MARENA, Managua, Nicaragua, 279 pp.
- Medina-Fitoria A, Saldaña O, MacCarthy T, Vilchez S (2010) Nuevos reportes y comentarios históricos de Murciélagos (Orden Chiroptera) para la fauna de Nicaragua. Biodiversidad Revista Nicaragüense 2: 93–102.
- Moscoso P, Tirira DG (2009) Nuevos registros y comentarios sobre la distribución del murciélago blanco común *Diclidurus albus* (Chiroptera: Emballonuridae) en Ecuador. Mastozoología Neotropical 16 (1): 233–238.
- Ojasti J, Linares OJ (1971) Adiciones a la fauna de murciélagos de Venezuela con notas sobre las especies de *Diclidurus*. Acta Biológica Venezuelica 7: 421–441.
- Pedrosa Ferreira A, de Carvalho Melo D, Loures-Ribeiro A (2013) *Diclidurus albus* Wied-Neuwied, 1820 (Chiroptera: Emballonuridae): First record of the species in the state of Paraíba, Brazil. Check List 9 (4): 793–796. <https://doi.org/10.15560/9.4.793>
- Reid FA (2009) A Field Guide to the Mammals of Central America and Southeast Mexico. 2nd edition. Oxford University Press, New York, 346 pp. <https://doi.org/10.2193/2010-132>
- Rodríguez-Herrera B, Medellín RA, Timm RM (2007) Murciélagos neotropicales que acampan en hojas. INbio, Santo Domingo de Heredia, Costa Rica, 178 pp.