

# New information on distribution of a marine leech, *Pontobdella muricata* (Linnaeus, 1758), from the Mediterranean coast of Turkey

Saadet Yağmur Bulguroğlu, Jale Korun\* and Mehmet Gökoğlu

Akdeniz University, Faculty of Fisheries, Aquaculture Division, Department of Fish Diseases, Antalya, Turkey, 07058

\* Corresponding author. E-mail: [jalekorun@akdeniz.edu.tr](mailto:jalekorun@akdeniz.edu.tr)

**Abstract:** New information on distribution of marine leech, *Pontobdella muricata* (Linnaeus, 1758) is given here. One leech was observed on the dorsal surface of Thornback Ray, *Raja clavata* (Linnaeus, 1758) that was caught in 2013 in the Antalya, Turkey. This is a new record for the eastern Mediterranean coast of Turkey.

**Key words:** marine leech, *Pontobdella muricata*, Thornback Ray, *Raja clavata*

Marine leeches are included in the class Hirudinea, which is divided into four orders, Acanthobdellida, Rhynchobdellida, Gnathobdellida, and Pharyngobdellida (Govedich et al. 2004). Members of these orders infest elasmobranch fishes. The subfamily Pontobdellinae (order Rhynchobdellida) is characterized by having metameric line (annulus) (Llwellyn 1966), which is a distinctive feature for classification of the genera (Klemm 1982). *Stibarobdella*, *Pontobdella*, and *Pentabdella* contain species that have trimeric, tetrameric, and pentameric somites, respectively (Llwellyn 1966). *Pontobdella* is characterized by the tubercles which occur on the major ( $a_2$ ) annulus of each somite and usually on most annuli of the testicular and caecal regions and by the body circular or slightly flattened in cross sections (Govedich et al. 2004).


Leeches damage the bodies of their hosts. *Pontobdella muricata* (Linnaeus, 1758) causes local hemorrhages, swelling, tissue necrosis, and disappearance on the surface of epidermal tissue on its attachment regions on the host (Sağlam et al. 2003; Bakopoulos and Ksida 2013).

A Thornback Ray, *Raja clavata* (Linnaeus, 1758) was caught in November 2013 by fishermen as a non-target species from Antalya Bay located in the Mediterranean coast of Turkey. The ray was captured at a depth of 70 m between 36°46' N, 031°12' E and 36°46' N, 031°08' E (Figure 1). A leech (Figure 2) was collected on the dorsal surface of the Thornback Ray and then fixed in 70%

ethanol without relaxation in the field. The fixed leech was examined in a Petri dish using forceps and parasitological needles under a stereomicroscope. The length, width and also diameters of oral and caudal suckers of the leech were measured, annulation on body surface of parasite was separated as  $a_1$ ,  $a_2$ ,  $b_5$ , and  $b_6$  annuli, the somites on these annuli were examined in terms of the morphological characteristics and then these were counted for dorsal, lateral and ventral regions. After counting, the parasite was preserved in 70% ethanol for long-term storage. Identification of the leech was performed according to Sawyer (1986) and Llwellyn (1966).

*Pontobdella muricata* presented a dark green colour when was fresh, but the colour changed to pale yellow after fixation processes. The leech had oral and caudal suckers and there was a marginal fringe on the oral sucker and did not present true eyes. The length and width of the parasite were 99 and 28 mm, respectively. The diameters of the oral and caudal suckers were 2 and 5 mm, respectively. The anterior sucker was wider than trachelosome. Complete urosomal somites had four annulate as  $a_1$ ,  $a_2$ ,  $b_5$  and  $b_6$ . Urosomal tubercle numbers were 12 tubercles on  $a_1$ , 8 tubercles on  $a_2$ , 10 tubercles on  $b_5$  and 14 small tubercles on  $b_6$ . Annulation of the leech was as follow; uniannulate somites: none; biannulate somites: XI, XII, XXV, XXVI, XXVII; triannulate somites: VII–X; quadriannulate somites: XII–XXIV. The characteristics of the leech are in agreement with previous descriptions of *Pontobdella muricata* reported by Sawyer (1986) and Llwellyn (1966).

*Pontobdella muricata* was reported on the elasmobranchs such as ray fish, *Raja* spp., marbled electric ray, *Torpedo* spp., common stingray, *Dasyatis pastinaca*, blacktip reef shark *Carcharhinus melanopterus*, and teleost fish species such as flounder, *Pleuronectes* spp. and horse mackerel, *Trachurus trachurus capensis* from many countries including Pakistan, Namibia, USA, Greece, Europe, Norway, Iceland, and Greenland (Piasecki 1982;



**Figure 1.** The map showing the previously records (yellow dots) and new record (green dot) for the species *Pontobdella muricata* in Turkey.



**Figure 2.** *Pontobdella muricata* on dorsal region of *Raja clavata*.

Sawyer 1986; Kazmi and Naushaba 2013; Bakopoulos and Ksidia 2013). In Turkey, *P. muricata* was collected from the elasmobranchs including *Torpedo marmorata* and *Raja clavata* from the Aegean Sea, *Raja* sp. from the Marmara Sea, and *R. clavata* from the Black Sea (Ergüven and Candan 1992; Sağlam *et al.* 2003; Ökter and Utevsky 2010).

*Pontobdella muricata* was previously recorded from the Black, Marmara, Aegean and western Mediterranean coastal waters of Turkey from elasmobranch fishes. This species was not previously reported from the eastern Mediterranean of Turkey and the present study is the first record in this area.

## LITERATURE CITED

- Bakopoulos, V. and V.C. Ksidia. 2013. *Pontobdella muricata* infection of *Raja clavata* and *Dayatis pastinaca* off the coast of Lesvos, Greece. *Journal of the Marine Biological Association of the United Kingdom* 94(2): 405–409. doi: 10.1017/S0025315413000830
- Ergüven, H. and A. Candan 1992. A parasitic Hirudinea (*Pontobdella muricata* Linnaeus) at *Raja* sp. in Marmara Sea. *Turkish Journal of Fisheries and Aquatic Sciences* 2(1): 1–4
- Govedich, F.R, W.E. Moser and R.W. Davies 2004. Annelida: Clitellata, Hirudinea, Euhirudinea; pp. 175–190, in: C.M. Yule and Y.H. Sen (eds.). *Freshwater Invertebrates of the Malaysian Region*. Kuala Lumpur: Academy of Sciences Malaysia.
- Kazmi, Q.B. and R. Naushaba. 2013. Checklist of marine worms reported from Pakistani marine waters. *Pakistan Journal of Nematology* 31(2): 187–280.
- Klemm, D.J. 1982. *Leeches (Annelida: Hirudinea) of North America*. United States Environmental Protection Agency. Cincinnati, Ohio: Research and Development Publications. 117 pp.
- Llewellyn, L.C. 1966. Pontobdellinae (Piscicolidae: Hirudinea) in the British Museum (Natural History) with a review of the subfamily. *Bulletin of the British Museum (Natural History) Zoology* 14: 391–439. <http://www.biodiversitylibrary.org/page/4057312>
- Ökter, A. and S.Y. Utevsky 2010. New information on the hosts and distribution of the marine fish leeches *Trachelobdella lubrica* and *Pontobdella muricata* (Clitellata, Hirudinida). *Vestnik Zoology* 44(4):33–36. doi: 10.2478/v10058-010-0023-9
- Piasecki, W. 1982. Parasitofauna of Cape Horse Mackerel *Trachurus trachurus capensis* Castelnau, 1861. *Parazytologia Ostroboka Kapskiego Trachurus trachurus capensis* Castelnau, 1861. *Acta Ichthyologica et Piscatoria, Szczecin* 12(1): 43–56. [http://www.aipep.pl/volumes/1980/2\\_1/pdf/12\\_1\\_03.pdf](http://www.aipep.pl/volumes/1980/2_1/pdf/12_1_03.pdf).
- Robertson, M. 1909. Further notes on a trypanosome found in the alimentary tract of *Pontobdella muricata*. *Quarterly Journal of Microscopical Science* 54(213): 119–139. <http://jcs.biologists.org/content/s2-54/213/119.full.pdf>.
- Sağlam, N., M.C. Oguz, E.S. Celik, S.A. Doyuk and A. Usta 2003. *Pontobdella muricata* and *Trachelobdella lubrica* (Hirudinea: Piscicolidae) on some marine fish in the Dardanelles, Turkey. *Journal of the Marine Biological Association of the United Kingdom* 83(6): 1315–1316. doi: 10.1017/S0025315403008749
- Sawyer, R. T. 1986. *Leech Biology and Behaviour*. Volumes II. Oxford: Clarendon Press. 1065 pp. doi: 10.1017/S0025315400026801

**Authors' contribution statement:** MG collected the data, JK and SYB wrote the text, and SYB made the analysis.

**Received:** March 2014

**Accepted:** January 2015

**Editorial responsibility:** Simone Chinicz Cohen